

Cuilin Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6931543/publications.pdf>

Version: 2024-02-01

10
papers

174
citations

1163117

8
h-index

1372567

10
g-index

10
all docs

10
docs citations

10
times ranked

321
citing authors

#	ARTICLE	IF	CITATIONS
1	P4HB and PDIA3 are associated with tumor progression and therapeutic outcome of diffuse gliomas. <i>Oncology Reports</i> , 2018, 39, 501-510.	2.6	46
2	Down-regulation of DANCR acts as a potential biomarker for papillary thyroid cancer diagnosis. <i>Bioscience Reports</i> , 2019, 39, .	2.4	34
3	Molecular mechanisms and potential prognostic effects of REST and REST4 in glioma. <i>Molecular Medicine Reports</i> , 2017, 16, 3707-3712.	2.4	16
4	Glucose metabolism-related gene polymorphisms as the risk predictors of type 2 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2020, 12, 97.	2.7	15
5	Effect of AMP-activated protein kinase subunit alpha 2 (PRKAA2) genetic polymorphisms on susceptibility to type 2 diabetes mellitus and diabetic nephropathy in a Chinese population. <i>Journal of Diabetes</i> , 2018, 10, 43-49.	1.8	13
6	SREBP1 as a potential biomarker predicts levothyroxine efficacy of differentiated thyroid cancer. <i>Biomedicine and Pharmacotherapy</i> , 2020, 123, 109791.	5.6	13
7	REST, not REST4, is a risk factor associated with radiotherapy plus chemotherapy efficacy in glioma. <i>Drug Design, Development and Therapy</i> , 2018, Volume 12, 1363-1371.	4.3	12
8	Genetic polymorphism contributes to ¹³¹ I radiotherapy-induced toxicities in patients with differentiated thyroid cancer. <i>Pharmacogenomics</i> , 2018, 19, 1335-1344.	1.3	9
9	STK11 rs2075604 Polymorphism Is Associated with Metformin Efficacy in Chinese Type 2 Diabetes Mellitus. <i>International Journal of Endocrinology</i> , 2017, 2017, 1-6.	1.5	8
10	LncRNAs SNHG12 and LINC00152 were associated with progression of patients with papillary thyroid carcinoma. <i>Future Oncology</i> , 2019, 15, 4167-4179.	2.4	8