Angelika Fruth

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6934787/publications.pdf

Version: 2024-02-01

26 papers 2,623 citations

623574 14 h-index 26 g-index

26 all docs

26 docs citations

26 times ranked 3170 citing authors

#	Article	IF	CITATIONS
1	Zoonotic bacteria in clinically healthy goats in petting zoo settings of zoological gardens in Germany. Zoonoses and Public Health, 2022, , .	0.9	3
2	Influence of <i>Salmonella</i> specific bacteriophages (O1; S16) on the shedding of naturally occurring <i>Salmonella</i> and an orally applied <i>Salmonella</i> Eastbourne strain in bearded dragons (<i>Pogona vitticeps</i>). Veterinary Medicine and Science, 2021, 7, 534-547.	0.6	3
3	The Superior Adherence Phenotype of E. coli O104:H4 is Directly Mediated by the Aggregative Adherence Fimbriae Type I. Virulence, 2021, 12, 346-359.	1.8	9
4	Genome-wide insights into population structure and host specificity of Campylobacter jejuni. Scientific Reports, 2021, 11, 10358.	1.6	18
5	Genetic diversity and pathogenic potential of Shiga toxin-producing Escherichia coli (STEC) derived from German flour. International Journal of Food Microbiology, 2021, 347, 109197.	2.1	7
6	Population structure-guided profiling of antibiotic resistance patterns in clinical <i>Listeria monocytogenes</i> isolates from Germany identifies <i>pbpB3</i> alleles associated with low levels of cephalosporin resistance. Emerging Microbes and Infections, 2020, 9, 1804-1813.	3.0	18
7	Simple differentiation of Salmonella Typhi, Paratyphi and Choleraesuis from Salmonella species using the eazyplex TyphiTyper LAMP assay. Journal of Medical Microbiology, 2020, 69, 817-823.	0.7	2
8	Whole-Genome-Based Public Health Surveillance of Less Common Shiga Toxin-Producing Escherichia coli Serovars and Untypeable Strains Identifies Four Novel O Genotypes. Journal of Clinical Microbiology, 2019, 57, .	1.8	25
9	The use of a salmonella bacteriophage in bearded dragons: application, passage time and reisolation. Tierarztliche Praxis Ausgabe K: Kleintiere - Heimtiere, 2019, 47, 247-256.	0.3	3
10	Novel type of pilus associated with a Shiga-toxigenic <i>E. coli</i> hybrid pathovar conveys aggregative adherence and bacterial virulence. Emerging Microbes and Infections, 2018, 7, 1-16.	3.0	21
11	Attack of the clones: whole genome-based characterization of two closely related enterohemorrhagic Escherichia coli O26 epidemic lineages. BMC Genomics, 2018, 19, 647.	1.2	7
12	Shiga toxin-producing Escherichia coli O103:H2 outbreak in Germany after school trip to Austria due to raw cow milk, 2017 – The important role of international collaboration for outbreak investigations. International Journal of Medical Microbiology, 2018, 308, 539-544.	1.5	35
13	Bacteriophage Sf6 Tailspike Protein for Detection of Shigella flexneri Pathogens. Viruses, 2018, 10, 431.	1.5	25
14	Molecular characterisation of extended-spectrum \hat{l}^2 -lactamase (ESBL)-producing Escherichia coli isolates from hospital and ambulatory patients in Germany. Veterinary Microbiology, 2017, 200, 130-137.	0.8	71
15	Host cell interactions of outer membrane vesicle-associated virulence factors of enterohemorrhagic Escherichia coli O157: Intracellular delivery, trafficking and mechanisms of cell injury. PLoS Pathogens, 2017, 13, e1006159.	2.1	176
16	Clonal spread and interspecies transmission of clinically relevant ESBL-producing ⟨i⟩Escherichia coli⟨ i⟩of ST410â€"another successful pandemic clone?. FEMS Microbiology Ecology, 2016, 92, fiv155.	1.3	120
17	Virulence from vesicles: Novel mechanisms of host cell injury by Escherichia coli O104:H4 outbreak strain. Scientific Reports, 2015, 5, 13252.	1.6	122
18	Extended-spectrum beta-lactamase-producing Shiga toxin gene (stx1)-positive Escherichia coli O91:H14 carrying blaCTX-M-15 on an Incl1-ST31 plasmid isolated from a human patient in Germany. International Journal of Medical Microbiology, 2015, 305, 404-407.	1.5	12

#	Article	IF	CITATIONS
19	Molecular epidemiological view on Shiga toxin-producing Escherichia coli causing human disease in Germany: Diversity, prevalence, and outbreaks. International Journal of Medical Microbiology, 2015, 305, 697-704.	1.5	46
20	Correlation between the genomic o454-nlpD region polymorphisms, virulence gene equipment and phylogenetic group of extraintestinal Escherichia coli (ExPEC) enables pathotyping irrespective of host, disease and source of isolation. Gut Pathogens, 2014, 6, 37.	1.6	6
21	Molecular epidemiology of Salmonella enterica serovar Kottbus isolated in Germany from humans, food and animals. Veterinary Microbiology, 2014, 170, 97-108.	0.8	12
22	A molecular scheme for Yersinia enterocolitica patho-serotyping derived from genome-wide analysis. International Journal of Medical Microbiology, 2014, 304, 275-283.	1.5	42
23	Characterization of Escherichia coli Isolates from Hospital Inpatients or Outpatients with Urinary Tract Infection. Journal of Clinical Microbiology, 2014, 52, 407-418.	1.8	120
24	Two Novel EHEC/EAEC Hybrid Strains Isolated from Human Infections. PLoS ONE, 2014, 9, e95379.	1.1	39
25	Epidemic Profile of Shiga-Toxin–Producing <i>Escherichia coli</i> O104:H4 Outbreak in Germany. New England Journal of Medicine, 2011, 365, 1771-1780.	13.9	1,008
26	Characterisation of the Escherichia coli strain associated with an outbreak of haemolytic uraemic syndrome in Germany, 2011: a microbiological study. Lancet Infectious Diseases, The, 2011, 11, 671-676.	4.6	673