

# Qiuli Wu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6935739/publications.pdf>

Version: 2024-02-01

28  
papers

1,597  
citations

236833

25  
h-index

501076

28  
g-index

28  
all docs

28  
docs citations

28  
times ranked

941  
citing authors

#	ARTICLE	IF	CITATIONS
1	Induction of Protective Response Associated with Expressional Alterations in Neuronal G Protein-Coupled Receptors in Polystyrene Nanoparticle Exposed <i>Caenorhabditis elegans</i> . <i>Chemical Research in Toxicology</i> , 2021, 34, 1308-1318.	1.7	28
2	Neuronal G $\alpha$ subunits required for the control of response to polystyrene nanoparticles in the range of 1/4g/L in <i>C. elegans</i> . <i>Ecotoxicology and Environmental Safety</i> , 2021, 225, 112732.	2.9	27
3	Dysregulation of G protein-coupled receptors in the intestine by nanoplastic exposure in <i>Caenorhabditis elegans</i> . <i>Environmental Science: Nano</i> , 2021, 8, 1019-1028.	2.2	19
4	Response of G protein-coupled receptor CED-1 in germline to polystyrene nanoparticles in <i>Caenorhabditis elegans</i> . <i>Nanoscale Advances</i> , 2021, 3, 1997-2006.	2.2	26
5	Graphene oxide disrupts the protein-protein interaction between Neuroligin/NLG-1 and DLG-1 or MAGI-1 in nematode <i>Caenorhabditis elegans</i> . <i>Science of the Total Environment</i> , 2020, 700, 134492.	3.9	40
6	Lipid metabolic response to polystyrene particles in nematode <i>Caenorhabditis elegans</i> . <i>Environmental Pollution</i> , 2020, 256, 113439.	3.7	69
7	The <i>C. elegans</i> miR-235 regulates the toxicity of graphene oxide via targeting the nuclear hormone receptor DAF-12 in the intestine. <i>Scientific Reports</i> , 2020, 10, 16933.	1.6	4
8	Epigenetic response to nanopolystyrene in germline of nematode <i>Caenorhabditis elegans</i> . <i>Ecotoxicology and Environmental Safety</i> , 2020, 206, 111404.	2.9	38
9	Arsenite-induced transgenerational glycometabolism is associated with up-regulation of H3K4me2 via inhibiting spr-5 in <i>caenorhabditis elegans</i> . <i>Toxicology Letters</i> , 2020, 326, 11-17.	0.4	43
10	Response of intestinal G $\alpha$ subunits to nanopolystyrene in nematode <i>Caenorhabditis elegans</i> . <i>Environmental Science: Nano</i> , 2020, 7, 2351-2359.	2.2	26
11	A circular RNA <i>circ_0000115</i> in response to graphene oxide in nematodes. <i>RSC Advances</i> , 2019, 9, 13722-13735.	1.7	31
12	Dysregulation of let-7 by PEG modified graphene oxide in nematodes with deficit in epidermal barrier. <i>Ecotoxicology and Environmental Safety</i> , 2019, 169, 1-7.	2.9	30
13	NPR-9 regulates the innate immune response in <i>Caenorhabditis elegans</i> by antagonizing the activity of AIB interneurons. <i>Cellular and Molecular Immunology</i> , 2018, 15, 27-37.	4.8	63
14	Toxicity of Graphene Oxide in Nematodes with a Deficit in the Epidermal Barrier Caused by RNA Interference Knockdown of <i>unc-52</i> . <i>Environmental Science and Technology Letters</i> , 2018, 5, 622-628.	3.9	29
15	Coal combustion related fine particulate matter (PM <sub>2.5</sub> ) induces toxicity in <i>Caenorhabditis elegans</i> by dysregulating microRNA expression. <i>Toxicology Research</i> , 2017, 6, 432-441.	0.9	38
16	Neuronal ERK signaling in response to graphene oxide in nematode <i>Caenorhabditis elegans</i> . <i>Nanotoxicology</i> , 2017, 11, 520-533.	1.6	55
17	Function of RSKS-1-AAK-2-DAF-16 signaling cascade in enhancing toxicity of multi-walled carbon nanotubes can be suppressed by mir-259 activation in <i>Caenorhabditis elegans</i> . <i>Scientific Reports</i> , 2016, 6, 32409.	1.6	50
18	An epigenetic signal encoded protection mechanism is activated by graphene oxide to inhibit its induced reproductive toxicity in <i>Caenorhabditis elegans</i> . <i>Biomaterials</i> , 2016, 79, 15-24.	5.7	111

#	ARTICLE	IF	CITATIONS
19	p38 MAPK-SKN-1/Nrf signaling cascade is required for intestinal barrier against graphene oxide toxicity in <i>Caenorhabditis elegans</i> . <i>Nanotoxicology</i> , 2016, 10, 1469-1479.	1.6	73
20	Genome-wide identification and functional analysis of long noncoding RNAs involved in the response to graphene oxide. <i>Biomaterials</i> , 2016, 102, 277-291.	5.7	85
21	Quantum dots increased fat storage in intestine of <i>Caenorhabditis elegans</i> by influencing molecular basis for fatty acid metabolism. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2016, 12, 1175-1184.	1.7	48
22	Crucial role of intestinal barrier in the formation of transgenerational toxicity in quantum dot exposed nematodes <i>Caenorhabditis elegans</i> . <i>RSC Advances</i> , 2015, 5, 94257-94266.	1.7	40
23	A microRNAs-mRNAs network involved in the control of graphene oxide toxicity in <i>Caenorhabditis elegans</i> . <i>RSC Advances</i> , 2015, 5, 92394-92405.	1.7	40
24	Immune response is required for the control of in vivo translocation and chronic toxicity of graphene oxide. <i>Nanoscale</i> , 2014, 6, 5894.	2.8	115
25	Response of MicroRNAs to <i>In Vitro</i> Treatment with Graphene Oxide. <i>ACS Nano</i> , 2014, 8, 2100-2110.	7.3	91
26	microRNAs control of in vivo toxicity from graphene oxide in <i>Caenorhabditis elegans</i> . <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2014, 10, 1401-1410.	1.7	70
27	Contributions of altered permeability of intestinal barrier and defecation behavior to toxicity formation from graphene oxide in nematode <i>Caenorhabditis elegans</i> . <i>Nanoscale</i> , 2013, 5, 9934.	2.8	170
28	Translocation, transfer, and in vivo safety evaluation of engineered nanomaterials in the non-mammalian alternative toxicity assay model of nematode <i>Caenorhabditis elegans</i> . <i>RSC Advances</i> , 2013, 3, 5741.	1.7	138