

# Nusrat Husain

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6937757/publications.pdf>

Version: 2024-02-01

139  
papers

3,750  
citations

117453

34  
h-index

161609

54  
g-index

140  
all docs

140  
docs citations

140  
times ranked

4207  
citing authors

#	ARTICLE	IF	CITATIONS
1	Minocycline benefits negative symptoms in early schizophrenia: a randomised double-blind placebo-controlled clinical trial in patients on standard treatment. <i>Journal of Psychopharmacology</i> , 2012, 26, 1185-1193.	2.0	247
2	Prevalence and social correlates of postnatal depression in a low income country. <i>Archives of Women's Mental Health</i> , 2006, 9, 197-202.	1.2	153
3	Depression and social stress in Pakistan. <i>Psychological Medicine</i> , 2000, 30, 395-402.	2.7	144
4	The current status of culturally adapted mental health interventions: a practice-focused review of meta-analyses. <i>Neuropsychiatric Disease and Treatment</i> , 2018, Volume 14, 165-178.	1.0	142
5	The benefit of minocycline on negative symptoms of schizophrenia in patients with recent-onset psychosis (BeneMin): a randomised, double-blind, placebo-controlled trial. <i>Lancet Psychiatry</i> , 2018, 5, 885-894.	3.7	133
6	Minocycline and celecoxib as adjunctive treatments for bipolar depression: a multicentre, factorial design randomised controlled trial. <i>Lancet Psychiatry</i> , 2020, 7, 515-527.	3.7	110
7	Minocycline as an adjunct for treatment-resistant depressive symptoms: A pilot randomised placebo-controlled trial. <i>Journal of Psychopharmacology</i> , 2017, 31, 1166-1175.	2.0	104
8	Multidimensional scale of perceived social support: Psychometric properties in a South Asian population. <i>Journal of Obstetrics and Gynaecology Research</i> , 2010, 36, 845-851.	0.6	91
9	Cluster randomized trial of a parent-based intervention to support early development of children in a low-income country. <i>Child: Care, Health and Development</i> , 2009, 35, 56-62.	0.8	78
10	Social stress and depression during pregnancy and in the postnatal period in British Pakistani mothers: A cohort study. <i>Journal of Affective Disorders</i> , 2012, 140, 268-276.	2.0	77
11	Brief culturally adapted CBT for psychosis (CaCBTp): A randomized controlled trial from a low income country. <i>Schizophrenia Research</i> , 2015, 164, 143-148.	1.1	75
12	An examination of the relationship between shame, guilt and self-harm: A systematic review and meta-analysis. <i>Clinical Psychology Review</i> , 2019, 73, 101779.	6.0	71
13	Adverse social circumstances and depression in people of Pakistani origin in the UK. <i>British Journal of Psychiatry</i> , 1997, 171, 434-438.	1.7	68
14	Brief psychological intervention after self-harm: randomised controlled trial from Pakistan. <i>British Journal of Psychiatry</i> , 2014, 204, 462-470.	1.7	67
15	Effect of delaying treatment of first-episode psychosis on symptoms and social outcomes: a longitudinal analysis and modelling study. <i>Lancet Psychiatry</i> , 2020, 7, 602-610.	3.7	66
16	Self-harm in the UK. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2006, 41, 782-788.	1.6	65
17	Ethnicity and psychopharmacology. <i>Journal of Psychopharmacology</i> , 2008, 22, 673-680.	2.0	63
18	A population-based study of irritable bowel syndrome in a non-Western population. <i>Neurogastroenterology and Motility</i> , 2008, 20, 1022-1029.	1.6	62

#	ARTICLE	IF	CITATIONS
19	Brief Culturally adapted CBT (CaCBT) for depression: A randomized controlled trial from Pakistan. <i>Journal of Affective Disorders</i> , 2015, 177, 101-107.	2.0	62
20	A Systematic Review of the Evidence Supporting Mobile and Internet Based Psychological Interventions For Self-Harm. <i>Suicide and Life-Threatening Behavior</i> , 2020, 50, 151-179.	0.9	61
21	Life stress and depression in a tribal area of Pakistan. <i>British Journal of Psychiatry</i> , 2007, 190, 36-41.	1.7	56
22	Social intervention for British Pakistani women with depression: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2010, 197, 227-233.	1.7	55
23	Prevalence and psychosocial correlates of perinatal depression: a cohort study from urban Pakistan. <i>Archives of Women's Mental Health</i> , 2011, 14, 395-403.	1.2	55
24	Barefoot therapists: barriers and facilitators to delivering maternal mental health care through peer volunteers in Pakistan: a qualitative study. <i>International Journal of Mental Health Systems</i> , 2016, 10, 24.	1.1	53
25	Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. <i>Lancet Psychiatry</i> , 2020, 7, 788-800.	3.7	51
26	A naturalistic, randomized, controlled trial combining cognitive remediation with cognitive behavioural therapy after first-episode non-affective psychosis. <i>Psychological Medicine</i> , 2014, 44, 1889-1899.	2.7	49
27	A multicentre randomised controlled trial of a carer supervised Culturally adapted CBT (CaCBT) based self-help for depression in Pakistan. <i>Journal of Affective Disorders</i> , 2014, 156, 224-227.	2.0	47
28	A systematic review of explanatory models of illness for psychosis in developing countries. <i>International Review of Psychiatry</i> , 2012, 24, 450-462.	1.4	45
29	Social factors associated with chronic depression among a population-based sample of women in rural Pakistan. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2004, 39, 618-24.	1.6	44
30	An Exploratory Randomized Controlled Study of a Healthy Living Intervention in Early Intervention Services for Psychosis. <i>Journal of Clinical Psychiatry</i> , 2014, 75, 498-505.	1.1	44
31	Self-harm in British South Asian women: psychosocial correlates and strategies for prevention. <i>Annals of General Psychiatry</i> , 2006, 5, 7.	1.2	42
32	Medically unexplained presentations and quality of life: A study of a predominantly South Asian primary care population in England. <i>Journal of Psychosomatic Research</i> , 2008, 65, 311-317.	1.2	39
33	A Qualitative Study to Explore Patients, Carers and Health Professionals Views to Culturally Adapt CBT for Psychosis (CBTp) in Pakistan. <i>Behavioural and Cognitive Psychotherapy</i> , 2016, 44, 43-55.	0.9	39
34	Impact of early intervention services on duration of untreated psychosis: Data from the National EDEN prospective cohort study. <i>Schizophrenia Research</i> , 2014, 159, 1-6.	1.1	38
35	mHealth based interventions for the assessment and treatment of psychotic disorders: a systematic review. <i>MHealth</i> , 2017, 3, 33-33.	0.9	33
36	Treatment of maternal depression in urban slums of Karachi, Pakistan: A randomized controlled trial (RCT) of an integrated maternal psychological and early child development intervention. <i>Asian Journal of Psychiatry</i> , 2017, 29, 63-70.	0.9	32

#	ARTICLE	IF	CITATIONS
37	A qualitative study to explore views of patients <sup>â€™</sup> , carers <sup>â€™</sup> and mental health professionals <sup>â€™</sup> to inform cultural adaptation of CBT for psychosis (CBTp) in China. <i>BMC Psychiatry</i> , 2017, 17, 131.	1.1	31
38	Minocycline as an adjunct for treatment-resistant depressive symptoms: study protocol for a pilot randomised controlled trial. <i>Trials</i> , 2015, 16, 410.	0.7	30
39	Increasing access to Cognitive Behaviour Therapy in Low and Middle Income Countries: A strategic framework. <i>Asian Journal of Psychiatry</i> , 2016, 22, 190-195.	0.9	28
40	Deconstructing depression and negative symptoms of schizophrenia; differential and longitudinal immune correlates, and response to minocycline treatment. <i>Brain, Behavior, and Immunity</i> , 2021, 91, 498-504.	2.0	28
41	Comparison of the Personal Health Questionnaire and the Self Reporting Questionnaire in rural Pakistan. <i>JPMA the Journal of the Pakistan Medical Association</i> , 2006, 56, 366-70.	0.1	26
42	Detecting Depression in Pregnancy: Validation of EPDS in British Pakistani Mothers. <i>Journal of Immigrant and Minority Health</i> , 2014, 16, 1085-1092.	0.8	25
43	Antidepressant and Group Psychosocial Treatment for Depression: A Rater Blind Exploratory RCT from a Low Income Country. <i>Behavioural and Cognitive Psychotherapy</i> , 2014, 42, 693-705.	0.9	23
44	Minocycline and celecoxib as adjunctive treatments for bipolar depression: a study protocol for a multicenter factorial design randomized controlled trial. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 1-8.	1.0	23
45	Group psychological intervention for postnatal depression: a nested qualitative study with British South Asian women. <i>BMC Women's Health</i> , 2015, 15, 109.	0.8	22
46	An evaluation of subjective experiences, effects and overall satisfaction with clozapine treatment in a UK forensic service. <i>Therapeutic Advances in Psychopharmacology</i> , 2015, 5, 146-150.	1.2	22
47	Group eye movement desensitization and reprocessing interventions in adults and children: A systematic review of randomized and nonrandomized trials. <i>Clinical Psychology and Psychotherapy</i> , 2021, 28, 784-806.	1.4	22
48	Psychotherapy or medication for depression? Using individual symptom meta-analyses to derive a Symptom-Oriented Therapy (SOt) metric for a personalised psychiatry. <i>BMC Medicine</i> , 2020, 18, 170.	2.3	21
49	Promising use of minocycline augmentation with clozapine in treatment-resistant schizophrenia. <i>Journal of Psychopharmacology</i> , 2014, 28, 707-708.	2.0	20
50	Culturally-adapted cognitive behavioural therapy based intervention for maternal depression: a mixed-methods feasibility study. <i>BMC Women's Health</i> , 2019, 19, 21.	0.8	20
51	Development and pilot testing of a social intervention for depressed women of Pakistani family origin in the UK. <i>Journal of Mental Health</i> , 2009, 18, 504-509.	1.0	19
52	Maternal depression and infant growth and development in British Pakistani women: a cohort study. <i>BMJ Open</i> , 2012, 2, e000523.	0.8	19
53	Explanatory models of psychosis amongst British South Asians. <i>Asian Journal of Psychiatry</i> , 2015, 16, 48-54.	0.9	19
54	Amoxapine as an Antipsychotic. <i>Journal of Clinical Psychopharmacology</i> , 2007, 27, 575-581.	0.7	17

#	ARTICLE	IF	CITATIONS
55	Somatization among ethnic minorities and immigrants: Why does it matter to Consultation Liaison Psychiatry?. <i>Journal of Psychosomatic Research</i> , 2015, 79, 85-86.	1.2	17
56	Pilot study of a culturally adapted psychoeducation (CaPE) intervention for bipolar disorder in Pakistan. <i>International Journal of Bipolar Disorders</i> , 2017, 5, 3.	0.8	17
57	Efficacy of learning through play plus intervention to reduce maternal depression in women with malnourished children: A randomized controlled trial from Pakistan. <i>Journal of Affective Disorders</i> , 2021, 278, 78-84.	2.0	17
58	TechCare: mobile assessment and therapy for psychosis – an intervention for clients in the Early Intervention Service: A feasibility study protocol. <i>SAGE Open Medicine</i> , 2016, 4, 205031211666961.	0.7	16
59	Cognitive Behavior Therapy for psychosis based Guided Self-help (CBTp-GSH) delivered by frontline mental health professionals: Results of a feasibility study. <i>Schizophrenia Research</i> , 2016, 173, 69-74.	1.1	16
60	Adjunctive simvastatin for treatment-resistant depression: study protocol of a 12-week randomised controlled trial. <i>BJPsych Open</i> , 2019, 5, e13.	0.3	16
61	Minocycline as adjunctive treatment for major depressive disorder: Pooled data from two randomized controlled trials. <i>Australian and New Zealand Journal of Psychiatry</i> , 2021, 55, 784-798.	1.3	16
62	Psychosocial family interventions for relatives of people living with psychotic disorders in the Arab world: systematic review. <i>BMC Psychiatry</i> , 2020, 20, 413.	1.1	16
63	Developing a healthy living intervention for people with early psychosis using the Medical Research Council's guidelines on complex interventions: Phase 1 of the HELPER – InterACT programme. <i>International Journal of Nursing Studies</i> , 2012, 49, 398-406.	2.5	15
64	Depression and congestive heart failure: A large prospective cohort study from Pakistan. <i>Journal of Psychosomatic Research</i> , 2019, 120, 46-52.	1.2	15
65	Psychological interventions for maternal depression among women of African and Caribbean origin: a systematic review. <i>BMC Women's Health</i> , 2021, 21, 83.	0.8	15
66	A prospective study of social difficulties, acculturation and persistent depression in Pakistani women living in the UK. <i>Psychological Medicine</i> , 2012, 42, 1217-1226.	2.7	14
67	Antenatal Depression is Not Associated with Low-Birth Weight: A Study from Urban Pakistan. <i>Frontiers in Psychiatry</i> , 2014, 5, 175.	1.3	14
68	A randomised double-blind placebo-controlled 12-week feasibility trial of methotrexate added to treatment as usual in early schizophrenia: study protocol for a randomised controlled trial. <i>Trials</i> , 2015, 16, 9.	0.7	14
69	Pilot randomised controlled trial of culturally adapted cognitive behavior therapy for psychosis (CaCBTp) in Pakistan. <i>BMC Health Services Research</i> , 2017, 17, 808.	0.9	14
70	Psychological distress among patients attending a general medical outpatient clinic in Pakistan. <i>General Hospital Psychiatry</i> , 2004, 26, 277-281.	1.2	13
71	The validation of personal health questionnaire amongst people of Pakistani family origin living in the United Kingdom. <i>Journal of Affective Disorders</i> , 2007, 97, 261-264.	2.0	13
72	A randomised, double-blind, placebo-controlled trial of minocycline and/or omega-3 fatty acids added to treatment as usual for at-risk mental states (NAYAB): study protocol. <i>Trials</i> , 2017, 18, 524.	0.7	13

#	ARTICLE	IF	CITATIONS
73	Demographic and psychosocial characteristics of self-harm: The Pakistan perspective. <i>Psychiatry Research</i> , 2019, 279, 201-206.	1.7	13
74	An integrated parenting intervention for maternal depression and child development in a low-resource setting: Cluster randomized controlled trial. <i>Depression and Anxiety</i> , 2021, 38, 925-939.	2.0	13
75	Relationship between childhood trauma, personality, social support and depression in women attending general medical clinics in a low and middle-income country. <i>Journal of Affective Disorders</i> , 2021, 292, 526-533.	2.0	13
76	Psychological distress among patients of an orthopaedic outpatient clinic: a study from a low-income country. <i>Annals of General Psychiatry</i> , 2010, 9, 9.	1.2	12
77	A roadmap to develop dementia research capacity and capability in Pakistan: A model for low and middle-income countries. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2019, 5, 939-952.	1.8	12
78	Protocol for a cluster randomised trial evaluating a multifaceted intervention starting preconceptionally: Early Interventions to Support Trajectories for Healthy Life in India (EINSTEIN): a Healthy Life Trajectories Initiative (HeLTI) Study. <i>BMJ Open</i> , 2021, 11, e045862.	0.8	12
79	Ondansetron and simvastatin added to treatment as usual in patients with schizophrenia: study protocol for a randomized controlled trial. <i>Trials</i> , 2013, 14, 101.	0.7	11
80	Mental health services in Libya. <i>BJPsych International</i> , 2016, 13, 70-71.	0.8	11
81	Validation of the self-reporting questionnaire (SRQ 20) in British Pakistani and White European population in the United Kingdom. <i>Journal of Affective Disorders</i> , 2016, 189, 392-396.	2.0	11
82	Stigma toward mental and physical illness: attitudes of healthcare professionals, healthcare students and the general public in Pakistan. <i>BJPsych Open</i> , 2020, 6, e81.	0.3	11
83	A Group Parenting Intervention for Depressed Fathers (LTP + Dads): A Feasibility Study from Pakistan. <i>Children</i> , 2021, 8, 26.	0.6	11
84	Maternal mental health and child well-being in Nigeria: A systematic review. <i>Health Psychology Open</i> , 2021, 8, 205510292110121.	0.7	10
85	Detecting Maternal Depression in a Low-Income Country: Comparison of the Self-Reporting Questionnaire and the Edinburgh Postnatal Depression Scale. <i>Journal of Tropical Pediatrics</i> , 2014, 60, 129-133.	0.7	9
86	Prevalence of Common Mental Disorders and its Association with Life Events and Social Support in Mothers Attending a Well-Child Clinic. <i>SAGE Open</i> , 2016, 6, 215824401667732.	0.8	9
87	Culturally adapted family intervention for schizophrenia in Pakistan: a feasibility study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2021, 25, 258-267.	1.2	9
88	Mental health experiences of mothers in Jos, Nigeria: An interpretative phenomenological analysis. <i>SAGE Open Medicine</i> , 2021, 9, 205031212097071.	0.7	9
89	Prevalence of depression and anxiety in patients with chronic obstructive pulmonary disease and their association with psychosocial outcomes: A cross-sectional study from Pakistan. <i>SAGE Open Medicine</i> , 2021, 9, 205031212110328.	0.7	9
90	The benefit of minocycline on negative symptoms in early-phase psychosis in addition to standard care - extent and mechanism (BeneMin): study protocol for a randomised controlled trial. <i>Trials</i> , 2015, 16, 71.	0.7	8

#	ARTICLE	IF	CITATIONS
91	The relationship between childhood trauma and adult psychosis in a UK Early Intervention Service: results of a retrospective case note study. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 269-273.	1.0	8
92	Posttraumatic stress symptomology and non-suicidal self-injury: The role of intrusion and arousal symptoms. <i>Journal of Affective Disorders</i> , 2020, 276, 920-926.	2.0	8
93	Suicidality in patients with bipolar depression: Findings from a lower middle-income country. <i>Journal of Affective Disorders</i> , 2021, 289, 1-6.	2.0	8
94	Exploring the experience of relatives living with individuals diagnosed with schizophrenia in Oman: A qualitative study. <i>Journal of Psychiatric and Mental Health Nursing</i> , 2021, 28, 1029-1040.	1.2	8
95	Prevalence and risk factors for psychological distress and functional disability in urban Pakistan. <i>WHO South-East Asia Journal of Public Health</i> , 2014, 3, 144.	1.7	8
96	Group psychological intervention for maternal depression: A nested qualitative study from Karachi, Pakistan. <i>World Journal of Psychiatry</i> , 2017, 7, 98.	1.3	8
97	Care co-ordinator in my pocket™: a feasibility study of mobile assessment and therapy for psychosis (TechCare). <i>BMJ Open</i> , 2021, 11, e046755.	0.8	8
98	Prevention of self harm in British South Asian women: study protocol of an exploratory RCT of culturally adapted manual assisted Problem Solving Training (C- MAP). <i>Trials</i> , 2011, 12, 159.	0.7	7
99	Recruitment strategies for British South Asians in 5 depression trials: A mixed method study. <i>Journal of Affective Disorders</i> , 2015, 185, 195-203.	2.0	7
100	Glycopyrrolate in comparison to hyoscine hydrobromide and placebo in the treatment of hypersalivation induced by clozapine (GOTHIC1): study protocol for a randomised controlled feasibility study. <i>Trials</i> , 2016, 17, 553.	0.7	7
101	Considerations for clinicians treating Muslim patients with psychiatric disorders during Ramadan. <i>Lancet Psychiatry</i> , 2019, 6, 556-557.	3.7	7
102	Exploring Lived Experiences of Adolescents Presenting with Self-Harm and Their Views about Suicide Prevention Strategies: A Qualitative Approach. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 4694.	1.2	7
103	Psychological intervention, antipsychotic medication or a combined treatment for adolescents with a first episode of psychosis: the MAPS feasibility three-arm RCT. <i>Health Technology Assessment</i> , 2021, 25, 1-124.	1.3	7
104	Self-esteem and obsessive compulsive disorder. <i>JPMA the Journal of the Pakistan Medical Association</i> , 2014, 64, 64-8.	0.1	7
105	Medically Unexplained Symptoms—A Perspective from General Practitioners in the Developing World. <i>International Journal of Psychiatry in Medicine</i> , 2011, 42, 1-11.	0.8	6
106	Video intervention for the psychiatric waiting room: proof-of-concept randomised controlled trial of RESOLVE (Relaxation Exercise, SOLVing problem and cognitive Errors). <i>BJPsych Open</i> , 2019, 5, e77.	0.3	6
107	Montessori intervention for individuals with dementia: feasibility study of a culturally adapted psychosocial intervention in Pakistan (MIRACLE). <i>BJPsych Open</i> , 2020, 6, e69.	0.3	6
108	A review of economic evaluations of health care for people at risk of psychosis and for first-episode psychosis. <i>BMC Psychiatry</i> , 2022, 22, 126.	1.1	6

#	ARTICLE	IF	CITATIONS
109	Impact of COVID-19 on mental health research: is this the breaking point?. <i>British Journal of Psychiatry</i> , 2022, 220, 254-256.	1.7	6
110	Treatment of maternal depression in low-income women: A feasibility study from Kilifi, Kenya. <i>International Journal of Clinical Practice</i> , 2021, 75, e14862.	0.8	5
111	Depressive disorder and social stress in Pakistan compared to people of Pakistani origin in the UK. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2011, 46, 1153-1159.	1.6	4
112	Which antidepressant would psychiatrists and nurses from a developing country choose for themselves?. <i>International Journal of Psychiatry in Clinical Practice</i> , 2011, 15, 74-78.	1.2	4
113	Medical professionalism education for psychiatry trainees: does it meet standards?. <i>Australasian Psychiatry</i> , 2015, 23, 72-75.	0.4	4
114	Cognitive function in early psychosis patients from a lower middle-income country. <i>International Journal of Psychiatry in Clinical Practice</i> , 2018, 22, 13-18.	1.2	4
115	Depression and suicidal ideation in schizophrenia spectrum disorder: a cross-sectional study from a lower middle-income country. <i>International Journal of Psychiatry in Clinical Practice</i> , 2021, 25, 245-251.	1.2	4
116	The epidemiology of common mental disorders in Libya: a systematic review. <i>Egyptian Journal of Neurology, Psychiatry and Neurosurgery</i> , 2021, 57, .	0.4	4
117	Psychosocial Correlates, Psychological Distress, and Quality of Life in Patients with Medically Unexplained Symptoms. <i>International Journal of Psychiatry in Medicine</i> , 2014, 48, 235-251.	0.8	3
118	Clinicians' Perspectives on Self-Harm in Pakistan: A Qualitative Study. <i>Frontiers in Psychiatry</i> , 2021, 12, 607549.	1.3	3
119	Brief Intervention Models in Psychosis for Developing Countries (Asia and Africa). , 2016, , 191-204.		3
120	Metabolic function in patients with bipolar depression receiving anti-inflammatory agents: Findings from the MINDCARE study, a multicentre, randomised controlled trial. <i>Journal of Affective Disorders</i> , 2022, 299, 135-141.	2.0	3
121	Multicentre randomised controlled trial of a group psychological intervention for postnatal depression in British mothers of South Asian origin (ROSHNI-2): study protocol. <i>BJPsych Open</i> , 2022, 8, .	0.3	3
122	Capacity & capability building for applied dementia research in low- & middle-income countries: Two exemplars from South Asia. <i>Indian Journal of Medical Research</i> , 2020, 152, 614.	0.4	3
123	Protocol of a feasibility trial for an online group parenting intervention with an integrated mental health component for parent refugees and asylum-seekers in the United Kingdom: (LTP + EMDR G-TEP). <i>SAGE Open Medicine</i> , 2021, 9, 205031212110678.	0.7	3
124	Cognitive function in early-phase schizophrenia-spectrum disorder: IQ subtypes, brain volume and immune markers. <i>Psychological Medicine</i> , 2022, , 1-10.	2.7	3
125	What works? Lessons from a pretrial qualitative study to inform a multi-component intervention for refugees and asylum seekers: Learning Through Play and EMDR Group Traumatic Episode Protocol. <i>Journal of Community Psychology</i> , 2023, 51, 361-381.	1.0	3
126	A Survey of British Senior Psychiatry Trainees' Ethnocultural Personal Values. <i>Academic Psychiatry</i> , 2009, 33, 423-426.	0.4	2



#	ARTICLE	IF	CITATIONS
127	SENSE-Cog Asia: A Feasibility Study of a Hearing Intervention to Improve Outcomes in People With Dementia. <i>Frontiers in Neurology</i> , 2021, 12, 654143.	1.1	2
128	Cost-Effectiveness of a Culturally Adapted Manual-Assisted Brief Psychological Intervention for Self-Harm in Pakistan: A Secondary Analysis of the Culturally Adapted Manual-Assisted Problem-Solving Training Randomized Controlled Trial. <i>Value in Health Regional Issues</i> , 2021, 25, 150-156.	0.5	2
129	Minocycline for negative symptoms of schizophrenia and possible mechanistic actions: the BeneMin RCT. <i>Efficacy and Mechanism Evaluation</i> , 2019, 6, 1-66.	0.9	2
130	Cost-Effectiveness of a Culturally Adapted Manual-Assisted Brief Psychological Intervention for Self-Harm in Pakistan: A Secondary Analysis of the Culturally Adapted Manual-Assisted Brief Psychological Randomized Controlled Trial. <i>Value in Health Regional Issues</i> , 2022, 27, 65-71.	0.5	2
131	Exploring views of relatives and mental health professionals to inform the cultural adaptation of a family intervention for schizophrenia in Oman: A qualitative study. <i>International Journal of Social Psychiatry</i> , 2022, 68, 1795-1805.	1.6	2
132	The clinical effectiveness and cost effectiveness of clozapine for inpatients with severe borderline personality disorder (CALMED study): a randomised placebo-controlled trial. <i>Therapeutic Advances in Psychopharmacology</i> , 2022, 12, 204512532210908.	1.2	2
133	Prevention of suicide in ethnic minorities in the UK. <i>Ethnicity and Inequalities in Health and Social Care</i> , 2009, 2, 10-17.	0.4	1
134	Tardive dyskinesia in a South Asian population with first episode psychosis treated with antipsychotics. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1953.	1.0	1
135	Glycopyrrolate in comparison to hyoscine hydrobromide and placebo in the treatment of hypersalivation induced by clozapine (GOTHIC1): a feasibility study. <i>Pilot and Feasibility Studies</i> , 2019, 5, 79.	0.5	1
136	Capacity & capability building for applied dementia research in low- & middle-income countries: Two exemplars from South Asia. <i>Indian Journal of Medical Research</i> , 2020, 152, 614-625.	0.4	1
137	Youth Culturally adapted Manual Assisted Problem Solving Training (YCMAP) in Pakistani adolescent with a history of self-harm: protocol for multicentre clinical and cost-effectiveness randomised controlled trial. <i>BMJ Open</i> , 2022, 12, e056301.	0.8	1
138	Cross-Cultural Issues. <i>Integrating Psychiatry and Primary Care</i> , 2019, , 117-136.	0.3	0
139	Building Resilience Against Violence (BRAVE): protocol of a parenting intervention for mothers and fathers with post-traumatic stress disorder in Pakistan. <i>Global Mental Health (Cambridge, England)</i> , 0, 1-8.	1.0	0