## Nusrat Husain

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6937757/publications.pdf

Version: 2024-02-01

139 papers

3,750 citations

34 h-index 54 g-index

140 all docs

140 docs citations

140 times ranked

4207 citing authors

#	Article	IF	CITATIONS
1	What works? Lessons from a pretrial qualitative study to inform a multiâ€component intervention for refugees and asylum seekers: Learning Through Play and EMDR Group Traumatic Episode Protocol. Journal of Community Psychology, 2023, 51, 361-381.	1.0	3
2	Cost-Effectiveness of a Culturally Adapted Manual-Assisted Brief Psychological Intervention for Self-Harm in Pakistan: A Secondary Analysis of the Culturally Adapted Manual-Assisted Brief Psychological Randomized Controlled Trial. Value in Health Regional Issues, 2022, 27, 65-71.	0.5	2
3	Metabolic function in patients with bipolar depression receiving anti-inflammatory agents: Findings from the MINDCARE study, a multicentre, randomised controlled trial. Journal of Affective Disorders, 2022, 299, 135-141.	2.0	3
4	Multicentre randomised controlled trial of a group psychological intervention for postnatal depression in British mothers of South Asian origin (ROSHNI-2): study protocol. BJPsych Open, 2022, 8,	0.3	3
5	Exploring views of relatives and mental health professionals to inform the cultural adaptation of a family intervention for schizophrenia in Oman: A qualitative study. International Journal of Social Psychiatry, 2022, 68, 1795-1805.	1.6	2
6	Cognitive function in early-phase schizophrenia-spectrum disorder: IQ subtypes, brain volume and immune markers. Psychological Medicine, 2022, , $1\text{-}10$ .	2.7	3
7	A review of economic evaluations of health care for people at risk of psychosis and for first-episode psychosis. BMC Psychiatry, 2022, 22, 126.	1.1	6
8	Impact of COVID-19 on mental health research: is this the breaking point?. British Journal of Psychiatry, 2022, 220, 254-256.	1.7	6
9	The clinical effectiveness and cost effectiveness of clozapine for inpatients with severe borderline personality disorder (CALMED study): a randomised placebo-controlled trial. Therapeutic Advances in Psychopharmacology, 2022, 12, 204512532210908.	1.2	2
10	Youth Culturally adapted Manual Assisted Problem Solving Training (YCMAP) in Pakistani adolescent with a history of self-harm: protocol for multicentre clinical and cost-effectiveness randomised controlled trial. BMJ Open, 2022, 12, e056301.	0.8	1
11	Culturally adapted family intervention for schizophrenia in Pakistan: a feasibility study. International Journal of Psychiatry in Clinical Practice, 2021, 25, 258-267.	1.2	9
12	Minocycline as adjunctive treatment for major depressive disorder: Pooled data from two randomized controlled trials. Australian and New Zealand Journal of Psychiatry, 2021, 55, 784-798.	1.3	16
13	Deconstructing depression and negative symptoms of schizophrenia; differential and longitudinal immune correlates, and response to minocycline treatment. Brain, Behavior, and Immunity, 2021, 91, 498-504.	2.0	28
14	Efficacy of learning through play plus intervention to reduce maternal depression in women with malnourished children: A randomized controlled trial from Pakistan✰. Journal of Affective Disorders, 2021, 278, 78-84.	2.0	17
15	Maternal mental health and child well-being in Nigeria: A systematic review. Health Psychology Open, 2021, 8, 205510292110121.	0.7	10
16	A Group Parenting Intervention for Depressed Fathers (LTP + Dads): A Feasibility Study from Pakistan. Children, 2021, 8, 26.	0.6	11
17	Group eye movement desensitization and reprocessing interventions in adults and children: A systematic review of randomized and nonrandomized trials. Clinical Psychology and Psychotherapy, 2021, 28, 784-806.	1.4	22
18	Protocol for a cluster randomised trial evaluating a multifaceted intervention starting preconceptionally—Early Interventions to Support Trajectories for Healthy Life in India (EINSTEIN): a Healthy Life Trajectories Initiative (HeLTI) Study. BMJ Open, 2021, 11, e045862.	0.8	12

#	Article	IF	CITATIONS
19	Psychological interventions for maternal depression among women of African and Caribbean origin: a systematic review. BMC Women's Health, 2021, 21, 83.	0.8	15
20	Exploring Lived Experiences of Adolescents Presenting with Self-Harm and Their Views about Suicide Prevention Strategies: A Qualitative Approach. International Journal of Environmental Research and Public Health, 2021, 18, 4694.	1.2	7
21	Clinicians' Perspectives on Self-Harm in Pakistan: A Qualitative Study. Frontiers in Psychiatry, 2021, 12, 607549.	1.3	3
22	An integrated parenting intervention for maternal depression and child development in a lowâ€resource setting: Cluster randomized controlled trial. Depression and Anxiety, 2021, 38, 925-939.	2.0	13
23	Suicidality in patients with bipolar depression: Findings from a lower middle-income country. Journal of Affective Disorders, 2021, 289, 1-6.	2.0	8
24	SENSE-Cog Asia: A Feasibility Study of a Hearing Intervention to Improve Outcomes in People With Dementia. Frontiers in Neurology, 2021, 12, 654143.	1.1	2
25	Depression and suicidal ideation in schizophrenia spectrum disorder: a cross-sectional study from a lower middle-income country. International Journal of Psychiatry in Clinical Practice, 2021, 25, 245-251.	1.2	4
26	Exploring the experience of relatives living with individuals diagnosed with schizophrenia in Oman: A qualitative study. Journal of Psychiatric and Mental Health Nursing, 2021, 28, 1029-1040.	1.2	8
27	Cost-Effectiveness of a Culturally Adapted Manual-Assisted Brief Psychological Intervention for Self-Harm in Pakistan: A Secondary Analysis of the Culturally Adapted Manual-Assisted Problem-Solving Training Randomized Controlled Trial. Value in Health Regional Issues, 2021, 25, 150-156.	0.5	2
28	Treatment of maternal depression in lowâ€income women: A feasibility study from Kilifi, Kenya. International Journal of Clinical Practice, 2021, 75, e14862.	0.8	5
29	Relationship between childhood trauma, personality, social support and depression in women attending general medical clinics in a low and middle-income country. Journal of Affective Disorders, 2021, 292, 526-533.	2.0	13
30	Psychological intervention, antipsychotic medication or a combined treatment for adolescents with a first episode of psychosis: the MAPS feasibility three-arm RCT. Health Technology Assessment, 2021, 25, 1-124.	1.3	7
31	Mental health experiences of mothers in Jos, Nigeria: An interpretative phenomenological analysis. SAGE Open Medicine, 2021, 9, 205031212097071.	0.7	9
32	Prevalence of depression and anxiety in patients with chronic obstructive pulmonary disease and their association with psychosocial outcomes: A cross-sectional study from Pakistan. SAGE Open Medicine, 2021, 9, 205031212110328.	0.7	9
33	<i>âe^</i> Care co-ordinator in my pocketâe™: a feasibility study of mobile assessment and therapy for psychosis (TechCare). BMJ Open, 2021, 11, e046755.	0.8	8
34	The epidemiology of common mental disorders in Libya: a systematic review. Egyptian Journal of Neurology, Psychiatry and Neurosurgery, 2021, 57, .	0.4	4
35	Protocol of a feasibility trial for an online group parenting intervention with an integrated mental health component for parent refugees and asylum-seekers in the United Kingdom: (LTP + EMDR G-TEP). SAGE Open Medicine, 2021, 9, 205031212110678.	0.7	3
36	A Systematic Review of the Evidence Supporting Mobile―and Internetâ€Based Psychological Interventions For Selfâ€Harm. Suicide and Life-Threatening Behavior, 2020, 50, 151-179.	0.9	61

#	Article	IF	CITATIONS
37	Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. Lancet Psychiatry,the, 2020, 7, 788-800.	3.7	51
38	Posttraumatic stress symptomology and non-suicidal self-injury: The role of intrusion and arousal symptoms. Journal of Affective Disorders, 2020, 276, 920-926.	2.0	8
39	Stigma toward mental and physical illness: attitudes of healthcare professionals, healthcare students and the general public in Pakistan. BJPsych Open, 2020, 6, e81.	0.3	11
40	Psychosocial family interventions for relatives of people living with psychotic disorders in the Arab world: systematic review. BMC Psychiatry, 2020, 20, 413.	1.1	16
41	Psychotherapy or medication for depression? Using individual symptom meta-analyses to derive a Symptom-Oriented Therapy (SOrT) metric for a personalised psychiatry. BMC Medicine, 2020, 18, 170.	2.3	21
42	Effect of delaying treatment of first-episode psychosis on symptoms and social outcomes: a longitudinal analysis and modelling study. Lancet Psychiatry, the, 2020, 7, 602-610.	3.7	66
43	Montessori intervention for individuals with dementia: feasibility study of a culturally adapted psychosocial intervention in Pakistan (MIRACLE). BJPsych Open, 2020, 6, e69.	0.3	6
44	Minocycline and celecoxib as adjunctive treatments for bipolar depression: a multicentre, factorial design randomised controlled trial. Lancet Psychiatry,the, 2020, 7, 515-527.	3.7	110
45	Capacity & capability building for applied dementia research in low- & middle-income countries: Two exemplars from South Asia. Indian Journal of Medical Research, 2020, 152, 614-625.	0.4	1
46	Capacity & Capability building for applied dementia research in low- & Capacity & Capacity & Capability building for applied dementia research in low- & Capacity & C	0.4	3
47	Glycopyrrolate in comparison to hyoscine hydrobromide and placebo in the treatment of hypersalivation induced by clozapine (GOTHIC1): a feasibility study. Pilot and Feasibility Studies, 2019, 5, 79.	0.5	1
48	An examination of the relationship between shame, guilt and self-harm: A systematic review and meta-analysis. Clinical Psychology Review, 2019, 73, 101779.	6.0	71
49	Video intervention for the psychiatric waiting room: proof-of-concept randomised controlled trial of RESOLVE (Relaxation Exercise, SOLVing problem and cognitive Errors). BJPsych Open, 2019, 5, e77.	0.3	6
50	Culturally-adapted cognitive behavioural therapy based intervention for maternal depression: a mixed-methods feasibility study. BMC Women's Health, 2019, 19, 21.	0.8	20
51	Considerations for clinicians treating Muslim patients with psychiatric disorders during Ramadan. Lancet Psychiatry,the, 2019, 6, 556-557.	3.7	7
52	Demographic and psychosocial characteristics of self-harm: The Pakistan perspective. Psychiatry Research, 2019, 279, 201-206.	1.7	13
53	Depression and congestive heart failure: A large prospective cohort study from Pakistan. Journal of Psychosomatic Research, 2019, 120, 46-52.	1.2	15
54	Adjunctive simvastatin for treatment-resistant depression: study protocol of a 12-week randomised controlled trial. BJPsych Open, 2019, 5, e13.	0.3	16

#	Article	IF	CITATIONS
55	A roadmap to develop dementia research capacity and capability in Pakistan: A model for low†and middleâ€income countries. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2019, 5, 939-952.	1.8	12
56	Minocycline for negative symptoms of schizophrenia and possible mechanistic actions: the BeneMin RCT. Efficacy and Mechanism Evaluation, 2019, 6, 1-66.	0.9	2
57	Cross-Cultural Issues. Integrating Psychiatry and Primary Care, 2019, , 117-136.	0.3	O
58	Cognitive function in early psychosis patients from a lower middle-income country. International Journal of Psychiatry in Clinical Practice, 2018, 22, 13-18.	1.2	4
59	The benefit of minocycline on negative symptoms of schizophrenia in patients with recent-onset psychosis (BeneMin): a randomised, double-blind, placebo-controlled trial. Lancet Psychiatry,the, 2018, 5, 885-894.	3.7	133
60	The current status of culturally adapted mental health interventions: a practice-focused review of meta-analyses. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 165-178.	1.0	142
61	Minocycline and celecoxib as adjunctive treatments for bipolar depression: a study protocol for a multicenter factorial design randomized controlled trial. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 1-8.	1.0	23
62	Pilot study of a culturally adapted psychoeducation (CaPE) intervention for bipolar disorder in Pakistan. International Journal of Bipolar Disorders, 2017, 5, 3.	0.8	17
63	Treatment of maternal depression in urban slums of Karachi, Pakistan: A randomized controlled trial (RCT) of an integrated maternal psychological and early child development intervention. Asian Journal of Psychiatry, 2017, 29, 63-70.	0.9	32
64	Minocycline as an adjunct for treatment-resistant depressive symptoms: A pilot randomised placebo-controlled trial. Journal of Psychopharmacology, 2017, 31, 1166-1175.	2.0	104
65	A qualitative study to explore views of patients', carers' and mental health professionals' to inform cultural adaptation of CBT for psychosis (CBTp) in China. BMC Psychiatry, 2017, 17, 131.	1.1	31
66	Pilot randomised controlled trial of culturally adapted cognitive behavior therapy for psychosis (CaCBTp) in Pakistan. BMC Health Services Research, 2017, 17, 808.	0.9	14
67	A randomised, double-blind, placebo-controlled trial of minocycline and/or omega-3 fatty acids added to treatment as usual for at-risk mental states (NAYAB): study protocol. Trials, 2017, 18, 524.	0.7	13
68	The relationship between childhood trauma and adult psychosis in a UK Early Intervention Service: results of a retrospective case note study. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 269-273.	1.0	8
69	mHealth based interventions for the assessment and treatment of psychotic disorders: a systematic review. MHealth, 2017, 3, 33-33.	0.9	33
70	Group psychological intervention for maternal depression: A nested qualitative study from Karachi, Pakistan. World Journal of Psychiatry, 2017, 7, 98.	1.3	8
71	Mental health services in Libya. BJPsych International, 2016, 13, 70-71.	0.8	11
72	A Qualitative Study to Explore Patients', Carers' and Health Professionals' Views to Culturally Adapt CBT for Psychosis (CBTp) in Pakistan. Behavioural and Cognitive Psychotherapy, 2016, 44, 43-55.	0.9	39

#	Article	IF	Citations
73	TechCare: mobile assessment and therapy for psychosis $\hat{a}\in$ an intervention for clients in the Early Intervention Service: A feasibility study protocol. SAGE Open Medicine, 2016, 4, 205031211666961.	0.7	16
74	Glycopyrrolate in comparison to hyoscine hydrobromide and placebo in the treatment of hypersalivation induced by clozapine (GOTHIC1): study protocol for a randomised controlled feasibility study. Trials, 2016, 17, 553.	0.7	7
75	Prevalence of Common Mental Disorders and its Association with Life Events and Social Support in Mothers Attending a Well-Child Clinic. SAGE Open, 2016, 6, 215824401667732.	0.8	9
76	Increasing access to Cognitive Behaviour Therapy in Low and Middle Income Countries: A strategic framework. Asian Journal of Psychiatry, 2016, 22, 190-195.	0.9	28
77	Barefoot therapists: barriers and facilitators to delivering maternal mental health care through peer volunteers in Pakistan: a qualitative study. International Journal of Mental Health Systems, 2016, 10, 24.	1.1	53
78	Cognitive Behavior Therapy for psychosis based Guided Self-help (CBTp-GSH) delivered by frontline mental health professionals: Results of a feasibility study. Schizophrenia Research, 2016, 173, 69-74.	1.1	16
79	Validation of the self-reporting questionnaire (SRQ 20) in British Pakistani and White European population in the United Kingdom. Journal of Affective Disorders, 2016, 189, 392-396.	2.0	11
80	Brief Intervention Models in Psychosis for Developing Countries (Asia and Africa)., 2016,, 191-204.		3
81	Minocycline as an adjunct for treatment-resistant depressive symptoms: study protocol for a pilot randomised controlled trial. Trials, 2015, 16, 410.	0.7	30
82	Group psychological intervention for postnatal depression: a nested qualitative study with British South Asian women. BMC Women's Health, 2015, 15, 109.	0.8	22
83	Medical professionalism education for psychiatry trainees: does it meet standards?. Australasian Psychiatry, 2015, 23, 72-75.	0.4	4
84	An evaluation of subjective experiences, effects and overall satisfaction with clozapine treatment in a UK forensic service. Therapeutic Advances in Psychopharmacology, 2015, 5, 146-150.	1.2	22
85	Recruitment strategies for British South Asians in 5 depression trials: A mixed method study. Journal of Affective Disorders, 2015, 185, 195-203.	2.0	7
86	Somatization among ethnic minorities and immigrants: Why does it matter to Consultation Liaison Psychiatry?. Journal of Psychosomatic Research, 2015, 79, 85-86.	1,2	17
87	Brief Culturally adapted CBT (CaCBT) for depression: A randomized controlled trial from Pakistan. Journal of Affective Disorders, 2015, 177, 101-107.	2.0	62
88	A randomised double-blind placebo-controlled 12- week feasibility trial of methotrexate added to treatment as usual in early schizophrenia: study protocol for a randomised controlled trial. Trials, 2015, 16, 9.	0.7	14
89	The benefit of minocycline on negative symptoms in early-phase psychosis in addition to standard care - extent and mechanism (BeneMin): study protocol for a randomised controlled trial. Trials, 2015, 16, 71.	0.7	8
90	Brief culturally adapted CBT for psychosis (CaCBTp): A randomized controlled trial from a low income country. Schizophrenia Research, 2015, 164, 143-148.	1.1	75

#	Article	IF	Citations
91	Explanatory models of psychosis amongst British South Asians. Asian Journal of Psychiatry, 2015, 16, 48-54.	0.9	19
92	Antenatal Depression is Not Associated with Low-Birth Weight: A Study from Urban Pakistan. Frontiers in Psychiatry, 2014, 5, 175.	1.3	14
93	Tardive dyskinesia in a South Asian population with first episode psychosis treated with antipsychotics. Neuropsychiatric Disease and Treatment, 2014, 10, 1953.	1.0	1
94	Antidepressant and Group Psychosocial Treatment for Depression: A Rater Blind Exploratory RCT from a Low Income Country. Behavioural and Cognitive Psychotherapy, 2014, 42, 693-705.	0.9	23
95	A naturalistic, randomized, controlled trial combining cognitive remediation with cognitive–behavioural therapy after first-episode non-affective psychosis. Psychological Medicine, 2014, 44, 1889-1899.	2.7	49
96	Detecting Maternal Depression in a Low-Income Country: Comparison of the Self-Reporting Questionnaire and the Edinburgh Postnatal Depression Scale. Journal of Tropical Pediatrics, 2014, 60, 129-133.	0.7	9
97	Impact of early intervention services on duration of untreated psychosis: Data from the National EDEN prospective cohort study. Schizophrenia Research, 2014, 159, 1-6.	1.1	38
98	Brief psychological intervention after self-harm: randomised controlled trial from Pakistan. British Journal of Psychiatry, 2014, 204, 462-470.	1.7	67
99	Detecting Depression in Pregnancy: Validation of EPDS in British Pakistani Mothers. Journal of Immigrant and Minority Health, 2014, 16, 1085-1092.	0.8	25
100	Promising use of minocycline augmentation with clozapine in treatment-resistant schizophrenia. Journal of Psychopharmacology, 2014, 28, 707-708.	2.0	20
101	A multicentre randomised controlled trial of a carer supervised Culturally adapted CBT (CaCBT) based self-help for depression in Pakistan. Journal of Affective Disorders, 2014, 156, 224-227.	2.0	47
102	Psychosocial Correlates, Psychological Distress, and Quality of Life in Patients with Medically Unexplained Symptoms. International Journal of Psychiatry in Medicine, 2014, 48, 235-251.	0.8	3
103	An Exploratory Randomized Controlled Study of a Healthy Living Intervention in Early Intervention Services for Psychosis. Journal of Clinical Psychiatry, 2014, 75, 498-505.	1.1	44
104	Prevalence and risk factors for psychological distress and functional disability in urban Pakistan. WHO South-East Asia Journal of Public Health, 2014, 3, 144.	1.7	8
105	Self-esteem and obsessive compulsive disorder. JPMA the Journal of the Pakistan Medical Association, 2014, 64, 64-8.	0.1	7
106	Ondansetron and simvastatin added to treatment as usual in patients with schizophrenia: study protocol for a randomized controlled trial. Trials, 2013, 14, 101.	0.7	11
107	A systematic review of explanatory models of illness for psychosis in developing countries. International Review of Psychiatry, 2012, 24, 450-462.	1.4	45
108	Maternal depression and infant growth and development in British Pakistani women: a cohort study. BMJ Open, 2012, 2, e000523.	0.8	19

7

#	Article	IF	Citations
109	A prospective study of social difficulties, acculturation and persistent depression in Pakistani women living in the UK. Psychological Medicine, 2012, 42, 1217-1226.	2.7	14
110	Minocycline benefits negative symptoms in early schizophrenia: a randomised double-blind placebo-controlled clinical trial in patients on standard treatment. Journal of Psychopharmacology, 2012, 26, 1185-1193.	2.0	247
111	Developing a healthy living intervention for people with early psychosis using the Medical Research Council's guidelines on complex interventions: Phase 1 of the HELPER $\hat{a} \in \mathbb{C}$ InterACT programme. International Journal of Nursing Studies, 2012, 49, 398-406.	2.5	15
112	Social stress and depression during pregnancy and in the postnatal period in British Pakistani mothers: A cohort study. Journal of Affective Disorders, 2012, 140, 268-276.	2.0	77
113	Medically Unexplained Symptomsâ€"A Perspective from General Practitioners in the Developing World. International Journal of Psychiatry in Medicine, 2011, 42, 1-11.	0.8	6
114	Prevalence and psychosocial correlates of perinatal depression: a cohort study from urban Pakistan. Archives of Women's Mental Health, 2011, 14, 395-403.	1.2	55
115	Depressive disorder and social stress in Pakistan compared to people of Pakistani origin in the UK. Social Psychiatry and Psychiatric Epidemiology, 2011, 46, 1153-1159.	1.6	4
116	Prevention of: self harm in British South Asian women: study protocol of an exploratory RCT of culturally adapted manual assisted Problem Solving Training (C- MAP). Trials, 2011, 12, 159.	0.7	7
117	Which antidepressant would psychiatrists and nurses from a developing country choose for themselves?. International Journal of Psychiatry in Clinical Practice, 2011, 15, 74-78.	1.2	4
118	Social intervention for British Pakistani women with depression: randomised controlled trial. British Journal of Psychiatry, 2010, 197, 227-233.	1.7	55
119	Psychological distress among patients of an orthopaedic outpatient clinic: a study from a low-income country. Annals of General Psychiatry, 2010, 9, 9.	1.2	12
120	Multidimensional scale of perceived social support: Psychometric properties in a South Asian population. Journal of Obstetrics and Gynaecology Research, 2010, 36, 845-851.	0.6	91
121	Development and pilot testing of a social intervention for depressed women of Pakistani family origin in the UK. Journal of Mental Health, 2009, 18, 504-509.	1.0	19
122	Cluster randomized trial of a parentâ€based intervention to support early development of children in a lowâ€income country. Child: Care, Health and Development, 2009, 35, 56-62.	0.8	78
123	A Survey of British Senior Psychiatry Trainees' Ethnocultural Personal Values. Academic Psychiatry, 2009, 33, 423-426.	0.4	2
124	Prevention of suicide in ethnic minorities in the UK. Ethnicity and Inequalities in Health and Social Care, 2009, 2, 10-17.	0.4	1
125	A populationâ€based study of irritable bowel syndrome in a nonâ€Western population. Neurogastroenterology and Motility, 2008, 20, 1022-1029.	1.6	62
126	Ethnicity and psychopharmacology. Journal of Psychopharmacology, 2008, 22, 673-680.	2.0	63

#	Article	lF	CITATIONS
127	Medically unexplained presentations and quality of life: A study of a predominantly South Asian primary care population in England. Journal of Psychosomatic Research, 2008, 65, 311-317.	1.2	39
128	Life stress and depression in a tribal area of Pakistan. British Journal of Psychiatry, 2007, 190, 36-41.	1.7	56
129	Amoxapine as an Antipsychotic. Journal of Clinical Psychopharmacology, 2007, 27, 575-581.	0.7	17
130	The validation of personal health questionnaire amongst people of Pakistani family origin living in the United Kingdom. Journal of Affective Disorders, 2007, 97, 261-264.	2.0	13
131	Self-harm in British South Asian women: psychosocial correlates and strategies for prevention. Annals of General Psychiatry, 2006, 5, 7.	1.2	42
132	Prevalence and social correlates of postnatal depression in a low income country. Archives of Women's Mental Health, 2006, 9, 197-202.	1.2	153
133	Self-harm in the UK. Social Psychiatry and Psychiatric Epidemiology, 2006, 41, 782-788.	1.6	65
134	Comparison of the Personal Health Questionnaire and the Self Reporting Questionnaire in rural Pakistan. JPMA the Journal of the Pakistan Medical Association, 2006, 56, 366-70.	0.1	26
135	Psychological distress among patients attending a general medical outpatient clinic in Pakistan. General Hospital Psychiatry, 2004, 26, 277-281.	1.2	13
136	Social factors associated with chronic depression among a population-based sample of women in rural Pakistan. Social Psychiatry and Psychiatric Epidemiology, 2004, 39, 618-24.	1.6	44
137	Depression and social stress in Pakistan. Psychological Medicine, 2000, 30, 395-402.	2.7	144
138	Adverse social circumstances and depression in people of Pakistani origin in the UK. British Journal of Psychiatry, 1997, 171, 434-438.	1.7	68
139	Building Resilience Against ViolencE (BRAVE): protocol of a parenting intervention for mothers and fathers with post-traumatic stress disorder in Pakistan. Global Mental Health (Cambridge, England), 0,	1.0	О