

Igor Bombin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6939280/publications.pdf>

Version: 2024-02-01

27
papers

1,184
citations

361413

20
h-index

552781

26
g-index

32
all docs

32
docs citations

32
times ranked

1817
citing authors

#	ARTICLE	IF	CITATIONS
1	Fronto-Parietal Gray Matter Volume Loss Is Associated with Decreased Working Memory Performance in Adolescents with a First Episode of Psychosis. <i>Journal of Clinical Medicine</i> , 2021, 10, 3929.	2.4	4
2	Premorbid adjustment and clinical correlates of cognitive impairment in first-episode psychosis. The PEPsCog Study. <i>Schizophrenia Research</i> , 2015, 164, 65-73.	2.0	92
3	Predictors of schizophrenia spectrum disorders in early-onset first episodes of psychosis: a support vector machine model. <i>European Child and Adolescent Psychiatry</i> , 2015, 24, 427-440.	4.7	37
4	Cognitive reserve as a predictor of two year neuropsychological performance in early onset first-episode schizophrenia. <i>Schizophrenia Research</i> , 2013, 143, 125-131.	2.0	61
5	Neuropsychological evidence for abnormal neurodevelopment associated with early-onset psychoses. <i>Psychological Medicine</i> , 2013, 43, 757-768.	4.5	56
6	Functional impairment as a defining feature of: amnestic MCI cognitive, emotional, and demographic correlates. <i>International Psychogeriatrics</i> , 2012, 24, 1494-1504.	1.0	23
7	Longitudinal study of neurological soft signs in first-episode early-onset psychosis. <i>Journal of Child Psychology and Psychiatry and Allied Disciplines</i> , 2012, 53, 323-331.	5.2	39
8	Genetic variation in dysbindin gene influences both the risk for functional psychosis and the cognitive functioning in a Spanish family based study. <i>International Clinical Psychopharmacology</i> , 2011, 26, e59-e60.	1.7	0
9	Dysbindin-1 gene contributes differentially to early- and adult-onset forms of functional psychosis. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2011, 156, 322-333.	1.7	22
10	Cognitive Efficacy of Quetiapine and Olanzapine in Early-Onset First-Episode Psychosis. <i>Schizophrenia Bulletin</i> , 2011, 37, 405-415.	4.3	18
11	Cognitive Functioning in Children and Adolescents in Their First Episode of Psychosis. <i>Journal of Nervous and Mental Disease</i> , 2010, 198, 159-162.	1.0	36
12	Neuropsychological functioning in early-onset first-episode psychosis: comparison of diagnostic subgroups. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2010, 260, 225-233.	3.2	69
13	GENETIC VARIABILITY IN DYSBINDIN-1 GENE (DTNBP1) CONTRIBUTES DIFFERENTIALLY TO EARLY AND ADULT ONSET FUNCTIONAL PSYCHOSES AND IT IS ASSOCIATED WITH THE FAMILIAL TRANSMISSION OF IQ AND PREFRONTAL COGNITIVE DEFICITS. <i>Schizophrenia Research</i> , 2010, 117, 220-221.	2.0	0
14	Predominance of Symptoms Over Time in Early-Onset Psychosis. <i>Journal of Clinical Psychiatry</i> , 2010, 71, 327-337.	2.2	26
15	Insight correlates in child- and adolescent-onset first episodes of psychosis: results from the CAFEPS study. <i>Psychological Medicine</i> , 2009, 39, 1433-1445.	4.5	43
16	Brain morphology and neurological soft signs in adolescents with first-episode psychosis. <i>British Journal of Psychiatry</i> , 2009, 195, 227-233.	2.8	50
17	Olanzapine compared to quetiapine in adolescents with a first psychotic episode. <i>European Child and Adolescent Psychiatry</i> , 2009, 18, 418-428.	4.7	66
18	DRD3, but not COMT or DRD2, genotype affects executive functions in healthy and first-episode psychosis adolescents. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2008, 147B, 873-879.	1.7	53

#	ARTICLE	IF	CITATIONS
19	COMT AND DRD3, BUT NOT DRD2, GENOTYPE AFFECT SOFT NEUROLOGICAL SIGNS IN HEALTHY AND FIRST-EPIISODE PSYCHOSIS ADOLESCENTS. <i>Schizophrenia Research</i> , 2008, 102, 183.	2.0	0
20	Neurological soft signs in adolescents with first episode psychosis: Two-year followup. <i>Psychiatry Research</i> , 2008, 161, 344-348.	3.3	30
21	Association study between obsessive-compulsive disorder and serotonergic candidate genes. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 765-770.	4.8	47
22	Neuropsychological functioning in adolescents with first episode psychosis: A two-year follow-up study. <i>European Psychiatry</i> , 2008, 23, 375-383.	0.2	40
23	The genetic contribution to first psychotic episodes in children and adolescents of the child and adolescent first-episode psychosis study. <i>Psychiatric Genetics</i> , 2008, 18, 151-152.	1.1	1
24	Randomised clinical trial comparing oral versus depot formulations of zuclopenthixol in patients with schizophrenia and previous violence. <i>European Psychiatry</i> , 2006, 21, 34-40.	0.2	70
25	Significance and Meaning of Neurological Signs in Schizophrenia: Two Decades Later. <i>Schizophrenia Bulletin</i> , 2005, 31, 962-977.	4.3	240
26	Assessment Tools for Soft Signs. <i>Psychiatric Annals</i> , 2003, 33, 170-176.	0.1	21
27	Assessment for subtle neurological signs. , 0, , 333-343.		1