
## Shin-Liang Pan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/694340/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effectiveness of fecal immunochemical testing in reducing colorectal cancer mortality from the <scp>O</scp> ne <scp>M</scp> illion <scp>T</scp> aiwanese <scp>S</scp> creening <scp>P</scp> rogram. Cancer, 2015, 121, 3221-3229.	2.0	205
2	Circulating Levels of Markers of Inflammation and Endothelial Activation are Increased in Men with Chronic Spinal Cord Injury. Journal of the Formosan Medical Association, 2007, 106, 919-928.	0.8	102
3	Association of Diabetes Mellitus With the Risk of Developing Adhesive Capsulitis of the Shoulder: A Longitudinal Populationâ€Based Followup Study. Arthritis Care and Research, 2013, 65, 1197-1202.	1.5	70
4	Population-Based Breast Cancer Screening With Risk-Based and Universal Mammography Screening Compared With Clinical Breast Examination. JAMA Oncology, 2016, 2, 915.	3.4	62
5	Parkinson's Disease Is Related to an Increased Risk of Ischemic Stroke—A Population-Based Propensity Score-Matched Follow-Up Study. PLoS ONE, 2013, 8, e68314.	1.1	58
6	Intracerebral hemorrhage secondary to autonomic dysreflexia in a young person with incomplete C8 tetraplegia: A case report. Archives of Physical Medicine and Rehabilitation, 2005, 86, 591-593.	0.5	52
7	Predicting Length of Stay among Patients Discharged from the Emergency Department—Using an Accelerated Failure Time Model. PLoS ONE, 2017, 12, e0165756.	1.1	51
8	Sonographic evaluations in elite college baseball athletes. Scandinavian Journal of Medicine and Science in Sports, 2005, 15, 29-35.	1.3	48
9	Increased Risk of Hemorrhagic Stroke in Patients with Migraine: A Population-Based Cohort Study. PLoS ONE, 2013, 8, e55253.	1.1	45
10	Increased risk of trigeminal neuralgia after hypertension. Neurology, 2011, 77, 1605-1610.	1.5	43
11	Increased Risk of Ischemic Heart Disease in Young Patients with Newly Diagnosed Ankylosing Spondylitis – A Population-Based Longitudinal Follow-Up Study. PLoS ONE, 2013, 8, e64155.	1.1	40
12	Parkinson disease and risk of acute myocardial infarction: A population-based, propensity score–matched, longitudinal follow-up study. American Heart Journal, 2015, 169, 508-514.	1.2	38
13	Responsiveness of SF-36 and Lower Extremity Functional Scale for assessing outcomes in traumatic injuries of lower extremities. Injury, 2014, 45, 1759-1763.	0.7	35
14	Location and size of infarct on functional outcome of noncardioembolic ischemic stroke. Disability and Rehabilitation, 2006, 28, 977-983.	0.9	32
15	Increased Risk of Ischemic Stroke in Young Patients with Ankylosing Spondylitis: A Population-Based Longitudinal Follow-Up Study. PLoS ONE, 2014, 9, e94027.	1.1	28
16	Increased risk of ischemic heart disease in young patients with migraine: A population-based, propensity score-matched, longitudinal follow-up study. International Journal of Cardiology, 2014, 172, 213-216.	0.8	27
17	Increased risk of rheumatoid arthritis in patients with migraine: a population-based, propensity score-matched cohort study. Rheumatology International, 2017, 37, 273-279.	1.5	27
18	Predicting Recovery of Voluntary Upper Extremity Movement in Subacute Stroke Patients with Severe Upper Extremity Paresis. PLoS ONE, 2015, 10, e0126857.	1.1	25

Shin-Liang Pan

#	Article	IF	CITATIONS
19	Predictors of long-term survival after stroke in Taiwan. Journal of Rehabilitation Medicine, 2008, 40, 844-849.	0.8	23
20	A Markov regression randomâ€effects model for remission of functional disability in patients following a first stroke: A Bayesian approach. Statistics in Medicine, 2007, 26, 5335-5353.	0.8	22
21	ls Higher Serum Total Cholesterol Level Associated With Better Long-Term Functional Outcomes After Noncardioembolic Ischemic Stroke?. Archives of Physical Medicine and Rehabilitation, 2010, 91, 913-918.	0.5	21
22	Increased risk of stroke after trigeminal neuralgia – a population-based follow-up study. Cephalalgia, 2011, 31, 937-942.	1.8	21
23	Spinal cord injury and Alzheimer's disease risk: a population-based, retrospective cohort study. Spinal Cord, 2018, 56, 151-157.	0.9	21
24	Dynamic Aspect of Functional Recovery After Stroke Using a Multistate Model. Archives of Physical Medicine and Rehabilitation, 2008, 89, 1054-1060.	0.5	19
25	Migraine is related to an increased risk of Parkinson's disease: A population-based, propensity score-matched, longitudinal follow-up study. Cephalalgia, 2016, 36, 1316-1323.	1.8	19
26	Auditory Cortical Evoked Potentials in Tinnitus Patients with Normal Audiological Presentation. Journal of the Formosan Medical Association, 2007, 106, 979-985.	0.8	18
27	Effect of Functional Status on Survival in Patients With Stroke: Is Independent Ambulation a Key Determinant?. Archives of Physical Medicine and Rehabilitation, 2012, 93, 527-531.	0.5	18
28	Thresholds of Tone Burst Auditory Brainstem Responses for Infants and Young Children with Normal Hearing in Taiwan. Journal of the Formosan Medical Association, 2007, 106, 847-853.	0.8	12
29	Increased risk of stroke after Bell's palsy: a population-based longitudinal follow-up study: Table 1. Journal of Neurology, Neurosurgery and Psychiatry, 2012, 83, 341-343.	0.9	11
30	Spinal Cord Injury Is Related to an Increased Risk of Multiple Sclerosis: A Population-Based, Propensity Score-Matched, Longitudinal Follow-Up Study. Journal of Neurotrauma, 2015, 32, 655-659.	1.7	11
31	Osteoporosis and the Risk of Parkinson's Disease: A Nationwide, Propensity Score–Matched, Longitudinal Follow-up Study. Journal of Clinical Endocrinology and Metabolism, 2021, 106, e763-e771.	1.8	10
32	Reduction of disability after stroke is a more informative predictor of long-time survival than initial disability status. Disability and Rehabilitation, 2007, 29, 417-423.	0.9	9
33	Effects of Age and Degree of Hearing Loss on the Agreement and Correlation Between Sound Field Audiometric Thresholds and Tone Burst Auditory Brainstem Response Thresholds in Infants and Young Children. Journal of the Formosan Medical Association, 2008, 107, 869-875.	0.8	9
34	Does Adhesive Capsulitis of the Shoulder Increase the Risk of Stroke? A Population-Based Propensity Score-Matched Follow-Up Study. PLoS ONE, 2012, 7, e49343.	1,1	9
35	Does Non-Central Nervous System Tuberculosis Increase the Risk of Ischemic Stroke? A Population-Based Propensity Score-Matched Follow-Up Study. PLoS ONE, 2014, 9, e98158.	1.1	8
36	Time-varying Markov regression random-effect model with Bayesian estimation procedures: Application to dynamics of functional recovery in patients with stroke. Mathematical Biosciences, 2010, 227, 72-79.	0.9	7

Shin-Liang Pan

#	Article	IF	CITATIONS
37	Asymptomatic Median Mononeuropathy Among Men With Chronic Paraplegia. Archives of Physical Medicine and Rehabilitation, 2007, 88, 1193-1197.	0.5	5
38	Sublesional Spinal Vertebral Bone Mineral Density Correlates With Neurological Level and Body Mass Index in Individuals With Chronic Complete Spinal Cord Injuries. Spine, 2010, 35, 958-962.	1.0	5
39	Are higher total serum cholesterol levels associated with better long-term motor function after ischemic stroke?. Nutritional Neuroscience, 2012, 15, 239-243.	1.5	5
40	Reduced Sympathetic Skin Response in the Isolated Spinal Cord of Subjects With Spinal Cord Injury. Archives of Physical Medicine and Rehabilitation, 2006, 87, 1201-1206.	0.5	4
41	Myasthenia gravis with superimposed spinal cord injury: a case report. Journal of Rehabilitation Medicine, 2008, 40, 684-686.	0.8	4
42	Increased Risk of Acute Cholecystitis in Patients With Spinal Cord Injury. Spine, 2018, 43, 934-939.	1.0	4
43	Osteoarthritis is associated with an increased risk of Parkinson's disease: A populationâ€based, longitudinal followâ€up study. Arthritis Care and Research, 2021, , .	1.5	3
44	Quantifying Dynamic Flow of Emergency Department (ED) Patient Managements: A Multistate Model Approach. Emergency Medicine International, 2020, 2020, 1-9.	0.3	2
45	Impacts of treatments on recurrence and 28-year survival of ischemic stroke patients. Scientific Reports, 2021, 11, 15258.	1.6	2
46	Increased risk of carpal tunnel syndrome in people with spinal cord injury: A nationwide longitudinal follow-up study. Archives of Physical Medicine and Rehabilitation, 2021, , .	0.5	2
47	Osteoporosis and the long-term risk of peripheral artery disease: a population-based longitudinal follow-up study in Taiwan. Osteoporosis International, 2022, 33, 1117-1123.	1.3	2
48	Physical Activity and the Risk of Hemorrhagic Stroke: A Population-Based Longitudinal Follow-Up Study in Taiwan. Frontiers in Medicine, 2021, 8, 791772.	1.2	2
49	Increased risk of acute pancreatitis in persons with spinal cord injury: a population-based, propensity score-matched longitudinal follow-up study. Spinal Cord, 2021, 59, 1170-1176.	0.9	1
50	Increased risk of sensorineural hearing loss in patients with spinal cord injury: a nationwide longitudinal follow-up study. Spinal Cord, 2021, 59, 1200-1205.	0.9	1
51	Use of Bayesian Modeling to Estimate the Sensitivity of Stereotactic Directional Vacuum-Assisted Breast Biopsy When the Gold Standard is Incomplete. Academic Radiology, 2009, 16, 1316-1322.	1.3	0
52	Corrigendum to "Increased risk of ischemic heart disease in young patients with migraine: A population-based, propensity score-matched, longitudinal follow-up study―[Int J Cardiol 172 (2014) 213–216]. International Journal of Cardiology, 2014, 175, 393.	0.8	0
53	OUP accepted manuscript. International Journal of Epidemiology, 2022, , .	0.9	0