

# Seyyed Mohammad Reza Kazemi-Bajes

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6945712/publications.pdf>

Version: 2024-02-01

31  
papers

1,283  
citations

535685

17  
h-index

488211

31  
g-index

31  
all docs

31  
docs citations

31  
times ranked

2568  
citing authors

#	ARTICLE	IF	CITATIONS
1	Rapid atrophy of cardiac left ventricular mass in patients with non-small cell carcinoma of the lung. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2019, 10, 1070-1082.	2.9	21
2	The Effect of Platelet Rich Plasma on Hair Regrowth in Patients with Alopecia Areata Totalis: a Clinical Pilot Study. <i>Dermatologic Therapy</i> , 2019, 32, e12989.	0.8	14
3	Undiagnosed cardiac deficits in non-small cell carcinoma patients in the candidate population for anti-cachexia clinical trials. <i>Supportive Care in Cancer</i> , 2019, 27, 1551-1561.	1.0	9
4	High fat mass associates with occurrence of targeted therapy-induced left ventricular ejection fraction reduction in patients with renal cell carcinoma. <i>Clinical Nutrition</i> , 2018, 37, 1070-1072.	2.3	1
5	Adherence to a Dash-style diet in relation to depression and aggression in adolescent girls. <i>Psychiatry Research</i> , 2018, 259, 104-109.	1.7	28
6	Dietary behaviors in relation to prevalence of irritable bowel syndrome in adolescent girls. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018, 33, 404-410.	1.4	29
7	Depression and anxiety both associate with serum level of hs-CRP: A gender-stratified analysis in a population-based study. <i>Psychoneuroendocrinology</i> , 2017, 81, 63-69.	1.3	95
8	Serum high C reactive protein concentrations are related to the intake of dietary macronutrients and fiber: Findings from a large representative Persian population sample. <i>Clinical Biochemistry</i> , 2017, 50, 750-755.	0.8	19
9	The prevalence of metabolic syndrome increases with serum high sensitivity C-reactive protein concentration in individuals without a history of cardiovascular disease: a report from a large Persian cohort. <i>Annals of Clinical Biochemistry</i> , 2017, 54, 644-648.	0.8	13
10	Rheumatoid Arthritis Disadvantages Younger Patients for Cardiovascular Diseases: A Meta-Analysis. <i>PLoS ONE</i> , 2016, 11, e0157360.	1.1	43
11	Concurrent depletion of skeletal muscle, fat, and left ventricular mass in patients with cirrhosis of the liver. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2016, 7, 97-99.	2.9	13
12	Development of a validated algorithm for the diagnosis of paediatric asthma in electronic medical records. <i>Npj Primary Care Respiratory Medicine</i> , 2016, 26, 16085.	1.1	14
13	Computed tomography-defined muscle and fat wasting are associated with cancer clinical outcomes. <i>Seminars in Cell and Developmental Biology</i> , 2016, 54, 2-10.	2.3	227
14	Obesity Paradox versus Frailty Syndrome in First-Ever Ischemic Stroke Survivors. <i>International Journal of Stroke</i> , 2015, 10, E75-E75.	2.9	2
15	Concurrent evolution of cancer cachexia and heart failure: bilateral effects exist. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2014, 5, 95-104.	2.9	62
16	Cardiovascular Risk Factors and Nutritional Intake are not Associated with Ultrasound-defined Increased Carotid Intima Media Thickness in Individuals Without a History of Cardiovascular Events. <i>International Journal of Preventive Medicine</i> , 2014, 5, 1412-21.	0.2	5
17	Clinical outcomes related to muscle mass in humans with cancer and catabolic illnesses. <i>International Journal of Biochemistry and Cell Biology</i> , 2013, 45, 2302-2308.	1.2	120
18	A High Prevalence of Carotid Artery Stenosis in Male Patients Older Than 65 Years, Irrespective of Presenting Clinical Manifestation of Atherosclerotic Diseases. <i>Angiology</i> , 2013, 64, 281-286.	0.8	3

#	ARTICLE	IF	CITATIONS
19	Concept of atherosclerosis velocity: is it a better measure of cardiovascular risk?. Iranian Journal of Medical Sciences, 2013, 38, 210-20.	0.3	1
20	Lunar phase cycle and psychiatric hospital emergency visits, inpatient admissions and aggressive behavior. Asian Journal of Psychiatry, 2011, 4, 45-50.	0.9	14
21	Changes in pro-oxidant-antioxidant balance after bare metal and drug eluting stent implantation in patients with stable coronary disease. Clinical Biochemistry, 2011, 44, 160-164.	0.8	8
22	Serum Inflammatory and Immune Marker Response After Bare-Metal or Drug-Eluting Stent Implantation Following Percutaneous Coronary Intervention. Angiology, 2011, 62, 184-190.	0.8	10
23	Metabolic Syndrome May Not be a Good Predictor of Coronary Artery Disease in the Iranian Population: Population-Specific Definitions are Required. Scientific World Journal, The, 2009, 9, 86-96.	0.8	23
24	Serum Selenium and Glutathione Peroxidase Concentrations in Iranian Patients with Angiography-Defined Coronary Artery Disease. Angiology, 2009, 60, 186-191.	0.8	17
25	The relationship between established coronary risk factors and serum copper and zinc concentrations in a large Persian Cohort. Journal of Trace Elements in Medicine and Biology, 2009, 23, 167-175.	1.5	65
26	Omega-3 fatty acid supplements improve the cardiovascular risk profile of subjects with metabolic syndrome, including markers of inflammation and auto-immunity. Acta Cardiologica, 2009, 64, 321-327.	0.3	102
27	Prooxidant-antioxidant balance as a new risk factor in patients with angiographically defined coronary artery disease. Clinical Biochemistry, 2008, 41, 375-380.	0.8	110
28	Natural Honey and Cardiovascular Risk Factors; Effects on Blood Glucose, Cholesterol, Triacylglycerole, CRP, and Body Weight Compared with Sucrose. Scientific World Journal, The, 2008, 8, 463-469.	0.8	134
29	Serum zinc and copper status in dyslipidaemic patients with and without established coronary artery disease. Clinical Laboratory, 2008, 54, 321-9.	0.2	18
30	Serum copper and zinc concentrations are lower in Iranian patients with angiographically defined coronary artery disease than in subjects with a normal angiogram. Journal of Trace Elements in Medicine and Biology, 2007, 21, 22-28.	1.5	42
31	C-reactive protein associated with coronary artery disease in Iranian patients with angiographically defined coronary artery disease. Clinical Laboratory, 2007, 53, 49-56.	0.2	21