

# May Arna Risberg

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/6946386/may-arna-risberg-publications-by-year.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

117  
papers

7,196  
citations

47  
h-index

83  
g-index

127  
ext. papers

8,260  
ext. citations

5.3  
avg, IF

6.06  
L-index

#	Paper	IF	Citations
117	Relationship between cam morphology, hip symptoms, and hip osteoarthritis: the Musculoskeletal pain in Ullersaker Study (MUST) cohort. <i>HIP International</i> , <b>2021</b> , 31, 789-796	1.7	2
116	Clinical, Functional, and Physical Activity Outcomes 5 Years Following the Treatment Algorithm of the Delaware-Oslo ACL Cohort Study. <i>Journal of Bone and Joint Surgery - Series A</i> , <b>2021</b> , 103, 1473-1481	5.6	3
115	On a Trajectory for Success-9 in Every 10 People With a Degenerative Meniscus Tear Have Improved Knee Function Within 2 Years After Treatment: A Secondary Exploratory Analysis of a Randomized Controlled Trial. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2021</b> , 51, 289-297	4.2	2
114	Infographic. Consensus recommendations on the classification, definition and diagnostic criteria of hip-related pain in young and middle-aged active adults from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2021</b> , 55, 115-117	10.3	1
113	Low Rates of Radiographic Knee Osteoarthritis 5 Years After ACL Reconstruction or Rehabilitation Alone: The Delaware-Oslo ACL Cohort Study. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2021</b> , 9, 23259671211027530	3.5	1
112	Assessing implementation, limited efficacy, and acceptability of the BEAST tool: A rehabilitation and return-to-sport decision tool for nonprofessional athletes with anterior cruciate ligament reconstruction. <i>Physical Therapy in Sport</i> , <b>2021</b> , 52, 147-154	3	
111	Comparing the Responsiveness of the Global Rating Scale With Legacy Knee Outcome Scores: A Delaware-Oslo Cohort Study. <i>American Journal of Sports Medicine</i> , <b>2020</b> , 48, 1953-1960	6.8	1
110	Patient-reported outcome measures for hip-related pain: a review of the available evidence and a consensus statement from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 848-857	10.3	36
109	Activity and functional readiness, not age, are the critical factors for second anterior cruciate ligament injury - the Delaware-Oslo ACL cohort study. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 1099-1102	10.3	23
108	Does surgery reduce knee osteoarthritis, meniscal injury and subsequent complications compared with non-surgery after ACL rupture with at least 10 years follow-up? A systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 592-598	10.3	23
107	Standardised measurement of physical capacity in young and middle-aged active adults with hip-related pain: recommendations from the first International Hip-related Pain Research Network (IHiPRN) meeting, Zurich, 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 702-710	10.3	12
106	Physiotherapist-led treatment for young to middle-aged active adults with hip-related pain: consensus recommendations from the International Hip-related Pain Research Network, Zurich 2018. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 504-511	10.3	18
105	Evaluating plasma extracellular vesicle microRNAs as possible biomarkers for osteoarthritis. <i>Osteoarthritis and Cartilage Open</i> , <b>2020</b> , 1, 100018	1.5	3
104	The Active with OsteoArthritis (AktivA) Physiotherapy Implementation Model: A Patient Education, Supervised Exercise and Self-Management Program for Patients with Mild to Moderate Osteoarthritis of the Knee or Hip Joint. A National Register Study with a Two-Year Follow-Up. <i>Journal of Clinical Medicine</i> , <b>2020</b> , 9, 100018	5.1	3
103	Is good muscle function a protective factor for early signs of knee osteoarthritis after anterior cruciate ligament reconstruction? The SHIELD cohort study protocol. <i>Osteoarthritis and Cartilage Open</i> , <b>2020</b> , 2, 100102	1.5	1
102	Association of Quadriceps Strength Symmetry and Surgical Status with Clinical Osteoarthritis 5 Years after Anterior Cruciate Ligament Rupture. <i>Arthritis Care and Research</i> , <b>2020</b> ,	4.7	10
101	Meniscus or Cartilage Injury at the Time of Anterior Cruciate Ligament Tear Is Associated With Worse Prognosis for Patient-Reported Outcome 2 to 10 Years After Anterior Cruciate Ligament Injury: A Systematic Review. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2020</b> , 50, 490-502	4.2	6

100	Coper Classification Early After ACL Rupture Changes With Progressive Neuromuscular and Strength Training and Is Associated With 2-Year Success: Response. <i>American Journal of Sports Medicine</i> , <b>2019</b> , 47, NP65-NP66	6.8	
99	Establishing outcome measures in early knee osteoarthritis. <i>Nature Reviews Rheumatology</i> , <b>2019</b> , 15, 438-448	8.1	50
98	The Maternal and Paternal Effects on Clinically and Surgically Defined Osteoarthritis. <i>Arthritis and Rheumatology</i> , <b>2019</b> , 71, 1844-1848	9.5	4
97	Knee Pathology in Young Adults After Pediatric Anterior Cruciate Ligament Injury: A Prospective Case Series of 47 Patients With a Mean 9.5-Year Follow-up. <i>American Journal of Sports Medicine</i> , <b>2019</b> , 47, 1557-1566	6.8	12
96	What's the rate of knee osteoarthritis 10 years after anterior cruciate ligament injury? An updated systematic review. <i>British Journal of Sports Medicine</i> , <b>2019</b> , 53, 1162-1167	10.3	70
95	Coper Classification Early After Anterior Cruciate Ligament Rupture Changes With Progressive Neuromuscular and Strength Training and Is Associated With 2-Year Success: The Delaware-Oslo ACL Cohort Study. <i>American Journal of Sports Medicine</i> , <b>2019</b> , 47, 807-814	6.8	21
94	Superior 2-Year Functional Outcomes Among Young Female Athletes After ACL Reconstruction in 10 Return-to-Sport Training Sessions: Comparison of ACL-SPORTS Randomized Controlled Trial With Delaware-Oslo and MOON Cohorts. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2019</b> , 7, 2325967119861311	3.5	16
93	Complex Tears, Extrusion, and Larger Excision Are Prognostic Factors for Worse Outcomes 1 and 2 Years After Arthroscopic Partial Meniscectomy for Degenerative Meniscal Tears: A Secondary Exploratory Study of the Surgically Treated Group From the Odense-Oslo Meniscectomy Versus Exercise (OMEX) Trial. <i>Arthroscopy</i> , <b>2019</b> , 47, 2478-2487	6.8	9
92	Efficacy of periacetabular osteotomy followed by progressive resistance training compared to progressive resistance training as non-surgical treatment in patients with hip dysplasia (PreserveHip) - a protocol for a randomised controlled trial. <i>BMJ Open</i> , <b>2019</b> , 9, e032782	3	4
91	Better before-better after: efficacy of prehabilitation for older patients with osteoarthritis awaiting total hip replacement-a study protocol for a randomised controlled trial in South-Eastern Norway. <i>BMJ Open</i> , <b>2019</b> , 9, e031626	3	2
90	A Pragmatic Approach to the Implementation of Osteoarthritis Guidelines Has Fewer Potential Barriers Than Recommended Implementation Frameworks. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2019</b> , 49, 1-4	4.2	11
89	Coping With Anterior Cruciate Ligament Injury From Childhood to Maturation: A Prospective Case Series of 44 Patients With Mean 8 Years' Follow-up. <i>American Journal of Sports Medicine</i> , <b>2019</b> , 47, 22-30	6.8	19
88	The 6-m timed hop test is a prognostic factor for outcomes in patients with meniscal tears treated with exercise therapy or arthroscopic partial meniscectomy: a secondary, exploratory analysis of the Odense-Oslo meniscectomy versus exercise (OMEX) trial. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2019</b> , 27, 2478-2487	5.5	5
87	Arthroscopic Surgical Procedures Versus Sham Surgery for Patients With Femoroacetabular Impingement and/or Labral Tears: Study Protocol for a Randomized Controlled Trial (HIPARTI) and a Prospective Cohort Study (HARP). <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2018</b> , 48, 325-335	4.2	16
86	Impairment-Based Rehabilitation Following Hip Arthroscopy: Postoperative Protocol for the HIP ARThroscopy International Randomized Controlled Trial. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2018</b> , 48, 336-342	4.2	13
85	Return to pivoting sport after ACL reconstruction: association with osteoarthritis and knee function at the 15-year follow-up. <i>British Journal of Sports Medicine</i> , <b>2018</b> , 52, 1199-1204	10.3	25
84	Functional performance 6 months after ACL reconstruction can predict return to participation in the same preinjury activity level 12 and 24 months after surgery. <i>British Journal of Sports Medicine</i> , <b>2018</b> , 52, 375	10.3	57
83	No Difference in the KOOS Quality of Life Subscore Between Anatomic Double-Bundle and Anatomic Single-Bundle Anterior Cruciate Ligament Reconstruction of the Knee: A Prospective Randomized Controlled Trial With 2 Years' Follow-up. <i>American Journal of Sports Medicine</i> , <b>2018</b> , 46, 2341-2354	6.8	22

82	Changes in the rate of publicly financed knee arthroscopies: an analysis of data from the Norwegian patient registry from 2012 to 2016. <i>BMJ Open</i> , <b>2018</b> , 8, e021199	3	8
81	Anterior Cruciate Ligament Injury-Who Succeeds Without Reconstructive Surgery? The Delaware-Oslo ACL Cohort Study. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2018</b> , 6, 2325967118774255	3.5	22
80	Do Patients Failing Return-to-Activity Criteria at 6 Months After Anterior Cruciate Ligament Reconstruction Continue Demonstrating Deficits at 2 Years?. <i>American Journal of Sports Medicine</i> , <b>2017</b> , 45, 1037-1048	6.8	52
79	Distribution of osteoarthritis in a Norwegian population-based cohort: associations to risk factor profiles and health-related quality of life. <i>Rheumatology International</i> , <b>2017</b> , 37, 1541-1550	3.6	5
78	Middle-aged patients with an MRI-verified medial meniscal tear report symptoms commonly associated with knee osteoarthritis. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , <b>2017</b> , 88, 664-669	4.3	16
77	We Need to Implement Current Evidence in Early Rehabilitation Programs to Improve Long-Term Outcome After Anterior Cruciate Ligament Injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2016</b> , 46, 710-3	4.2	26
76	Does Extended Preoperative Rehabilitation Influence Outcomes 2 Years After ACL Reconstruction? A Comparative Effectiveness Study Between the MOON and Delaware-Oslo ACL Cohorts. <i>American Journal of Sports Medicine</i> , <b>2016</b> , 44, 2608-2614	6.8	82
75	Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. <i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 804-8	10.3	547
74	Patients With Chondrolabral Pathology Have Bilateral Functional Impairments 12 to 24 Months After Unilateral Hip Arthroscopy: A Cross-sectional Study. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2016</b> , 46, 947-956	4.2	28
73	Evaluation of focal cartilage lesions of the knee using MRI T2 mapping and delayed Gadolinium Enhanced MRI of Cartilage (dGEMRIC). <i>BMC Musculoskeletal Disorders</i> , <b>2016</b> , 17, 73	2.8	11
72	Changes in Knee Osteoarthritis, Symptoms, and Function After Anterior Cruciate Ligament Reconstruction: A 20-Year Prospective Follow-up Study. <i>American Journal of Sports Medicine</i> , <b>2016</b> , 44, 1215-24	6.8	109
71	Long-Term Effect of Exercise Therapy and Patient Education on Impairments and Activity Limitations in People With Hip Osteoarthritis: Secondary Outcome Analysis of a Randomized Clinical Trial. <i>Physical Therapy</i> , <b>2016</b> , 96, 818-27	3.3	17
70	Authors' reply to Joshi. <i>BMJ, The</i> , <b>2016</b> , 354, i4623	5.9	1
69	Physical impairments in symptomatic femoroacetabular impingement: a systematic review of the evidence. <i>British Journal of Sports Medicine</i> , <b>2016</b> , 50, 1180	10.3	79
68	Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. <i>BMJ, The</i> , <b>2016</b> , 354, i3740	5.9	133
67	Quantifying Quadriceps Muscle Strength in Patients With ACL Injury, Focal Cartilage Lesions, and Degenerative Meniscus Tears: Differences and Clinical Implications. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2016</b> , 4, 2325967116667717	3.5	11
66	Defining the presence of radiographic knee osteoarthritis: a comparison between the Kellgren and Lawrence system and OARSI atlas criteria. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2015</b> , 23, 3532-9	5.5	51
65	Exercise therapy may postpone total hip replacement surgery in patients with hip osteoarthritis: a long-term follow-up of a randomised trial. <i>Annals of the Rheumatic Diseases</i> , <b>2015</b> , 74, 164-9	2.4	70

64	Effect of exercise therapy compared with arthroscopic surgery on knee muscle strength and functional performance in middle-aged patients with degenerative meniscus tears: a 3-mo follow-up of a randomized controlled trial. <i>American Journal of Physical Medicine and Rehabilitation</i> , <b>2015</b> , 94, 460-73	2.6	37
63	Gait Characteristics, Symptoms, and Function in Persons With Hip Osteoarthritis: A Longitudinal Study With 6 to 7 Years of Follow-up. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2015</b> , 45, 539-49	4.2	10
62	Inter- and intrarater reliability of four single-legged hop tests and isokinetic muscle torque measurements in children. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2015</b> , 23, 1907-16	5.5	20
61	Consensus criteria for defining 'successful outcome' after ACL injury and reconstruction: a Delaware-Oslo ACL cohort investigation. <i>British Journal of Sports Medicine</i> , <b>2015</b> , 49, 335-42	10.3	170
60	Exercises including weight vests and a patient education program for women with osteopenia: a feasibility study of the OsteoACTIVE rehabilitation program. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2015</b> , 45, 97-105, C1-4	4.2	5
59	Upper limb physical function and adverse effects after breast cancer surgery: a prospective 2.5-year follow-up study and preoperative measures. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2014</b> , 95, 875-81	2.8	67
58	Self-reported knee function can identify athletes who fail return-to-activity criteria up to 1 year after anterior cruciate ligament reconstruction: a delaware-oslo ACL cohort study. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2014</b> , 44, 914-23	4.2	99
57	Nonsurgical or Surgical Treatment of ACL Injuries: Knee Function, Sports Participation, and Knee Reinjury: The Delaware-Oslo ACL Cohort Study. <i>Journal of Bone and Joint Surgery - Series A</i> , <b>2014</b> , 96, 1233-1241	5.6	105
56	Weight-bearing asymmetries during Sit-To-Stand in patients with mild-to-moderate hip osteoarthritis. <i>Gait and Posture</i> , <b>2014</b> , 39, 683-8	2.6	31
55	Knee function and knee muscle strength in middle-aged patients with degenerative meniscal tears eligible for arthroscopic partial meniscectomy. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 784-8	10.3	23
54	Online registration of monthly sports participation after anterior cruciate ligament injury: a reliability and validity study. <i>British Journal of Sports Medicine</i> , <b>2014</b> , 48, 748-53	10.3	21
53	The prevalence of patellofemoral osteoarthritis 12 years after anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2013</b> , 21, 942-9	5.5	61
52	The feasibility of a 3-month active rehabilitation program for patients with knee full-thickness articular cartilage lesions: the Oslo Cartilage Active Rehabilitation and Education Study. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2013</b> , 43, 310-24	4.2	27
51	Prevalence and incidence of new meniscus and cartilage injuries after a nonoperative treatment algorithm for ACL tears in skeletally immature children: a prospective MRI study. <i>American Journal of Sports Medicine</i> , <b>2013</b> , 41, 1771-9	6.8	80
50	Functional outcomes following a non-operative treatment algorithm for anterior cruciate ligament injuries in skeletally immature children 12 years and younger. A prospective cohort with 2 years follow-up. <i>British Journal of Sports Medicine</i> , <b>2013</b> , 47, 488-94	10.3	72
49	A 12-week exercise therapy program in middle-aged patients with degenerative meniscus tears: a case series with 1-year follow-up. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2012</b> , 42, 919-31	4.2	38
48	Management of anterior cruciate ligament injuries in skeletally immature individuals. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2012</b> , 42, 172-83	4.2	34
47	Ultrasound-guided sclerosis of neovessels in patellar tendinopathy: a prospective study of 101 patients. <i>American Journal of Sports Medicine</i> , <b>2012</b> , 40, 542-7	6.8	43

46	Inter- and intrarater reliability of isokinetic thigh muscle strength tests in postmenopausal women with osteopenia. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2012</b> , 93, 420-7	2.8	11
45	Patients with focal full-thickness cartilage lesions benefit less from ACL reconstruction at 2-5 years follow-up. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2012</b> , 20, 1533-9	5.5	36
44	No differences in prevalence of osteoarthritis or function after open versus endoscopic technique for anterior cruciate ligament reconstruction: 12-year follow-up report of a randomized controlled trial. <i>American Journal of Sports Medicine</i> , <b>2012</b> , 40, 2492-8	6.8	31
43	Single-legged hop tests as predictors of self-reported knee function after anterior cruciate ligament reconstruction: the Delaware-Oslo ACL cohort study. <i>American Journal of Sports Medicine</i> , <b>2012</b> , 40, 2348-56	6.8	206
42	A pair-matched comparison of return to pivoting sports at 1 year in anterior cruciate ligament-injured patients after a nonoperative versus an operative treatment course. <i>American Journal of Sports Medicine</i> , <b>2012</b> , 40, 2509-16	6.8	91
41	Muscle strength and hop performance criteria prior to return to sports after ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2011</b> , 19, 1798-805	5.5	265
40	Single-legged hop tests as predictors of self-reported knee function in nonoperatively treated individuals with anterior cruciate ligament injury. <i>American Journal of Sports Medicine</i> , <b>2011</b> , 39, 2347-54	6.8	104
39	No difference in knee function or prevalence of osteoarthritis after reconstruction of the anterior cruciate ligament with 4-strand hamstring autograft versus patellar tendon-bone autograft: a randomized study with 10-year follow-up. <i>American Journal of Sports Medicine</i> , <b>2010</b> , 38, 448-54	6.8	163
38	A progressive 5-week exercise therapy program leads to significant improvement in knee function early after anterior cruciate ligament injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2010</b> , 40, 705-21	4.2	153
37	Development of a therapeutic exercise program for patients with osteoarthritis of the hip. <i>Physical Therapy</i> , <b>2010</b> , 90, 592-601	3.3	28
36	Anterior cruciate ligament-deficient potential copers and noncopers reveal different isokinetic quadriceps strength profiles in the early stage after injury. <i>American Journal of Sports Medicine</i> , <b>2010</b> , 38, 586-93	6.8	62
35	Knee function and prevalence of knee osteoarthritis after anterior cruciate ligament reconstruction: a prospective study with 10 to 15 years of follow-up. <i>American Journal of Sports Medicine</i> , <b>2010</b> , 38, 2201-10	6.8	325
34	Functioning and disability in patients with hip osteoarthritis with mild to moderate pain. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2010</b> , 40, 616-24	4.2	39
33	No effect of group-based aerobic interval training on N-terminal pro-B-type natriuretic peptide levels in patients with chronic heart failure. <i>Scandinavian Cardiovascular Journal</i> , <b>2010</b> , 44, 223-9	2	15
32	Full-thickness cartilage lesion do not affect knee function in patients with ACL injury. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2010</b> , 18, 298-303	5.5	17
31	Functional tests should be accentuated more in the decision for ACL reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2010</b> , 18, 1517-25	5.5	44
30	Quadriceps muscle weakness after anterior cruciate ligament reconstruction: a risk factor for knee osteoarthritis?. <i>Arthritis Care and Research</i> , <b>2010</b> , 62, 1706-14	4.7	61
29	Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. <i>American Journal of Sports Medicine</i> , <b>2009</b> , 37, 1434-43	6.8	597

28	Outcome after knee dislocations: a 2-9 years follow-up of 85 consecutive patients. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2009</b> , 17, 1013-26	5.5	134
27	The long-term effect of 2 postoperative rehabilitation programs after anterior cruciate ligament reconstruction: a randomized controlled clinical trial with 2 years of follow-up. <i>American Journal of Sports Medicine</i> , <b>2009</b> , 37, 1958-66	6.8	76
26	Validity for the simplified water displacement instrument to measure arm lymphedema as a result of breast cancer surgery. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2009</b> , 90, 803-9	2.8	43
25	Changes in arm morbidities and health-related quality of life after breast cancer surgery - a five-year follow-up study. <i>Acta Oncologica</i> , <b>2009</b> , 48, 1111-8	3.2	57
24	Physical activity for the affected limb and arm lymphedema after breast cancer surgery. A prospective, randomized controlled trial with two years follow-up. <i>Acta Oncologica</i> , <b>2009</b> , 48, 1102-10	3.2	99
23	Arthroscopic surgery provides no additional benefit over physiotherapy and medication for the treatment of knee osteoarthritis. <i>Australian Journal of Physiotherapy</i> , <b>2009</b> , 55, 137		7
22	Long-term effects of a group-based high-intensity aerobic interval-training program in patients with chronic heart failure. <i>American Journal of Cardiology</i> , <b>2008</b> , 102, 1220-4	3	54
21	Effects of group-based high-intensity aerobic interval training in patients with chronic heart failure. <i>American Journal of Cardiology</i> , <b>2008</b> , 102, 1361-5	3	63
20	Short-term beneficial effects of physiotherapy exercise after knee arthroplasty for osteoarthritis. <i>Australian Journal of Physiotherapy</i> , <b>2008</b> , 54, 73		
19	Group-based aerobic interval training in patients with chronic heart failure: Norwegian Ullevaal Model. <i>Physical Therapy</i> , <b>2008</b> , 88, 523-35	3.3	27
18	Individuals with an anterior cruciate ligament-deficient knee classified as noncopers may be candidates for nonsurgical rehabilitation. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2008</b> , 38, 586-95	4.2	81
17	Performance-based functional outcome for children 12 years or younger following anterior cruciate ligament injury: a two to nine-year follow-up study. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , <b>2008</b> , 16, 214-23	5.5	54
16	Neuromuscular training versus strength training during first 6 months after anterior cruciate ligament reconstruction: a randomized clinical trial. <i>Physical Therapy</i> , <b>2007</b> , 87, 737-50	3.3	170
15	Warm-up exercise prevents acute knee and ankle injuries in young handball players. <i>Australian Journal of Physiotherapy</i> , <b>2005</b> , 51, 131		3
14	Outpatient physical therapy influences the patients health-related quality of life. <i>Advances in Physiotherapy</i> , <b>2005</b> , 7, 40-47		6
13	The reliability of a simplified water displacement instrument: a method for measuring arm volume. <i>Archives of Physical Medicine and Rehabilitation</i> , <b>2005</b> , 86, 86-9	2.8	16
12	A systematic review of evidence for anterior cruciate ligament rehabilitation: how much and what type?. <i>Physical Therapy in Sport</i> , <b>2004</b> , 5, 125-145	3	105
11	Effect of neuromuscular training on proprioception, balance, muscle strength, and lower limb function in female team handball players. <i>Clinical Journal of Sport Medicine</i> , <b>2004</b> , 14, 88-94	3.2	146

10	Dynamic knee stability after anterior cruciate ligament rupture. <i>Exercise and Sport Sciences Reviews</i> , <b>2003</b> , 31, 195-200	6.7	25
9	The effect of bracing on proprioception of knees with anterior cruciate ligament injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2002</b> , 32, 11-5	4.2	54
8	A new device to measure knee laxity during weightbearing and non-weightbearing conditions. <i>Journal of Orthopaedic Research</i> , <b>2001</b> , 19, 1185-91	3.8	52
7	Four-strand hamstring tendon autograft compared with patellar tendon-bone autograft for anterior cruciate ligament reconstruction. A randomized study with two-year follow-up. <i>American Journal of Sports Medicine</i> , <b>2001</b> , 29, 722-8	6.8	350
6	Design and implementation of a neuromuscular training program following anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>2001</b> , 31, 620-31	4.2	102
5	Muscle strength recovery following anterior cruciate ligament reconstruction: A prospective study of 151 patients with a two-year follow-up. <i>Isokinetics and Exercise Science</i> , <b>2000</b> , 8, 57-63	0.6	8
4	Prospective study of changes in impairments and disabilities after anterior cruciate ligament reconstruction. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>1999</b> , 29, 400-12	4.2	125
3	The effect of knee bracing after anterior cruciate ligament reconstruction. A prospective, randomized study with two years' follow-up. <i>American Journal of Sports Medicine</i> , <b>1999</b> , 27, 76-83	6.8	119
2	Evaluation of knee joint laxity and the structural properties of the anterior cruciate ligament graft in the human. A case report. <i>American Journal of Sports Medicine</i> , <b>1997</b> , 25, 203-6	6.8	38
1	Assessment of functional tests after anterior cruciate ligament surgery. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , <b>1994</b> , 19, 212-7	4.2	106