

Li-Xin Wei

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6953702/li-xin-wei-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

133
papers

7,053
citations

36
h-index

82
g-index

135
ext. papers

7,950
ext. citations

6.7
avg, IF

5.09
L-index

#	Paper	IF	Citations
133	Extracellular microparticles derived from hepatic progenitor cells deliver a death signal to hepatoma-initiating cells.. <i>Journal of Nanobiotechnology</i> , 2022 , 20, 79	9.4	
132	LPS/Bcl3/YAP1 signaling promotes Sox9HNF4 β hepatocyte-mediated liver regeneration after hepatectomy.. <i>Cell Death and Disease</i> , 2022 , 13, 277	9.8	
131	Single cell transcriptional diversity and intercellular crosstalk of human liver cancer.. <i>Cell Death and Disease</i> , 2022 , 13, 261	9.8	0
130	Efficacy and Safety of TACE Combined With Lenvatinib Plus PD-1 Inhibitors Compared With TACE Alone for Unresectable Hepatocellular Carcinoma Patients: A Prospective Cohort Study.. <i>Frontiers in Oncology</i> , 2022 , 12, 874473	5.3	0
129	Autophagy Is Required for Hepatic Differentiation of Hepatic Progenitor Cells via Wnt Signaling Pathway. <i>BioMed Research International</i> , 2021 , 2021, 6627506	3	1
128	Autophagy deficiency downregulates Omethylguanine-DNA methyltransferase and increases chemosensitivity of liver cancer cells. <i>Aging</i> , 2021 , 13, 14289-14303	5.6	0
127	The stemness of hepatocytes is maintained by high levels of lipopolysaccharide via YAP1 activation. <i>Stem Cell Research and Therapy</i> , 2021 , 12, 342	8.3	0
126	A TNFR2-hnRNPK Axis Promotes Primary Liver Cancer Development via Activation of YAP Signaling in Hepatic Progenitor Cells. <i>Cancer Research</i> , 2021 , 81, 3036-3050	10.1	4
125	Oncostatin M promotes hepatic progenitor cell activation and hepatocarcinogenesis via macrophage-derived tumor necrosis factor- α <i>Cancer Letters</i> , 2021 , 517, 46-54	9.9	7
124	An RNA-RNA crosstalk network involving HMGB1 and RICTOR facilitates hepatocellular carcinoma tumorigenesis by promoting glutamine metabolism and impedes immunotherapy by PD-L1+ exosomes activity.. <i>Signal Transduction and Targeted Therapy</i> , 2021 , 6, 421	21	3
123	Sirt1-Overexpressing Mesenchymal Stem Cells Drive the Anti-tumor Effect through Their Pro-inflammatory Capacity. <i>Molecular Therapy</i> , 2020 , 28, 874-888	11.7	7
122	Correction: Kupffer cells-dependent inflammation in the injured liver increases recruitment of mesenchymal stem cells in aging mice. <i>Oncotarget</i> , 2020 , 11, 3805-3806	3.3	
121	Autophagy and Tumorigenesis. <i>Advances in Experimental Medicine and Biology</i> , 2020 , 1207, 275-299	3.6	5
120	Autophagy and Tumour Stem Cells. <i>Advances in Experimental Medicine and Biology</i> , 2020 , 1207, 301-313	3.6	1
119	Autophagy and Tumour Metastasis. <i>Advances in Experimental Medicine and Biology</i> , 2020 , 1207, 315-338	3.6	4
118	Autophagy and Tumour Chemotherapy. <i>Advances in Experimental Medicine and Biology</i> , 2020 , 1207, 351-374	3.6	4
117	Autophagy and Tumour Radiotherapy. <i>Advances in Experimental Medicine and Biology</i> , 2020 , 1207, 375-387	3.6	5

116	The concentration of tumor necrosis factor- α determines its protective or damaging effect on liver injury by regulating Yap activity. <i>Cell Death and Disease</i> , 2020 , 11, 70	9.8	18
115	Mesenchymal stem cell therapy for liver disease: full of chances and challenges. <i>Cell and Bioscience</i> , 2020 , 10, 123	9.8	18
114	Ribosomal protein L34 is a potential prognostic biomarker and therapeutic target in hilar cholangiocarcinoma. <i>Cell and Bioscience</i> , 2020 , 10,	9.8	1
113	Lipopolysaccharide induces the differentiation of hepatic progenitor cells into myofibroblasts constitutes the hepatocarcinogenesis-associated microenvironment. <i>Cell Death and Differentiation</i> , 2020 , 27, 85-101	12.7	12
112	Macrophages and hepatocellular carcinoma. <i>Cell and Bioscience</i> , 2019 , 9, 79	9.8	42
111	BabaoDan attenuates high-fat diet-induced non-alcoholic fatty liver disease via activation of AMPK signaling. <i>Cell and Bioscience</i> , 2019 , 9, 77	9.8	22
110	Babao Dan attenuates acute ethanol-induced liver injury via Nrf2 activation and autophagy. <i>Cell and Bioscience</i> , 2019 , 9, 80	9.8	8
109	Inhibition of Autophagy with Chloroquine Enhanced Sinoporphyrin Sodium Mediated Photodynamic Therapy-induced Apoptosis in Human Colorectal Cancer Cells. <i>International Journal of Biological Sciences</i> , 2019 , 15, 12-23	11.2	19
108	A Pretreatment CT Model Predicts Survival Following Chemolipiodolization in Patients With Hepatocellular Carcinoma. <i>Technology in Cancer Research and Treatment</i> , 2019 , 18, 1533033819844488	2.7	2
107	Lipopolysaccharide protects against acetaminophen-induced hepatotoxicity by reducing oxidative stress via the TNF- α /TNFR1 pathway. <i>Biochemical and Biophysical Research Communications</i> , 2019 , 513, 623-630	3.4	4
106	Glycochenodeoxycholate promotes hepatocellular carcinoma invasion and migration by AMPK/mTOR dependent autophagy activation. <i>Cancer Letters</i> , 2019 , 454, 215-223	9.9	35
105	The distinct roles of mesenchymal stem cells in the initial and progressive stage of hepatocarcinoma. <i>Cell Death and Disease</i> , 2018 , 9, 345	9.8	20
104	Hippo Cascade Controls Lineage Commitment of Liver Tumors in Mice and Humans. <i>American Journal of Pathology</i> , 2018 , 188, 995-1006	5.8	20
103	Inhibition of DNMT suppresses the stemness of colorectal cancer cells through down-regulating Wnt signaling pathway. <i>Cellular Signalling</i> , 2018 , 47, 79-87	4.9	17
102	Enhanced doxorubicin delivery to hepatocellular carcinoma cells via CD147 antibody-conjugated immunoliposomes. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2018 , 14, 1949-1961	6	32
101	Immune response involved in liver damage and the activation of hepatic progenitor cells during liver tumorigenesis. <i>Cellular Immunology</i> , 2018 , 326, 52-59	4.4	18
100	Cancer nanomedicine: mechanisms, obstacles and strategies. <i>Nanomedicine</i> , 2018 , 13, 1639-1656	5.6	27
99	Tumor necrosis factor- α promotes hepatocellular carcinogenesis through the activation of hepatic progenitor cells. <i>Cancer Letters</i> , 2018 , 434, 22-32	9.9	32

98	LPS-induced CXCR4-dependent migratory properties and a mesenchymal-like phenotype of colorectal cancer cells. <i>Cell Adhesion and Migration</i> , 2017 , 11, 13-23	3.2	18
97	Lipopolysaccharide induces the differentiation of hepatic progenitor cells into myofibroblasts via activation of the Hedgehog signaling pathway. <i>Cell Cycle</i> , 2017 , 16, 1357-1365	4.7	8
96	Lipopolysaccharide promotes tumorigenicity of hepatic progenitor cells by promoting proliferation and blocking normal differentiation. <i>Cancer Letters</i> , 2017 , 386, 35-46	9.9	13
95	Autophagy-deficient Kupffer cells promote tumorigenesis by enhancing mtROS-NF-B-IL1 β -dependent inflammation and fibrosis during the preneoplastic stage of hepatocarcinogenesis. <i>Cancer Letters</i> , 2017 , 388, 198-207	9.9	52
94	Methylation mediated Gadd45 β enhanced the chemosensitivity of hepatocellular carcinoma by inhibiting the stemness of liver cancer cells. <i>Cell and Bioscience</i> , 2017 , 7, 63	9.8	6
93	Bcl-3 is a novel biomarker of renal fibrosis in chronic kidney disease. <i>Oncotarget</i> , 2017 , 8, 97206-97216	3.3	1
92	Pan-mTOR inhibitor MLN0128 is effective against intrahepatic cholangiocarcinoma in mice. <i>Journal of Hepatology</i> , 2017 , 67, 1194-1203	13.4	54
91	Peri-tumor associated fibroblasts promote intrahepatic metastasis of hepatocellular carcinoma by recruiting cancer stem cells. <i>Cancer Letters</i> , 2017 , 404, 19-28	9.9	27
90	Suppression of p53 potentiates chemosensitivity in nutrient-deprived cholangiocarcinoma cells via inhibition of autophagy. <i>Oncology Letters</i> , 2017 , 14, 1959-1966	2.6	11
89	Hepatocyte nuclear factor-1 β enhances the stemness of hepatocellular carcinoma cells through activation of the Notch pathway. <i>Scientific Reports</i> , 2017 , 7, 4793	4.9	14
88	Look into hepatic progenitor cell associated trait: Histological heterogeneity of hepatitis B-related combined hepatocellular-cholangiocarcinoma. <i>Current Medical Science</i> , 2017 , 37, 873-879	2.8	2
87	Involvement of proapoptotic genes in autophagic cell death induced by irradiation. <i>Cell Death Discovery</i> , 2017 , 3, 17068	6.9	4
86	The protective or damaging effect of Tumor necrosis factor- α in acute liver injury is concentration-dependent. <i>Cell and Bioscience</i> , 2016 , 6, 8	9.8	24
85	Autophagy regulates biliary differentiation of hepatic progenitor cells through Notch1 signaling pathway. <i>Cell Cycle</i> , 2016 , 15, 1602-10	4.7	15
84	Meta-analysis of laparoscopic versus open liver resection for colorectal liver metastases. <i>Oncotarget</i> , 2016 , 7, 84544-84555	3.3	18
83	Babao Dan attenuates hepatic fibrosis by inhibiting hepatic stellate cells activation and proliferation via TLR4 signaling pathway. <i>Oncotarget</i> , 2016 , 7, 82554-82566	3.3	15
82	Kupffer cells-dependent inflammation in the injured liver increases recruitment of mesenchymal stem cells in aging mice. <i>Oncotarget</i> , 2016 , 7, 1084-95	3.3	12
81	Inhibition of Growth and Metastasis of Colon Cancer by Delivering 5-Fluorouracil-loaded Pluronic P85 Copolymer Micelles. <i>Scientific Reports</i> , 2016 , 6, 20896	4.9	20

80	Lipopolysaccharide supports maintaining the stemness of CD133(+) hepatoma cells through activation of the NF- κ B/HIF-1 β pathway. <i>Cancer Letters</i> , 2016 , 378, 131-41	9.9	22
79	Hepatic stellate cell promoted hepatoma cell invasion via the HGF/c-Met signaling pathway regulated by p53. <i>Cell Cycle</i> , 2016 , 15, 886-94	4.7	25
78	Targeted and controlled drug delivery using a temperature and ultra-violet responsive liposome with excellent breast cancer suppressing ability. <i>RSC Advances</i> , 2015 , 5, 27630-27639	3.7	12
77	The role of autophagy induced by tumor microenvironment in different cells and stages of cancer. <i>Cell and Bioscience</i> , 2015 , 5, 14	9.8	95
76	TGF- β regulates hepatocellular carcinoma progression by inducing Treg cell polarization. <i>Cellular Physiology and Biochemistry</i> , 2015 , 35, 1623-32	3.9	70
75	Toll like receptor 4 facilitates invasion and migration as a cancer stem cell marker in hepatocellular carcinoma. <i>Cancer Letters</i> , 2015 , 358, 136-143	9.9	70
74	Contribution and Mobilization of Mesenchymal Stem Cells in a mouse model of carbon tetrachloride-induced liver fibrosis. <i>Scientific Reports</i> , 2015 , 5, 17762	4.9	28
73	Overexpression Of Hepatocyte Nuclear Factor-1beta Predicting Poor Prognosis Is Associated With Biliary Phenotype In Patients With Hepatocellular Carcinoma. <i>Scientific Reports</i> , 2015 , 5, 13319	4.9	13
72	A review on hepatocyte nuclear factor-1beta and tumor. <i>Cell and Bioscience</i> , 2015 , 5, 58	9.8	36
71	Controlled and Targeted Drug Delivery by a UV-responsive Liposome for Overcoming Chemo-resistance in Non-Hodgkin Lymphoma. <i>Chemical Biology and Drug Design</i> , 2015 , 86, 783-94	2.9	20
70	Effect of Autophagy on Chemotherapy-Induced Apoptosis and Growth Inhibition 2015 , 145-156		0
69	Corticosterone mediates the inhibitory effect of restraint stress on the migration of mesenchymal stem cell to carbon tetrachloride-induced fibrotic liver by downregulating CXCR4/7 expression. <i>Stem Cells and Development</i> , 2015 , 24, 587-96	4.4	11
68	The injured liver induces hyperimmunoglobulinemia by failing to dispose of antigens and endotoxins in the portal system. <i>PLoS ONE</i> , 2015 , 10, e0122739	3.7	10
67	Hepatitis B virus (HBV) receptors: Deficiency in tumor results in scant HBV infection and overexpression in peritumor leads to higher recurrence risk. <i>Oncotarget</i> , 2015 , 6, 42952-62	3.3	17
66	Cell-based therapy for acute and chronic liver failures: distinct diseases, different choices. <i>Scientific Reports</i> , 2014 , 4, 6494	4.9	29
65	Autophagy inhibits chemotherapy-induced apoptosis through downregulating Bad and Bim in hepatocellular carcinoma cells. <i>Scientific Reports</i> , 2014 , 4, 5382	4.9	31
64	Peritumoral ductular reaction: a poor postoperative prognostic factor for hepatocellular carcinoma. <i>BMC Cancer</i> , 2014 , 14, 65	4.8	15
63	Inhibition of tumor necrosis factor alpha reduces the outgrowth of hepatic micrometastasis of colorectal tumors in a mouse model of liver ischemia-reperfusion injury. <i>Journal of Biomedical Science</i> , 2014 , 21, 1	13.3	48

62	Proliferative ductular reactions correlate with hepatic progenitor cell and predict recurrence in HCC patients after curative resection. <i>Cell and Bioscience</i> , 2014 , 4, 50	9.8	26
61	Autophagy protects against palmitate-induced apoptosis in hepatocytes. <i>Cell and Bioscience</i> , 2014 , 4, 28	9.8	54
60	Inhibition of p53 increases chemosensitivity to 5-FU in nutrient-deprived hepatocarcinoma cells by suppressing autophagy. <i>Cancer Letters</i> , 2014 , 346, 278-84	9.9	32
59	Autophagy inhibition switches low-dose camptothecin-induced premature senescence to apoptosis in human colorectal cancer cells. <i>Biochemical Pharmacology</i> , 2014 , 90, 265-75	6	32
58	Expression of epithelial cell adhesion molecule associated with elevated ductular reactions in hepatocellular carcinoma. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2014 , 38, 699-705	2.4	14
57	Tumor-associated macrophages promote cancer stem cell-like properties via transforming growth factor-beta1-induced epithelial-mesenchymal transition in hepatocellular carcinoma. <i>Cancer Letters</i> , 2014 , 352, 160-8	9.9	267
56	High levels of SIRT1 expression enhance tumorigenesis and associate with a poor prognosis of colorectal carcinoma patients. <i>Scientific Reports</i> , 2014 , 4, 7481	4.9	92
55	Peptidylarginine deiminase IV promotes the development of chemoresistance through inducing autophagy in hepatocellular carcinoma. <i>Cell and Bioscience</i> , 2014 , 4, 49	9.8	18
54	Chronic restraint stress decreases the repair potential from mesenchymal stem cells on liver injury by inhibiting TGF- β 1 generation. <i>Cell Death and Disease</i> , 2014 , 5, e1308	9.8	11
53	Overexpression of SIRT1 promotes metastasis through epithelial-mesenchymal transition in hepatocellular carcinoma. <i>BMC Cancer</i> , 2014 , 14, 978	4.8	75
52	Activation of autophagy protects against cholestasis-induced hepatic injury. <i>Cell and Bioscience</i> , 2014 , 4, 47	9.8	25
51	Mesenchymal stem cells contribute to the chemoresistance of hepatocellular carcinoma cells in inflammatory environment by inducing autophagy. <i>Cell and Bioscience</i> , 2014 , 4, 22	9.8	22
50	Risk of treatment-related mortality with sorafenib in patients with cancer. <i>Asian Pacific Journal of Cancer Prevention</i> , 2014 , 14, 6681-6	1.7	9
49	Inhibition of autophagy enhances anticancer effects of bevacizumab in hepatocarcinoma. <i>Journal of Molecular Medicine</i> , 2013 , 91, 473-83	5.5	83
48	Autophagy lessens ischemic liver injury by reducing oxidative damage. <i>Cell and Bioscience</i> , 2013 , 3, 26	9.8	34
47	One cell, multiple roles: contribution of mesenchymal stem cells to tumor development in tumor microenvironment. <i>Cell and Bioscience</i> , 2013 , 3, 5	9.8	53
46	The silence of MUC2 mRNA induced by promoter hypermethylation associated with HBV in Hepatocellular Carcinoma. <i>BMC Medical Genetics</i> , 2013 , 14, 14	2.1	9
45	Autophagy contributes to the survival of CD133+ liver cancer stem cells in the hypoxic and nutrient-deprived tumor microenvironment. <i>Cancer Letters</i> , 2013 , 339, 70-81	9.9	109

44	Decreased PADI4 mRNA association with global hypomethylation in hepatocellular carcinoma during HBV exposure. <i>Cell Biochemistry and Biophysics</i> , 2013 , 65, 187-95	3.2	10
43	Rosiglitazone protects against palmitate-induced pancreatic beta-cell death by activation of autophagy via 5HAMP-activated protein kinase modulation. <i>Endocrine</i> , 2013 , 44, 87-98	4	52
42	Paradoxical roles of autophagy in different stages of tumorigenesis: protector for normal or cancer cells. <i>Cell and Bioscience</i> , 2013 , 3, 35	9.8	64
41	Tumor necrosis factor- α attenuates starvation-induced apoptosis through upregulation of ferritin heavy chain in hepatocellular carcinoma cells. <i>BMC Cancer</i> , 2013 , 13, 438	4.8	11
40	Pemetrexed plus platinum or gemcitabine plus platinum for advanced non-small cell lung cancer: final survival analysis from a multicentre randomized phase II trial in the East Asia region and a meta-analysis. <i>Respirology</i> , 2013 , 18, 131-139	3.6	12
39	Autophagy prevents irradiation injury and maintains stemness through decreasing ROS generation in mesenchymal stem cells. <i>Cell Death and Disease</i> , 2013 , 4, e844	9.8	124
38	Paradoxical role of autophagy in the dysplastic and tumor-forming stages of hepatocarcinoma development in rats. <i>Cell Death and Disease</i> , 2013 , 4, e501	9.8	54
37	Chloroquine promotes the anticancer effect of TACE in a rabbit VX2 liver tumor model. <i>International Journal of Biological Sciences</i> , 2013 , 9, 322-30	11.2	23
36	Hepatic stellate cells secreted hepatocyte growth factor contributes to the chemoresistance of hepatocellular carcinoma. <i>PLoS ONE</i> , 2013 , 8, e73312	3.7	48
35	The role of immunosuppression of mesenchymal stem cells in tissue repair and tumor growth. <i>Cell and Bioscience</i> , 2012 , 2, 8	9.8	68
34	Stem cells deployed for bone repair hijacked by T cells. <i>Cell Stem Cell</i> , 2012 , 10, 6-8	18	2
33	Targeting autophagy potentiates chemotherapy-induced apoptosis and proliferation inhibition in hepatocarcinoma cells. <i>Cancer Letters</i> , 2012 , 320, 171-9	9.9	141
32	CD133(+)CXCR4(+) colon cancer cells exhibit metastatic potential and predict poor prognosis of patients. <i>BMC Medicine</i> , 2012 , 10, 85	11.4	119
31	Toll-like receptor 4 signaling promotes epithelial-mesenchymal transition in human hepatocellular carcinoma induced by lipopolysaccharide. <i>BMC Medicine</i> , 2012 , 10, 98	11.4	91
30	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012 , 8, 445-544	14.2	2783
29	Combined EGFR and VEGFR versus single EGFR signaling pathways inhibition therapy for NSCLC: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2012 , 7, e40178	3.7	6
28	Mesenchymal stem cells in inflammation microenvironment accelerates hepatocellular carcinoma metastasis by inducing epithelial-mesenchymal transition. <i>PLoS ONE</i> , 2012 , 7, e43272	3.7	78
27	The silencing of RECK gene is associated with promoter hypermethylation and poor survival in hepatocellular carcinoma. <i>International Journal of Biological Sciences</i> , 2012 , 8, 451-8	11.2	32

26	Background progenitor activation is associated with recurrence after hepatectomy of combined hepatocellular-cholangiocarcinoma. <i>Hepatology</i> , 2012 , 56, 1804-16	11.2	59
25	Anti-tumor effect of 5-aza-2-deoxycytidine by inhibiting telomerase activity in hepatocellular carcinoma cells. <i>World Journal of Gastroenterology</i> , 2012 , 18, 2334-43	5.6	24
24	Association of alpha fetoprotein in hepatocellular carcinoma with activation of hepatic progenitor cells and patient prognosis. <i>Academic Journal of Second Military Medical University</i> , 2012 , 32, 136-139		
23	Maintenance therapy with continuous or switch strategy in advanced non-small cell lung cancer: a systematic review and meta-analysis. <i>Chest</i> , 2011 , 140, 117-126	5.3	61
22	Immunosuppressive effect of bone marrow-derived mesenchymal stem cells in inflammatory microenvironment favours the growth of B16 melanoma cells. <i>Journal of Cellular and Molecular Medicine</i> , 2011 , 15, 2343-52	5.6	56
21	Elevated expression of the stem cell marker CD133 associated with Line-1 demethylation in hepatocellular carcinoma. <i>Annals of Surgical Oncology</i> , 2011 , 18, 2373-80	3.1	28
20	Epithelial-Mesenchymal Transition in tumor microenvironment. <i>Cell and Bioscience</i> , 2011 , 1, 29	9.8	183
19	Autophagy in hypoxia protects cancer cells against apoptosis induced by nutrient deprivation through a Beclin1-dependent way in hepatocellular carcinoma. <i>Journal of Cellular Biochemistry</i> , 2011 , 112, 3406-20	4.7	39
18	CpG island methylator phenotype of cell-cycle regulators associated with TNM stage and poor prognosis in patients with oesophageal squamous cell carcinoma. <i>Journal of Clinical Pathology</i> , 2011 , 64, 246-51	3.9	22
17	Effects of inflammatory factors on mesenchymal stem cells and their role in the promotion of tumor angiogenesis in colon cancer. <i>Journal of Biological Chemistry</i> , 2011 , 286, 25007-15	5.4	137
16	Tumor necrosis factor-alpha promotes tumor growth by inducing vascular endothelial growth factor. <i>Cancer Investigation</i> , 2011 , 29, 485-93	2.1	16
15	Hepatoblast-like progenitor cells derived from embryonic stem cells can repopulate livers of mice. <i>Gastroenterology</i> , 2010 , 139, 2158-2169.e8	13.3	53
14	Autophagic cell death induced by 5-FU in Bax or PUMA deficient human colon cancer cell. <i>Cancer Letters</i> , 2010 , 288, 68-74	9.9	70
13	Correlation of CpG island methylator phenotype with poor prognosis in hepatocellular carcinoma. <i>Experimental and Molecular Pathology</i> , 2010 , 88, 112-7	4.4	26
12	Expression of differentiation inhibitory factor and prognosis of malignant tumors. <i>Academic Journal of Second Military Medical University</i> , 2010 , 30, 97-100		
11	Hypoxia-induced autophagy contributes to the chemoresistance of hepatocellular carcinoma cells. <i>Autophagy</i> , 2009 , 5, 1131-44	10.2	150
10	Up-regulation of hTERT expression by low-dose cisplatin contributes to chemotherapy resistance in human hepatocellular cancer cells. <i>Oncology Reports</i> , 2009 , 22, 549-56	3.5	15
9	siRNA-mediated inhibition of hTERT enhances chemosensitivity of hepatocellular carcinoma. <i>Cancer Biology and Therapy</i> , 2008 , 7, 1555-60	4.6	21

8	Methylation-related silencing of p14ARF gene correlates with telomerase activity and mRNA expression of human telomerase reverse transcriptase in hepatocellular carcinoma. <i>Journal of Surgical Oncology</i> , 2008 , 98, 462-8	2.8	13
7	CpG island methylator phenotype association with upregulated telomerase activity in hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2008 , 123, 998-1004	7.5	50
6	Lentiviral vector-mediated siRNA knockdown of SR-PSOX inhibits foam cell formation in vitro. <i>Acta Pharmacologica Sinica</i> , 2008 , 29, 847-52	8	23
5	Detection of telomerase activity in biopsy samples for predicting prognosis in cirrhotic patients with hepatocellular carcinoma after laparoscopic radiofrequency ablation therapy. <i>Chinese-German Journal of Clinical Oncology</i> , 2007 , 6, P210-P214		
4	CpG island methylator phenotype association with elevated serum alpha-fetoprotein level in hepatocellular carcinoma. <i>Clinical Cancer Research</i> , 2007 , 13, 944-52	12.9	56
3	Expression level of Bcl-XL critically affects sensitivity of hepatocellular carcinoma cells to LIGHT-enhanced and interferon- β -induced apoptosis. <i>Oncology Reports</i> , 2007 ,	3.5	1
2	Enhancement of immunogenicity of tumor cells by cotransfection with genes encoding antisense insulin-like growth factor-1 and B7.1 molecules. <i>Cancer Gene Therapy</i> , 2000 , 7, 456-65	5.4	11
1	Detection of human telomerase activity by telomerase TRAP ELISA assay. <i>Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research</i> , 1997 , 9, 277-280	3.8	2