

Chien-Feng Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6954293/publications.pdf>

Version: 2024-02-01

260
papers

7,973
citations

57681

46
h-index

104191

69
g-index

266
all docs

266
docs citations

266
times ranked

13186
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Disruption of the pentraxin 3/CD44 interaction as an efficient therapy for triple-negative breast cancers. <i>Clinical and Translational Medicine</i> , 2022, 12, e724. | 1.7 | 17 |
| 2 | Molecular Characterization of Dermatofibrosarcoma Protuberans. <i>American Journal of Surgical Pathology</i> , 2022, 46, 942-955. | 2.1 | 8 |
| 3 | Adult NTRK-rearranged spindle cell neoplasms of the viscera: with an emphasis on rare locations and heterologous elements. <i>Modern Pathology</i> , 2022, 35, 911-921. | 2.9 | 13 |
| 4 | Angiogenesis Driven by the CEBPD-hsa-miR-429-VEGFA Signaling Axis Promotes Urothelial Carcinoma Progression. <i>Cells</i> , 2022, 11, 638. | 1.8 | 9 |
| 5 | High Stromal SFRP2 Expression in Urothelial Carcinoma Confers an Unfavorable Prognosis. <i>Frontiers in Oncology</i> , 2022, 12, 834249. | 1.3 | 1 |
| 6 | Characterization of Aberrations in DNA Damage Repair Pathways in Gastrointestinal Stromal Tumors: The Clinicopathologic Relevance of H2AX and 53BP1 in Correlation with Heterozygous Deletions of CHEK2, BRCA2, and RB1. <i>Cancers</i> , 2022, 14, 1787. | 1.7 | 1 |
| 7 | Open the Technical Black Box of Tumor Mutational Burden (TMB): Factors Affecting Harmonization and Standardization of Panel-Based TMB. <i>International Journal of Molecular Sciences</i> , 2022, 23, 5097. | 1.8 | 3 |
| 8 | Discrimination of Methicillin-resistant <i>Staphylococcus aureus</i> by MALDI-TOF Mass Spectrometry with Machine Learning Techniques in Patients with <i>Staphylococcus aureus</i> Bacteremia. <i>Pathogens</i> , 2022, 11, 586. | 1.2 | 1 |
| 9 | High MT2A Expression Predicts Worse Prognosis in Patients with Urothelial Carcinoma. <i>Oncology</i> , 2022, 100, 485-497. | 0.9 | 1 |
| 10 | Lysine demethylase 2A expression in cancer-associated fibroblasts promotes breast tumour growth. <i>British Journal of Cancer</i> , 2021, 124, 484-493. | 2.9 | 17 |
| 11 | Metformin Resensitizes Sorafenib-Resistant HCC Cells Through AMPK-Dependent Autophagy Activation. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 8, . | 1.8 | 3 |
| 12 | Molecular biology of urothelial carcinoma. <i>Journal of Cancer Research and Practice</i> , 2021, 8, 1. | 0.2 | 1 |
| 13 | O6-methylguanine-DNA methyltransferase modulates cisplatin-induced DNA double-strand breaks by targeting the homologous recombination pathway in nasopharyngeal carcinoma. <i>Journal of Biomedical Science</i> , 2021, 28, 2. | 2.6 | 12 |
| 14 | RSF1 requires CEBP β and hSNF2H to promote IL-1 β -mediated angiogenesis: the clinical and therapeutic relevance of RSF1 overexpression and amplification in myxofibrosarcomas. <i>Angiogenesis</i> , 2021, 24, 533-548. | 3.7 | 4 |
| 15 | MIG is essential for promoting glucose metabolic reprogramming and tumor growth in triple-negative breast cancer. <i>EMBO Reports</i> , 2021, 22, e50781. | 2.0 | 8 |
| 16 | Identification of potential genes in upper tract urothelial carcinoma using next-generation sequencing with bioinformatics and in vitro analyses. <i>PeerJ</i> , 2021, 9, e11343. | 0.9 | 2 |
| 17 | High FRMD3 expression is prognostic for worse survival in rectal cancer patients treated with CCRT. <i>International Journal of Clinical Oncology</i> , 2021, 26, 1689-1697. | 1.0 | 9 |
| 18 | MIG Is Essential for Promoting Glucose Metabolic Reprogramming and Tumor Growth in Triple-Negative Breast Cancer. <i>FASEB Journal</i> , 2021, 35, . | 0.2 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Utility of EFEMP1 in the Prediction of Oncologic Outcomes of Urothelial Carcinoma. <i>Genes</i> , 2021, 12, 872. | 1.0 | 11 |
| 20 | Utility of Acute Physiology and Chronic Health Evaluation (APACHE II) in Predicting Mortality in Patients with Pyogenic Liver Abscess: A Retrospective Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 2644. | 1.0 | 3 |
| 21 | S100P as a marker for poor survival and advanced stage in gallbladder carcinoma. <i>Annals of Diagnostic Pathology</i> , 2021, 52, 151736. | 0.6 | 1 |
| 22 | High SPINK4 Expression Predicts Poor Outcomes among Rectal Cancer Patients Receiving CCRT. <i>Current Oncology</i> , 2021, 28, 2373-2384. | 0.9 | 7 |
| 23 | Upregulated MUC2 Is an Unfavorable Prognostic Indicator for Rectal Cancer Patients Undergoing Preoperative CCRT. <i>Journal of Clinical Medicine</i> , 2021, 10, 3030. | 1.0 | 6 |
| 24 | CTSE Overexpression Is an Adverse Prognostic Factor for Survival among Rectal Cancer Patients Receiving CCRT. <i>Life</i> , 2021, 11, 646. | 1.1 | 3 |
| 25 | Suppression of Extracellular Vesicle VEGF-Ca ²⁺ -mediated Lymphangiogenesis and Pancreatic Cancer Early Dissemination By a Selective HDAC1/2 Inhibitor. <i>Molecular Cancer Therapeutics</i> , 2021, 20, 1550-1560. | 1.9 | 14 |
| 26 | High G protein subunit beta 4 protein level is correlated to poor prognosis of urothelial carcinoma. <i>Medical Molecular Morphology</i> , 2021, 54, 356-367. | 0.4 | 2 |
| 27 | High glycosyltransferase ⁸ domain containing two protein levels contribute to poor prognosis in urothelial carcinoma. <i>International Journal of Urology</i> , 2021, 28, 1178-1187. | 0.5 | 0 |
| 28 | EMP2 induces cytotaxis and apoptosis via the TGF β ² /SMAD/SP1 axis and recruitment of P2RX7 in urinary bladder urothelial carcinoma. <i>Cellular Oncology (Dordrecht)</i> , 2021, 44, 1133-1150. | 2.1 | 3 |
| 29 | Prognostic Significance of ROR2 Expression in Patients with Urothelial Carcinoma. <i>Biomedicines</i> , 2021, 9, 1054. | 1.4 | 3 |
| 30 | Overexpression of Pyruvate Dehydrogenase Kinase-3 Predicts Poor Prognosis in Urothelial Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 749142. | 1.3 | 8 |
| 31 | Arginine starvation elicits chromatin leakage and cGAS-STING activation via epigenetic silencing of metabolic and DNA-repair genes. <i>Theranostics</i> , 2021, 11, 7527-7545. | 4.6 | 25 |
| 32 | Phase 1b study of pegylated arginine deiminase (ADI-PEG 20) plus Pembrolizumab in advanced solid cancers. <i>Oncolmmunology</i> , 2021, 10, 1943253. | 2.1 | 20 |
| 33 | Complete and Durable Response to Nivolumab in Recurrent Poorly Differentiated Pancreatic Neuroendocrine Carcinoma with High Tumor Mutational Burden. <i>Current Oncology</i> , 2021, 28, 4587-4596. | 0.9 | 4 |
| 34 | Myoepithelial and oral intracranial myxoid mesenchymal tumor-like neoplasms as diagnostic considerations of the ever-expanding extracranial myxocollagenous tumors harboring FET-CREB fusions. <i>Pathology Research and Practice</i> , 2021, 229, 153700. | 1.0 | 4 |
| 35 | High Level of Aristolochic Acid Detected With a Unique Genomic Landscape Predicts Early UTUC Onset After Renal Transplantation in Taiwan. <i>Frontiers in Oncology</i> , 2021, 11, 828314. | 1.3 | 4 |
| 36 | Biological significance of MYC and CEBPD coamplification in urothelial carcinoma: Multilayered genomic, transcriptional and posttranscriptional positive feedback loops enhance oncogenic glycolysis. <i>Clinical and Translational Medicine</i> , 2021, 11, e674. | 1.7 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Amplification-driven BCL6-suppressed cytostasis is mediated by transrepression of FOXO3 and post-translational modifications of FOXO3 in urinary bladder urothelial carcinoma. <i>Theranostics</i> , 2020, 10, 707-724. | 4.6 | 12 |
| 38 | Expression and prognostic utility of SSX2IP in patients with nasopharyngeal carcinoma. <i>Apmis</i> , 2020, 128, 287-297. | 0.9 | 6 |
| 39 | SLC14A1 prevents oncometabolite accumulation and recruits HDAC1 to transrepress oncometabolite genes in urothelial carcinoma. <i>Theranostics</i> , 2020, 10, 11775-11793. | 4.6 | 18 |
| 40 | Treatment of chronic hepatitis C regimens containing with recombinant interferon in patients with sustained virological response predicts risk of hepatocellular carcinoma. <i>Medicine (United States)</i> , 2020, 99, e22435. | 0.4 | 2 |
| 41 | Phosphorylation of PDHA by AMPK Drives TCA Cycle to Promote Cancer Metastasis. <i>Molecular Cell</i> , 2020, 80, 263-278.e7. | 4.5 | 120 |
| 42 | Prognostic Utility of FBLN2 Expression in Patients With Urothelial Carcinoma. <i>Frontiers in Oncology</i> , 2020, 10, 570340. | 1.3 | 7 |
| 43 | High Expression of UBE2B as a Poor Prognosis Factor in Patients With Rectal Cancer Following Chemoradiotherapy. <i>Anticancer Research</i> , 2020, 40, 6305-6317. | 0.5 | 13 |
| 44 | BMI1-KLF4 axis deficiency improves responses to neoadjuvant concurrent chemoradiotherapy in patients with rectal cancer. <i>Radiotherapy and Oncology</i> , 2020, 149, 249-258. | 0.3 | 12 |
| 45 | Role of Microtubule-Associated Protein 1b in Urothelial Carcinoma: Overexpression Predicts Poor Prognosis. <i>Cancers</i> , 2020, 12, 630. | 1.7 | 20 |
| 46 | DUSP2 regulates extracellular vesicle-VEGF secretion and pancreatic cancer early dissemination. <i>Journal of Extracellular Vesicles</i> , 2020, 9, 1746529. | 5.5 | 29 |
| 47 | Metformin Resensitizes Sorafenib-Resistant HCC Cells Through AMPK-Dependent Autophagy Activation. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 596655. | 1.8 | 19 |
| 48 | Abstract 1452: Hypoxia-induced MIG-6 promotes glucose metabolic reprogramming and tumorigenesis in triple-negative breast cancer. , 2020, , . | | 0 |
| 49 | KDM8/JMJD5 as a dual coactivator of AR and PKM2 integrates AR/EZH2 network and tumor metabolism in CRPC. <i>Oncogene</i> , 2019, 38, 17-32. | 2.6 | 77 |
| 50 | Oct4 upregulates osteopontin via Egr1 and is associated with poor outcome in human lung cancer. <i>BMC Cancer</i> , 2019, 19, 791. | 1.1 | 25 |
| 51 | Nuclear KIT induces a NFKBIB-RELA-KIT autoregulatory loop in imatinib-resistant gastrointestinal stromal tumors. <i>Oncogene</i> , 2019, 38, 6550-6565. | 2.6 | 14 |
| 52 | Identification of phenothiazine as an ETV1-targeting agent in gastrointestinal stromal tumors using the Connectivity Map. <i>International Journal of Oncology</i> , 2019, 55, 536-546. | 1.4 | 3 |
| 53 | Over-Expression of CHD4 Is an Independent Biomarker of Poor Prognosis in Patients with Rectal Cancers Receiving Concurrent Chemoradiotherapy. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4087. | 1.8 | 21 |
| 54 | Deduction of Novel Genes Potentially Involved in Upper Tract Urothelial Carcinoma Using Next-Generation Sequencing and Bioinformatics Approaches. <i>International Journal of Medical Sciences</i> , 2019, 16, 93-105. | 1.1 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Non-proteolytic ubiquitination of Hexokinase 2 by HectH9 controls tumor metabolism and cancer stem cell expansion. <i>Nature Communications</i> , 2019, 10, 2625. | 5.8 | 82 |
| 56 | Therapeutic benefit of second-look transurethral resection of bladder tumors for newly diagnosed T1 bladder cancer: a single-center experience. <i>International Urology and Nephrology</i> , 2019, 51, 1335-1342. | 0.6 | 8 |
| 57 | Expression of prothymosin $\hat{\pm}$ in lung cancer is associated with squamous cell carcinoma and smoking. <i>Oncology Letters</i> , 2019, 17, 5740-5746. | 0.8 | 2 |
| 58 | Hepatoma-derived growth factor supports the antiapoptosis and profibrosis of pancreatic stellate cells. <i>Cancer Letters</i> , 2019, 457, 180-190. | 3.2 | 18 |
| 59 | Myoepithelial carcinoma of tibia mimic giant cell tumor: a case report with emphasis on MR features. <i>Skeletal Radiology</i> , 2019, 48, 1637-1641. | 1.2 | 4 |
| 60 | YWHAZ amplification/overexpression defines aggressive bladder cancer and contributes to chemo- and radio-resistance by suppressing caspase-mediated apoptosis. <i>Journal of Pathology</i> , 2019, 248, 476-487. | 2.1 | 17 |
| 61 | Downregulation of the cytochrome P450 4B1 protein confers a poor prognostic factor in patients with urothelial carcinomas of upper urinary tracts and urinary bladder. <i>Apmis</i> , 2019, 127, 170-180. | 0.9 | 14 |
| 62 | High TNFAIP6 level is associated with poor prognosis of urothelial carcinomas. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019, 37, 293.e11-293.e24. | 0.8 | 21 |
| 63 | Inhibition of the formation of autophagosome but not autolysosome augments ABT-751-induced apoptosis in <i>TP53</i> -deficient Hep-3B cells. <i>Journal of Cellular Physiology</i> , 2019, 234, 9551-9563. | 2.0 | 5 |
| 64 | A phase II study of ADI-PEG 20 and FOLFOX6 in patients (pts) with advanced hepatocellular carcinoma (HCC).. <i>Journal of Clinical Oncology</i> , 2019, 37, TPS477-TPS477. | 0.8 | 2 |
| 65 | Decreased succinate dehydrogenase B in human hepatocellular carcinoma accelerates tumor malignancy by inducing the Warburg effect. <i>Scientific Reports</i> , 2018, 8, 3081. | 1.6 | 67 |
| 66 | Prediction of progression in skull base meningiomas: additional benefits of apparent diffusion coefficient value. <i>Journal of Neuro-Oncology</i> , 2018, 138, 63-71. | 1.4 | 17 |
| 67 | Draft genome sequence of CTX-M-type $\hat{2}$ -lactamase-producing <i>Klebsiella quasipneumoniae</i> subsp. <i>similipneumoniae</i> isolated from a Box turtle. <i>Journal of Global Antimicrobial Resistance</i> , 2018, 12, 235-236. | 0.9 | 8 |
| 68 | Higher nuclear EGFR expression is a better predictor of survival in rectal cancer patients following neoadjuvant chemoradiotherapy than cytoplasmic EGFR expression. <i>Oncology Letters</i> , 2018, 17, 1551-1558. | 0.8 | 10 |
| 69 | The critical role of AMPK in driving Akt activation under stress, tumorigenesis and drug resistance. <i>Nature Communications</i> , 2018, 9, 4728. | 5.8 | 125 |
| 70 | Hydroxysteroid 11-Beta Dehydrogenase 1 Overexpression with Copy-Number Gain and Missense Mutations in Primary Gastrointestinal Stromal Tumors. <i>Journal of Clinical Medicine</i> , 2018, 7, 408. | 1.0 | 9 |
| 71 | A phase 1 study of ADI-PEG 20 and modified FOLFOX6 in patients with advanced hepatocellular carcinoma and other gastrointestinal malignancies. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 429-440. | 1.1 | 35 |
| 72 | Autophagic reliance promotes metabolic reprogramming in oncogenic KRAS-driven tumorigenesis. <i>Autophagy</i> , 2018, 14, 1481-1498. | 4.3 | 37 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Proteomic Identification of the Galectin-1-Involved Molecular Pathways in Urinary Bladder Urothelial Carcinoma. <i>International Journal of Molecular Sciences</i> , 2018, 19, 1242. | 1.8 | 7 |
| 74 | Deciphering the Molecular Mechanism Underlying the Inhibitory Efficacy of Taiwanese Local Pomegranate Peels against Urinary Bladder Urothelial Carcinoma. <i>Nutrients</i> , 2018, 10, 543. | 1.7 | 11 |
| 75 | The diagnostic ureteroscopy before radical nephroureterectomy in upper urinary tract urothelial carcinoma is not associated with higher intravesical recurrence. <i>World Journal of Surgical Oncology</i> , 2018, 16, 135. | 0.8 | 26 |
| 76 | Periostin overexpression is associated with worse prognosis in nasopharyngeal carcinoma from endemic area: a cohort study. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 3205-3213. | 1.0 | 6 |
| 77 | Role of second-look transurethral resection of bladder tumors for newly diagnosed T1 bladder cancer: Experience at a single center. <i>Urological Science</i> , 2018, 29, 95. | 0.2 | 2 |
| 78 | ADI-PEG 20 and FOLFOX6: A phase 1 study in pts (pts) with advanced hepatocellular carcinoma (HCC).. <i>Journal of Clinical Oncology</i> , 2018, 36, 4085-4085. | 0.8 | 0 |
| 79 | Preprogramming therapeutic response of PI3K/mTOR dual inhibitor via the regulation of EHMT2 and p27 in pancreatic cancer. <i>American Journal of Cancer Research</i> , 2018, 8, 1812-1822. | 1.4 | 12 |
| 80 | Prognostic Value of Leptin Receptor Overexpression in Upper Tract Urothelial Carcinomas in Taiwan. <i>Clinical Genitourinary Cancer</i> , 2017, 15, e653-e659. | 0.9 | 11 |
| 81 | Overexpressed Fatty Acid Synthase in Gastrointestinal Stromal Tumors: Targeting a Progression-Associated Metabolic Driver Enhances the Antitumor Effect of Imatinib. <i>Clinical Cancer Research</i> , 2017, 23, 4908-4918. | 3.2 | 27 |
| 82 | Skp2 deficiency restricts the progression and stem cell features of castration-resistant prostate cancer by destabilizing Twist. <i>Oncogene</i> , 2017, 36, 4299-4310. | 2.6 | 56 |
| 83 | A hypoxia-responsive TRAF6-ATM-H2AX signalling axis promotes HIF1 α activation, tumorigenesis and metastasis. <i>Nature Cell Biology</i> , 2017, 19, 38-51. | 4.6 | 83 |
| 84 | Inhibition of the EGFR/STAT3/CEBPD Axis Reverses Cisplatin Cross-resistance with Paclitaxel in the Urothelial Carcinoma of the Urinary Bladder. <i>Clinical Cancer Research</i> , 2017, 23, 503-513. | 3.2 | 56 |
| 85 | Transmembrane and Coiled-Coil Domain 1 Impairs the AKT Signaling Pathway in Urinary Bladder Urothelial Carcinoma: A Characterization of a Tumor Suppressor. <i>Clinical Cancer Research</i> , 2017, 23, 7650-7663. | 3.2 | 22 |
| 86 | Anti-IL-20 monoclonal antibody inhibited tumor growth in hepatocellular carcinoma. <i>Scientific Reports</i> , 2017, 7, 17609. | 1.6 | 7 |
| 87 | ALK oncoproteins in atypical inflammatory myofibroblastic tumours: novel RRBP1-ALK fusions in epithelioid inflammatory myofibroblastic sarcoma. <i>Journal of Pathology</i> , 2017, 241, 316-323. | 2.1 | 87 |
| 88 | High Expression of Aldolase B Confers a Poor Prognosis for Rectal Cancer Patients Receiving Neoadjuvant Chemoradiotherapy. <i>Journal of Cancer</i> , 2017, 8, 1197-1204. | 1.2 | 13 |
| 89 | Overexpression of Transcobalamin 1 is an Independent Negative Prognosticator in Rectal Cancers Receiving Concurrent Chemoradiotherapy. <i>Journal of Cancer</i> , 2017, 8, 1330-1337. | 1.2 | 18 |
| 90 | Hypoxia-regulated MicroRNA-210 Overexpression is Associated with Tumor Development and Progression in Upper Tract Urothelial Carcinoma. <i>International Journal of Medical Sciences</i> , 2017, 14, 578-584. | 1.1 | 22 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | High Expression of EphA4 Predicted Lesser Degree of Tumor Regression after Neoadjuvant Chemoradiotherapy in Rectal Cancer. <i>Journal of Cancer</i> , 2017, 8, 1089-1096. | 1.2 | 14 |
| 92 | DPP4/CD26 overexpression in urothelial carcinoma confers an independent prognostic impact and correlates with intrinsic biological aggressiveness. <i>Oncotarget</i> , 2017, 8, 2995-3008. | 0.8 | 24 |
| 93 | Metformin promotes apoptosis in hepatocellular carcinoma through the CEBPD-induced autophagy pathway. <i>Oncotarget</i> , 2017, 8, 13832-13845. | 0.8 | 56 |
| 94 | PLCB4 copy gain and PLC γ 4 overexpression in primary gastrointestinal stromal tumors: Integrative characterization of a lipid-catabolizing enzyme associated with worse disease-free survival. <i>Oncotarget</i> , 2017, 8, 19997-20010. | 0.8 | 20 |
| 95 | Sulfatase-1 overexpression indicates poor prognosis in urothelial carcinoma of the urinary bladder and upper tract. <i>Oncotarget</i> , 2017, 8, 47216-47229. | 0.8 | 26 |
| 96 | Necdin Overexpression Predicts Poor Prognosis in Patients with Urothelial Carcinomas of the Upper Urinary Tract and Urinary Bladder. <i>Journal of Cancer</i> , 2016, 7, 304-313. | 1.2 | 11 |
| 97 | Complement Component 1, s Subcomponent Overexpression is an Independent Poor Prognostic Indicator in Patients with Urothelial Carcinomas of the Upper Urinary Tract and Urinary Bladder. <i>Journal of Cancer</i> , 2016, 7, 1396-1405. | 1.2 | 13 |
| 98 | Glutamate Decarboxylase 1 Overexpression as a Poor Prognostic Factor in Patients with Nasopharyngeal Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 1716-1723. | 1.2 | 16 |
| 99 | Prognostic Impact of Thrombospondin-2 (THBS2) Overexpression on Patients with Urothelial Carcinomas of Upper Urinary Tracts and Bladders. <i>Journal of Cancer</i> , 2016, 7, 1541-1549. | 1.2 | 44 |
| 100 | Overexpression of the PSAT1 Gene in Nasopharyngeal Carcinoma Is an Indicator of Poor Prognosis. <i>Journal of Cancer</i> , 2016, 7, 1088-1094. | 1.2 | 29 |
| 101 | Malignant Peripheral Nerve Sheath Tumor of Prostate: A Rare Case Report and Literature Review. <i>Case Reports in Urology</i> , 2016, 2016, 1-3. | 0.1 | 6 |
| 102 | Downregulation of RNF128 Predicts Progression and Poor Prognosis in Patients with Urothelial Carcinoma of the Upper Tract and Urinary Bladder. <i>Journal of Cancer</i> , 2016, 7, 2187-2196. | 1.2 | 17 |
| 103 | SPOCK1 Overexpression Confers a Poor Prognosis in Urothelial Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 467-476. | 1.2 | 37 |
| 104 | CSF2 Overexpression Is Associated with STAT5 Phosphorylation and Poor Prognosis in Patients with Urothelial Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 711-721. | 1.2 | 34 |
| 105 | Inhibition of HDAC3- and HDAC6-Promoted Survivin Expression Plays an Important Role in SAHA-Induced Autophagy and Viability Reduction in Breast Cancer Cells. <i>Frontiers in Pharmacology</i> , 2016, 7, 81. | 1.6 | 53 |
| 106 | Overexpression of Transient Receptor Protein Cation Channel Subfamily A Member 1, Confers an Independent Prognostic Indicator in Nasopharyngeal Carcinoma. <i>Journal of Cancer</i> , 2016, 7, 1181-1188. | 1.2 | 16 |
| 107 | High Expression of 17 β -hydroxysteroid Dehydrogenase Type 2 is Associated with a Better Prognosis in Urothelial Carcinoma of the Urinary Tract. <i>Journal of Cancer</i> , 2016, 7, 2221-2230. | 1.2 | 4 |
| 108 | Clinicopathological and genetic heterogeneity of the head and neck solitary fibrous tumours: a comparative histological, immunohistochemical and molecular study of 36 cases. <i>Histopathology</i> , 2016, 68, 492-501. | 1.6 | 49 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | <i>NAB2-STAT6</i> gene fusion and STAT6 immunoeexpression in extrathoracic solitary fibrous tumors: the association between fusion variants and locations. <i>Pathology International</i> , 2016, 66, 288-296. | 0.6 | 42 |
| 110 | Matrix metalloproteinase-11 as a marker of metastasis and predictor of poor survival in urothelial carcinomas. <i>Journal of Surgical Oncology</i> , 2016, 113, 700-707. | 0.8 | 17 |
| 111 | Metabolic Stress-Induced Phosphorylation of KAP1 Ser473 Blocks Mitochondrial Fusion in Breast Cancer Cells. <i>Cancer Research</i> , 2016, 76, 5006-5018. | 0.4 | 50 |
| 112 | Prognostic significance of DSG3 in rectal adenocarcinoma treated with preoperative chemoradiotherapy. <i>Future Oncology</i> , 2016, 12, 1457-1467. | 1.1 | 7 |
| 113 | Overexpression of JAK2: a predictor of unfavorable prognosis for nasopharyngeal carcinoma. <i>Future Oncology</i> , 2016, 12, 1887-1896. | 1.1 | 12 |
| 114 | Is preoperative anemia a risk factor for upper tract urothelial carcinoma following radical nephroureterectomy?. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016, 34, 337.e1-337.e9. | 0.8 | 10 |
| 115 | Role of galectin-1 in urinary bladder urothelial carcinoma cell invasion through the JNK pathway. <i>Cancer Science</i> , 2016, 107, 1390-1398. | 1.7 | 21 |
| 116 | A microtubule inhibitor, ABT-751, induces autophagy and delays apoptosis in Huh-7 cells. <i>Toxicology and Applied Pharmacology</i> , 2016, 311, 88-98. | 1.3 | 19 |
| 117 | The clinicopathological significance of <i>NAB2-STAT6</i> gene fusions in 52 cases of intrathoracic solitary fibrous tumors. <i>Cancer Medicine</i> , 2016, 5, 159-168. | 1.3 | 61 |
| 118 | The DNA Damage Transducer RNF8 Facilitates Cancer Chemoresistance and Progression through Twist Activation. <i>Molecular Cell</i> , 2016, 63, 1021-1033. | 4.5 | 82 |
| 119 | Characterization of FN1-FGFR1 and novel FN1-FGFR1 fusion genes in a large series of phosphaturic mesenchymal tumors. <i>Modern Pathology</i> , 2016, 29, 1335-1346. | 2.9 | 139 |
| 120 | TRAF6 Restricts p53 Mitochondrial Translocation, Apoptosis, and Tumor Suppression. <i>Molecular Cell</i> , 2016, 64, 803-814. | 4.5 | 63 |
| 121 | Inflammatory myofibroblastic tumor with RANBP2 and ALK gene rearrangement with bland cytological features mimicking desmoid-type fibromatosis: A case report and review of the literature. <i>Oncology Letters</i> , 2016, 11, 1429-1434. | 0.8 | 10 |
| 122 | BCAT1 overexpression is an indicator of poor prognosis in patients with urothelial carcinomas of the upper urinary tract and urinary bladder. <i>Histopathology</i> , 2016, 68, 520-532. | 1.6 | 29 |
| 123 | Overexpression of PTP4A3 is associated with metastasis and unfavorable prognosis in bladder cancer. <i>World Journal of Urology</i> , 2016, 34, 835-846. | 1.2 | 6 |
| 124 | The Prognostic Significance of Inflammation-Associated Blood Cell Markers in Patients with Upper Tract Urothelial Carcinoma. <i>Annals of Surgical Oncology</i> , 2016, 23, 343-351. | 0.7 | 43 |
| 125 | IL-18-induced interaction between IMP3 and HuR contributes to <i>COX-2</i> mRNA stabilization in acute myeloid leukemia. <i>Journal of Leukocyte Biology</i> , 2016, 99, 131-141. | 1.5 | 21 |
| 126 | An Effective and Well Tolerated Strategy of Bladder Preservation Therapy in Cisplatin-Ineligible Patients With Muscle-Invasive Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2016, 14, e67-e74. | 0.9 | 2 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Laparoscopic partial nephrectomy without intracorporeal suturing. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2016, 30, 1585-1591. | 1.3 | 9 |
| 128 | Differentiation between Glioblastoma Multiforme and Primary Cerebral Lymphoma: Additional Benefits of Quantitative Diffusion-Weighted MR Imaging. <i>PLoS ONE</i> , 2016, 11, e0162565. | 1.1 | 29 |
| 129 | Transcriptomic reappraisal identifies <i>MGLL</i> overexpression as an unfavorable prognosticator in primary gastrointestinal stromal tumors. <i>Oncotarget</i> , 2016, 7, 49986-49997. | 0.8 | 12 |
| 130 | MCM10 overexpression implicates adverse prognosis in urothelial carcinoma. <i>Oncotarget</i> , 2016, 7, 77777-77792. | 0.8 | 34 |
| 131 | DDR2 overexpression in urothelial carcinoma indicates an unfavorable prognosis: a large cohort study. <i>Oncotarget</i> , 2016, 7, 78918-78931. | 0.8 | 11 |
| 132 | Lysine demethylase 2A promotes stemness and angiogenesis of breast cancer by upregulating Jagged1. <i>Oncotarget</i> , 2016, 7, 27689-27710. | 0.8 | 51 |
| 133 | Overexpression of regulator of G protein signaling 11 promotes cell migration and associates with advanced stages and aggressiveness of lung adenocarcinoma. <i>Oncotarget</i> , 2016, 7, 31122-31136. | 0.8 | 15 |
| 134 | Advances in molecular genetics of early-stage urothelial carcinoma. <i>Translational Cancer Research</i> , 2016, 5, S1126-S1130. | 0.4 | 1 |
| 135 | Ductal activation of oncogenic KRAS alone induces sarcomatoid phenotype. <i>Scientific Reports</i> , 2015, 5, 13347. | 1.6 | 13 |
| 136 | Aspiration cytology of an ectopic cervical thymoma misinterpreted as a lymphoproliferative lesion of the thyroid: A case report. <i>Oncology Letters</i> , 2015, 10, 1255-1258. | 0.8 | 10 |
| 137 | Anesthetic Propofol Overdose Causes Vascular Hyperpermeability by Reducing Endothelial Glycocalyx and ATP Production. <i>International Journal of Molecular Sciences</i> , 2015, 16, 12092-12107. | 1.8 | 26 |
| 138 | Targeting chemotherapy-induced PTX3 in tumor stroma to prevent the progression of drug-resistant cancers. <i>Oncotarget</i> , 2015, 6, 23987-24001. | 0.8 | 51 |
| 139 | HAS3 underexpression as an indicator of poor prognosis in patients with urothelial carcinoma of the upper urinary tract and urinary bladder. <i>Tumor Biology</i> , 2015, 36, 5441-5450. | 0.8 | 26 |
| 140 | Characterization and Prognostic Significance of Methylthioadenosine Phosphorylase Deficiency in Nasopharyngeal Carcinoma. <i>Medicine (United States)</i> , 2015, 94, e2271. | 0.4 | 6 |
| 141 | Chromatophagy: Autophagy goes nuclear and captures broken chromatin during arginine-starvation. <i>Autophagy</i> , 2015, 11, 419-421. | 4.3 | 20 |
| 142 | Galectin-1 Dysregulation Independently Predicts Disease Specific Survival in Bladder Urothelial Carcinoma. <i>Journal of Urology</i> , 2015, 193, 1002-1008. | 0.2 | 23 |
| 143 | Skp2-Dependent Ubiquitination and Activation of LKB1 Is Essential for Cancer Cell Survival under Energy Stress. <i>Molecular Cell</i> , 2015, 57, 1022-1033. | 4.5 | 97 |
| 144 | NAB2-STAT6 fusion types account for clinicopathological variations in solitary fibrous tumors. <i>Modern Pathology</i> , 2015, 28, 1324-1335. | 2.9 | 133 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Overexpression of DNAJC12 predicts poor response to neoadjuvant concurrent chemoradiotherapy in patients with rectal cancer. <i>Experimental and Molecular Pathology</i> , 2015, 98, 338-345. | 0.9 | 25 |
| 146 | GPX2 underexpression indicates poor prognosis in patients with urothelial carcinomas of the upper urinary tract and urinary bladder. <i>World Journal of Urology</i> , 2015, 33, 1777-1789. | 1.2 | 27 |
| 147 | FGF7 Over Expression is an Independent Prognosticator in Patients with Urothelial Carcinoma of the Upper Urinary Tract and Bladder. <i>Journal of Urology</i> , 2015, 194, 223-229. | 0.2 | 37 |
| 148 | The prognostic impact of lipid biosynthesis-associated markers, HSD17B2 and HMGCS2, in rectal cancer treated with neoadjuvant concurrent chemoradiotherapy. <i>Tumor Biology</i> , 2015, 36, 7675-7683. | 0.8 | 31 |
| 149 | Skp2â€‘MacroH2A1â€‘CDK8 axis orchestrates G2/M transition and tumorigenesis. <i>Nature Communications</i> , 2015, 6, 6641. | 5.8 | 87 |
| 150 | Positive effects of cell-free porous PLGA implants and early loading exercise on hyaline cartilage regeneration in rabbits. <i>Acta Biomaterialia</i> , 2015, 28, 128-137. | 4.1 | 37 |
| 151 | <scp>PLA</scp>2G2A overexpression is associated with poor therapeutic response and inferior outcome in rectal cancer patients receiving neoadjuvant concurrent chemoradiotherapy. <i>Histopathology</i> , 2015, 66, 991-1002. | 1.6 | 15 |
| 152 | Subcellular localisation of anillin is associated with different survival outcomes in upper urinary tract urothelial carcinoma. <i>Journal of Clinical Pathology</i> , 2015, 68, 1026-1032. | 1.0 | 14 |
| 153 | PTP4A3 Independently Predicts Metastasis and Survival in Upper Tract Urothelial Carcinoma Treated with Radical Nephroureterectomy. <i>Journal of Urology</i> , 2015, 194, 1449-1455. | 0.2 | 10 |
| 154 | HMDB and 5-AzadC Combination Reverses Tumor Suppressor CCAAT/Enhancer-Binding Protein Delta to Strengthen the Death of Liver Cancer Cells. <i>Molecular Cancer Therapeutics</i> , 2015, 14, 2623-2633. | 1.9 | 19 |
| 155 | Saikosaponin b2 is a naturally occurring terpenoid that efficiently inhibits hepatitis C virus entry. <i>Journal of Hepatology</i> , 2015, 62, 541-548. | 1.8 | 99 |
| 156 | INHBA overexpression indicates poor prognosis in urothelial carcinoma of urinary bladder and upper tract. <i>Journal of Surgical Oncology</i> , 2015, 111, 414-422. | 0.8 | 39 |
| 157 | The cAMP responsive element binding protein 1 transactivates epithelial membrane protein 2, a potential tumor suppressor in the urinary bladder urothelial carcinoma. <i>Oncotarget</i> , 2015, 6, 9220-9239. | 0.8 | 30 |
| 158 | <i>CEBPD</i> amplification and overexpression in urothelial carcinoma: a driver of tumor metastasis indicating adverse prognosis. <i>Oncotarget</i> , 2015, 6, 31069-31084. | 0.8 | 35 |
| 159 | CDCA5 overexpression is an indicator of poor prognosis in patients with urothelial carcinomas of the upper urinary tract and urinary bladder. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 710-22. | 0.0 | 54 |
| 160 | The prognostic impact of RAP2A expression in patients with early and locoregionally advanced nasopharyngeal carcinoma in an endemic area. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 912-21. | 0.0 | 12 |
| 161 | Low thrombospondin 2 expression is predictive of low tumor regression after neoadjuvant chemoradiotherapy in rectal cancer. <i>American Journal of Translational Research (discontinued)</i> , 2015, 7, 2423-32. | 0.0 | 10 |
| 162 | Overexpression of REG4 confers an independent negative prognosticator in rectal cancers receiving concurrent chemoradiotherapy. <i>Journal of Surgical Oncology</i> , 2014, 110, 1002-1010. | 0.8 | 13 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 163 | Overexpression of CPS1 is an independent negative prognosticator in rectal cancers receiving concurrent chemoradiotherapy. <i>Tumor Biology</i> , 2014, 35, 11097-11105. | 0.8 | 25 |
| 164 | Angiomatoid fibrous histiocytoma: clinicopathological and molecular characterisation with emphasis on variant histomorphology. <i>Journal of Clinical Pathology</i> , 2014, 67, 210-215. | 1.0 | 53 |
| 165 | Galectin-1 overexpression in nasopharyngeal carcinoma: effect on survival. <i>Acta Oto-Laryngologica</i> , 2014, 134, 536-542. | 0.3 | 7 |
| 166 | AMACR Amplification in Myxofibrosarcomas: A Mechanism of Overexpression That Promotes Cell Proliferation with Therapeutic Relevance. <i>Clinical Cancer Research</i> , 2014, 20, 6141-6152. | 3.2 | 32 |
| 167 | Overexpression of CDC28 protein kinase regulatory subunit 1B confers an independent prognostic factor in nasopharyngeal carcinoma. <i>Apmis</i> , 2014, 122, 206-214. | 0.9 | 10 |
| 168 | Fibroblast growth factor receptor 2 overexpression is predictive of poor prognosis in rectal cancer patients receiving neoadjuvant chemoradiotherapy. <i>Journal of Clinical Pathology</i> , 2014, 67, 1056-1061. | 1.0 | 23 |
| 169 | Dedifferentiated liposarcoma can induce a leukemoid reaction. <i>Formosan Journal of Surgery</i> , 2014, 47, 42-48. | 0.1 | 0 |
| 170 | TOP2A overexpression as a poor prognostic factor in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 179-187. | 0.8 | 72 |
| 171 | Deficiency in expression and epigenetic DNA Methylation of ASS1 gene in nasopharyngeal carcinoma: negative prognostic impact and therapeutic relevance. <i>Tumor Biology</i> , 2014, 35, 161-169. | 0.8 | 54 |
| 172 | Overexpression of stathmin 1 confers an independent prognostic indicator in nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 2619-2629. | 0.8 | 40 |
| 173 | Fibronectin overexpression is associated with latent membrane protein 1 expression and has independent prognostic value for nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 1703-1712. | 0.8 | 40 |
| 174 | BCL6 overexpression is associated with decreased p19ARF expression and confers an independent prognosticator in gallbladder carcinoma. <i>Tumor Biology</i> , 2014, 35, 1417-1426. | 0.8 | 9 |
| 175 | Prognostic and Therapeutic Impact of Argininosuccinate Synthetase 1 Control in Bladder Cancer as Monitored Longitudinally by PET Imaging. <i>Cancer Research</i> , 2014, 74, 896-907. | 0.4 | 122 |
| 176 | Cancer/stroma interplay via cyclooxygenase-2 and indoleamine 2,3-dioxygenase promotes breast cancer progression. <i>Breast Cancer Research</i> , 2014, 16, 410. | 2.2 | 119 |
| 177 | Clinical Aggressiveness of Myxofibrosarcomas Associates with Down-Regulation of p12CDK2AP1: Prognostic Implication of a Putative Tumor Suppressor that Induces Cell Cycle Arrest and Apoptosis Via Mitochondrial Pathway. <i>Annals of Surgical Oncology</i> , 2014, 21, 711-720. | 0.7 | 2 |
| 178 | Cyclin-dependent kinase 4 overexpression is mostly independent of gene amplification and constitutes an independent prognosticator for nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 7209-7216. | 0.8 | 9 |
| 179 | Deficiency in asparagine synthetase expression in rectal cancers receiving concurrent chemoradiotherapy: negative prognostic impact and therapeutic relevance. <i>Tumor Biology</i> , 2014, 35, 6823-6830. | 0.8 | 18 |
| 180 | Overexpression of ANXA1 confers independent negative prognostic impact in rectal cancers receiving concurrent chemoradiotherapy. <i>Tumor Biology</i> , 2014, 35, 7755-7763. | 0.8 | 17 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 181 | AMACR overexpression as a poor prognostic factor in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 7983-7991. | 0.8 | 7 |
| 182 | Downregulated <i>MTAP</i> expression in myxofibrosarcoma: A characterization of inactivating mechanisms, tumor suppressive function, and therapeutic relevance. <i>Oncotarget</i> , 2014, 5, 11428-11441. | 0.8 | 21 |
| 183 | <i>AMACR</i> amplification and overexpression in primary imatinib-naïve gastrointestinal stromal tumors: a driver of cell proliferation indicating adverse prognosis. <i>Oncotarget</i> , 2014, 5, 11588-11603. | 0.8 | 22 |
| 184 | MTOR inhibition enhances NVP-AUY922-induced autophagy-mediated KIT degradation and cytotoxicity in imatinib-resistant gastrointestinal stromal tumors. <i>Oncotarget</i> , 2014, 5, 11723-11736. | 0.8 | 21 |
| 185 | The protective effects of glutamine in a rat model of ventilator-induced lung injury. <i>Journal of Thoracic Disease</i> , 2014, 6, 1704-13. | 0.6 | 3 |
| 186 | Nicotinamide N-methyltransferase overexpression is associated with Akt phosphorylation and indicates worse prognosis in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2013, 34, 3923-3931. | 0.8 | 35 |
| 187 | Clinicopathological significance of HuR expression in gallbladder carcinoma: with special emphasis on the implications of its nuclear and cytoplasmic expression. <i>Tumor Biology</i> , 2013, 34, 3059-3069. | 0.8 | 18 |
| 188 | Fatty acid synthase overexpression confers an independent prognosticator and associates with radiation resistance in nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2013, 34, 759-768. | 0.8 | 34 |
| 189 | SKP2 overexpression is associated with a poor prognosis of rectal cancer treated with chemoradiotherapy and represents a therapeutic target with high potential. <i>Tumor Biology</i> , 2013, 34, 1107-1117. | 0.8 | 34 |
| 190 | Pharmacological Inactivation of Skp2 SCF Ubiquitin Ligase Restricts Cancer Stem Cell Traits and Cancer Progression. <i>Cell</i> , 2013, 154, 556-568. | 13.5 | 335 |
| 191 | A rare malignant tumor of scalp in a 3-month-old Taiwanese infancy: case report of primitive myxoid mesenchymal tumor of infancy with molecular study. <i>Medical Molecular Morphology</i> , 2013, 46, 109-113. | 0.4 | 11 |
| 192 | E2F transcription factor 1 overexpression as a poor prognostic factor in patients with nasopharyngeal carcinomas. <i>Biomarkers and Genomic Medicine</i> , 2013, 5, 23-30. | 0.2 | 0 |
| 193 | A biosensor of high-density lipoprotein of human serum on a liquid crystal and polymer composite film. <i>Proceedings of SPIE</i> , 2013, , . | 0.8 | 4 |
| 194 | <i>ASS1</i> as a Novel Tumor Suppressor Gene in Myxofibrosarcomas: Aberrant Loss via Epigenetic DNA Methylation Confers Aggressive Phenotypes, Negative Prognostic Impact, and Therapeutic Relevance. <i>Clinical Cancer Research</i> , 2013, 19, 2861-2872. | 3.2 | 118 |
| 195 | Transplantation of autologous endothelial progenitor cells in porous PLGA scaffolds create a microenvironment for the regeneration of hyaline cartilage in rabbits. <i>Osteoarthritis and Cartilage</i> , 2013, 21, 1613-1622. | 0.6 | 47 |
| 196 | Cycles of Ubiquitination and Deubiquitination Critically Regulate Growth Factor-Mediated Activation of Akt Signaling. <i>Science Signaling</i> , 2013, 6, ra3. | 1.6 | 59 |
| 197 | The Effect of Osteochondral Regeneration Using Polymer Constructs and Continuous Passive Motion Therapy in the Lower Weight-Bearing Zone of Femoral Trochlear Groove in Rabbits. <i>Annals of Biomedical Engineering</i> , 2013, 41, 385-397. | 1.3 | 8 |
| 198 | Loss of epithelial membrane protein-2 expression confers an independent prognosticator in gallbladder carcinoma. <i>Biomarkers and Genomic Medicine</i> , 2013, 5, 31-38. | 0.2 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 199 | Overexpression of thymidylate synthetase confers an independent prognostic indicator in nasopharyngeal carcinoma. <i>Experimental and Molecular Pathology</i> , 2013, 95, 83-90. | 0.9 | 34 |
| 200 | Ectopic hamartomatous thymoma is distinct from lipomatous pleomorphic adenoma in lacking <i>PLAG1</i> aberration. <i>Histopathology</i> , 2013, 62, 518-522. | 1.6 | 11 |
| 201 | The E2F Transcription Factor 1 Transactivates Stathmin 1 in Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2013, 20, 4041-4054. | 0.7 | 53 |
| 202 | CCAAT/Enhancer Binding Protein β in Macrophages Contributes to Immunosuppression and Inhibits Phagocytosis in Nasopharyngeal Carcinoma. <i>Science Signaling</i> , 2013, 6, ra59. | 1.6 | 42 |
| 203 | IGFBP-5 overexpression as a poor prognostic factor in patients with urothelial carcinomas of upper urinary tracts and urinary bladder. <i>Journal of Clinical Pathology</i> , 2013, 66, 573-582. | 1.0 | 43 |
| 204 | HuR cytoplasmic expression is associated with increased cyclin A expression and inferior disease-free survival in patients with gastrointestinal stromal tumours (<i>GIST</i> s). <i>Histopathology</i> , 2013, 63, 445-454. | 1.6 | 7 |
| 205 | Autophagy is involved in endogenous and NVP-AUY922-induced KIT degradation in gastrointestinal stromal tumors. <i>Autophagy</i> , 2013, 9, 220-233. | 4.3 | 27 |
| 206 | Selecting Tyrosine Kinase Inhibitors for Gastrointestinal Stromal Tumor with Secondary KIT Activation-Loop Domain Mutations. <i>PLoS ONE</i> , 2013, 8, e65762. | 1.1 | 20 |
| 207 | Inhibiting Interleukin-19 Activity Ameliorates Esophageal Squamous Cell Carcinoma Progression. <i>PLoS ONE</i> , 2013, 8, e75254. | 1.1 | 15 |
| 208 | Interleukin-19 in Breast Cancer. <i>Clinical and Developmental Immunology</i> , 2013, 2013, 1-9. | 3.3 | 34 |
| 209 | Loss of ZBRK1 Contributes to the Increase of KAP1 and Promotes KAP1-Mediated Metastasis and Invasion in Cervical Cancer. <i>PLoS ONE</i> , 2013, 8, e73033. | 1.1 | 35 |
| 210 | β -Adrenoceptor agonist dexmedetomidine protects septic acute kidney injury through increasing BMP-7 and inhibiting HDAC2 and HDAC5. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, F1443-F1453. | 1.3 | 76 |
| 211 | Rsf-1 expression in rectal cancer: with special emphasis on the independent prognostic value after neoadjuvant chemoradiation. <i>Journal of Clinical Pathology</i> , 2012, 65, 687-692. | 1.0 | 24 |
| 212 | Vav3-Rac1 Signaling Regulates Prostate Cancer Metastasis with Elevated Vav3 Expression Correlating with Prostate Cancer Progression and Posttreatment Recurrence. <i>Cancer Research</i> , 2012, 72, 3000-3009. | 0.4 | 56 |
| 213 | Alpha-methylacyl coenzyme A racemase overexpression in gallbladder carcinoma confers an independent prognostic indicator. <i>Journal of Clinical Pathology</i> , 2012, 65, 309-314. | 1.0 | 15 |
| 214 | Expression of cyclin-dependent kinase 2-associated protein 1 confers an independent prognosticator in nasopharyngeal carcinoma: a cohort study. <i>Journal of Clinical Pathology</i> , 2012, 65, 795-801. | 1.0 | 8 |
| 215 | Characterization of Gene Amplification-Driven SKP2 Overexpression in Myxofibrosarcoma: Potential Implications in Tumor Progression and Therapeutics. <i>Clinical Cancer Research</i> , 2012, 18, 1598-1610. | 3.2 | 40 |
| 216 | Upregulated IL-19 in Breast Cancer Promotes Tumor Progression and Affects Clinical Outcome. <i>Clinical Cancer Research</i> , 2012, 18, 713-725. | 3.2 | 54 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 217 | Associations of Rsf-1 overexpression with poor therapeutic response and worse survival in patients with nasopharyngeal carcinoma. <i>Journal of Clinical Pathology</i> , 2012, 65, 248-253. | 1.0 | 32 |
| 218 | Anti-IL-20 Monoclonal Antibody Suppresses Breast Cancer Progression and Bone Osteolysis in Murine Models. <i>Journal of Immunology</i> , 2012, 188, 1981-1991. | 0.4 | 60 |
| 219 | Loss of epithelial membrane protein-2 expression confers an independent prognosticator in nasopharyngeal carcinoma: a cohort study. <i>BMJ Open</i> , 2012, 2, e000900. | 0.8 | 11 |
| 220 | Enhancer of Zeste Homolog 2 Overexpression in Nasopharyngeal Carcinoma: An Independent Poor Prognosticator That Enhances Cell Growth. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012, 82, 597-604. | 0.4 | 27 |
| 221 | Rsf-1/HBXAP overexpression is independent of gene amplification and is associated with poor outcome in patients with urinary bladder urothelial carcinoma. <i>Journal of Clinical Pathology</i> , 2012, 65, 802-807. | 1.0 | 15 |
| 222 | Recurrent Amplification at 7q21.2 Targets CDK6 Gene in Primary Myxofibrosarcomas and Identifies CDK6 Overexpression as an Independent Adverse Prognosticator. <i>Annals of Surgical Oncology</i> , 2012, 19, 2716-2725. | 0.7 | 23 |
| 223 | The Skp2-SCF E3 Ligase Regulates Akt Ubiquitination, Glycolysis, Herceptin Sensitivity, and Tumorigenesis. <i>Cell</i> , 2012, 149, 1098-1111. | 13.5 | 332 |
| 224 | HuR cytoplasmic expression is associated with increased cyclin A expression and poor outcome with upper urinary tract urothelial carcinoma. <i>BMC Cancer</i> , 2012, 12, 611. | 1.1 | 37 |
| 225 | EGFR Nuclear Import in Gallbladder Carcinoma: Nuclear Phosphorylated EGFR Upregulates iNOS Expression and Confers Independent Prognostic Impact. <i>Annals of Surgical Oncology</i> , 2012, 19, 443-454. | 0.7 | 28 |
| 226 | The combined effects of continuous passive motion treatment and acellular PLGA implants on osteochondral regeneration in the rabbit. <i>Biomaterials</i> , 2012, 33, 3153-3163. | 5.7 | 42 |
| 227 | The expression and prognostic significance of hepatoma-derived growth factor in oral cancer. <i>Oral Oncology</i> , 2012, 48, 629-635. | 0.8 | 31 |
| 228 | Overexpression of a Chromatin Remodeling Factor, RSF-1/HBXAP, Correlates with Aggressive Oral Squamous Cell Carcinoma. <i>American Journal of Pathology</i> , 2011, 178, 2407-2415. | 1.9 | 44 |
| 229 | Loss of lactate dehydrogenase B subunit expression is correlated with tumour progression and independently predicts inferior disease-specific survival in urinary bladder urothelial carcinoma. <i>Pathology</i> , 2011, 43, 707-712. | 0.3 | 21 |
| 230 | <i>Candida</i> invasion and influences in smoking patients with multiple oral leucoplakias – a retrospective study. <i>Mycoses</i> , 2011, 54, e377-83. | 1.8 | 20 |
| 231 | Rsf-1/HBXAP overexpression is associated with disease-specific survival of patients with gallbladder carcinoma. <i>Apmis</i> , 2011, 119, 808-814. | 0.9 | 22 |
| 232 | Involvement of phorbol-12-myristate-13-acetate-induced protein 1 in goniothalamin-induced TP53-dependent and -independent apoptosis in hepatocellular carcinoma-derived cells. <i>Toxicology and Applied Pharmacology</i> , 2011, 256, 8-23. | 1.3 | 20 |
| 233 | CCAAT/Enhancer-binding Protein β Mediates Tumor Necrosis Factor α -induced Aurora Kinase C Transcription and Promotes Genomic Instability. <i>Journal of Biological Chemistry</i> , 2011, 286, 28662-28670. | 1.6 | 26 |
| 234 | Overexpression of Nuclear Protein Kinase CK2 β Catalytic Subunit (CK2 β) as a Poor Prognosticator in Human Colorectal Cancer. <i>PLoS ONE</i> , 2011, 6, e17193. | 1.1 | 86 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | CKS1B Overexpression Implicates Clinical Aggressiveness of Hepatocellular Carcinomas but Not p27Kip1 Protein Turnover: an Independent Prognosticator with Potential p27Kip1-Independent Oncogenic Attributes?. <i>Annals of Surgical Oncology</i> , 2010, 17, 907-922. | 0.7 | 26 |
| 236 | Overexpression of Nuclear Protein Kinase CK2 $\hat{1}^2$ Subunit and Prognosis in Human Gastric Carcinoma. <i>Annals of Surgical Oncology</i> , 2010, 17, 1695-1702. | 0.7 | 44 |
| 237 | Prognostic Implication of Ezrin Overexpression in Myxofibrosarcomas. <i>Annals of Surgical Oncology</i> , 2010, 17, 3212-3219. | 0.7 | 39 |
| 238 | Annexin-I overexpression is associated with tumour progression and independently predicts inferior disease-specific and metastasis-free survival in urinary bladder urothelial carcinoma. <i>Pathology</i> , 2010, 42, 43-49. | 0.3 | 37 |
| 239 | Deciphering the transcriptional complex critical for RhoA gene expression and cancer metastasis. <i>Nature Cell Biology</i> , 2010, 12, 457-467. | 4.6 | 190 |
| 240 | ZBRK1 Acts as a Metastatic Suppressor by Directly Regulating <i>MMP9</i> in Cervical Cancer. <i>Cancer Research</i> , 2010, 70, 192-201. | 0.4 | 50 |
| 241 | CEBPD Reverses RB/E2F1-Mediated Gene Repression and Participates in HMDB-Induced Apoptosis of Cancer Cells. <i>Clinical Cancer Research</i> , 2010, 16, 5770-5780. | 3.2 | 41 |
| 242 | Prognostic implication of MET overexpression in myxofibrosarcomas: an integrative array comparative genomic hybridization, real-time quantitative PCR, immunoblotting, and immunohistochemical analysis. <i>Modern Pathology</i> , 2010, 23, 1379-1392. | 2.9 | 31 |
| 243 | Primary urothelial carcinoma of the upper tract: Important clinicopathological factors predicting bladder recurrence after surgical resection. <i>Pathology International</i> , 2009, 59, 642-649. | 0.6 | 50 |
| 244 | Ezrin overexpression in gastrointestinal stromal tumors: an independent adverse prognosticator associated with the non-gastric location. <i>Modern Pathology</i> , 2009, 22, 1351-1360. | 2.9 | 25 |
| 245 | Effect of S-Phase Kinase-Associated Protein 2 Expression on Distant Metastasis and Survival in Nasopharyngeal Carcinoma Patients. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009, 73, 202-207. | 0.4 | 36 |
| 246 | Activin A Enhances Prostate Cancer Cell Migration Through Activation of Androgen Receptor and Is Overexpressed in Metastatic Prostate Cancer. <i>Journal of Bone and Mineral Research</i> , 2009, 24, 1180-1193. | 3.1 | 47 |
| 247 | Homozygous Deletion of <i>MTAP</i> Gene as a Poor Prognosticator in Gastrointestinal Stromal Tumors. <i>Clinical Cancer Research</i> , 2009, 15, 6963-6972. | 3.2 | 41 |
| 248 | Flow Cytometric Analysis of DNA Ploidy and S-Phase Fraction in Primary Localized Myxofibrosarcoma: Correlations with Clinicopathological Factors, Skp2 Expression, and Patient Survival. <i>Annals of Surgical Oncology</i> , 2008, 15, 2239-2249. | 0.7 | 10 |
| 249 | Interleukin-20 targets renal mesangial cells and is associated with lupus nephritis. <i>Clinical Immunology</i> , 2008, 129, 277-285. | 1.4 | 38 |
| 250 | Heat Shock Protein 90 Overexpression Independently Predicts Inferior Disease-Free Survival with Differential Expression of the $\hat{1}^{\pm}$ and $\hat{1}^2$ Isoforms in Gastrointestinal Stromal Tumors. <i>Clinical Cancer Research</i> , 2008, 14, 7822-7831. | 3.2 | 40 |
| 251 | Shiunko and Acetylshikonin Promote Reepithelialization, Angiogenesis, and Granulation Tissue Formation in Wounded Skin. <i>The American Journal of Chinese Medicine</i> , 2008, 36, 115-123. | 1.5 | 40 |
| 252 | Malignant Diffuse-type Tenosynovial Giant Cell Tumors. <i>American Journal of Surgical Pathology</i> , 2008, 32, 587-599. | 2.1 | 57 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 253 | Skp2 is an independent prognosticator of gallbladder carcinoma among p27Kip1-interacting cell cycle regulators: an immunohistochemical study of 62 cases by tissue microarray. <i>Modern Pathology</i> , 2007, 20, 497-507. | 2.9 | 43 |
| 254 | Fatal HHV-8-Associated Hemophagocytic Syndrome in an HIV-Negative Immunocompetent Patient With Plasmablastic Variant of Multicentric Castleman Disease (Plasmablastic Microlymphoma). <i>American Journal of Surgical Pathology</i> , 2006, 30, 123-127. | 2.1 | 47 |
| 255 | Immunohistochemical Expression of p16INK4A, Ki-67, and Mcm2 Proteins in Gastrointestinal Stromal Tumors: Prognostic Implications and Correlations with Risk Stratification of NIH Consensus Criteria. <i>Annals of Surgical Oncology</i> , 2006, 13, 1633-1644. | 0.7 | 46 |
| 256 | p53 and p27 as predictors of clinical outcome for rectal-cancer patients receiving neoadjuvant therapy. <i>Surgical Oncology</i> , 2006, 15, 211-216. | 0.8 | 42 |
| 257 | Prognostic factors of myxofibrosarcomas: Implications of margin status, tumor necrosis, and mitotic rate on survival. <i>Journal of Surgical Oncology</i> , 2006, 93, 294-303. | 0.8 | 72 |
| 258 | Skp2 Overexpression Is Highly Representative of Intrinsic Biological Aggressiveness and Independently Associated with Poor Prognosis in Primary Localized Myxofibrosarcomas. <i>Clinical Cancer Research</i> , 2006, 12, 487-498. | 3.2 | 50 |
| 259 | Gastrointestinal stromal tumor (GIST) in southern Taiwan: A clinicopathologic study of 93 resected cases. <i>Pathology Research and Practice</i> , 2005, 201, 1-9. | 1.0 | 8 |
| 260 | Biological significance of chromosomal imbalance aberrations in gastrointestinal stromal tumors. <i>Journal of Biomedical Science</i> , 2004, 11, 65-71. | 2.6 | 37 |