## Christine Chen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6954579/publications.pdf

Version: 2024-02-01

932766 887659 1,117 23 10 17 citations h-index g-index papers 23 23 23 1726 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Fixed duration <i>vs.</i> prolonged duration treatment after first line therapy in patients with systemic light chain amyloidosis. Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis, 2022, 29, 23-30.	1.4	O
2	Selinexor, daratumumab, and dexamethasone in patients with relapsed or refractory multiple myeloma. EJHaem, 2021, 2, 56-65.	0.4	15
3	Computerized cognitive training in post-treatment hematological cancer survivors: a feasibility study. Pilot and Feasibility Studies, 2021, 7, 36.	0.5	6
4	Integrated safety profile of selinexor in multiple myeloma: experience from 437 patients enrolled in clinical trials. Leukemia, 2020, 34, 2430-2440.	3.3	54
5	Outcomes of patients with relapsed low-grade lymphoma retreated with rituximab are similar to rituximab naÃ-ve patients. Leukemia and Lymphoma, 2019, 60, 2576-2579.	0.6	О
6	Addition of Cyclophosphamide "On Demand―to Lenalidomide and Corticosteroids in Patients With Relapsed/Refractory Multiple Myeloma—A Retrospective Review of a Single-center Experience. Clinical Lymphoma, Myeloma and Leukemia, 2019, 19, e195-e203.	0.2	4
7	Single-center Experience in Treating Patients With t(4;14) Multiple Myeloma With and Without Planned Frontline Autologous Stem Cell Transplantation. Clinical Lymphoma, Myeloma and Leukemia, 2018, 18, 225-234.	0.2	7
8	Selinexor plus low-dose bortezomib and dexamethasone for patients with relapsed or refractory multiple myeloma. Blood, 2018, 132, 2546-2554.	0.6	110
9	Safety and efficacy of selinexor in relapsed or refractory multiple myeloma and Waldenstrom macroglobulinemia. Blood, 2018, 131, 855-863.	0.6	105
10	Secondary primary malignancies during the lenalidomide–dexamethasone regimen in relapsed/refractory multiple myeloma patients. Cancer Medicine, 2017, 6, 3-11.	1.3	24
11	Cyclophosphamide and Bortezomib With Prednisone or Dexamethasone for the Treatment of Relapsed and Refractory Multiple Myeloma. Clinical Lymphoma, Myeloma and Leukemia, 2016, 16, 387-394.	0.2	11
12	Pattern of First Relapse in Multiple Myeloma (MM) Patients (Pts) after a Cybord Induction Regimen and Autologous Stem Cell Transplantation (ASCT): Impact of Maintenance Therapy in the Real-World Setting. Blood, 2016, 128, 2137-2137.	0.6	3
13	Selinexor Shows Synergy in Combination with Pomalidomide and Low Dose Dexamethasone in Patients with Relapsed / Refractory Multiple Myeloma. Blood, 2016, 128, 3330-3330.	0.6	6
14	A Phase 1/2 Study of the Second Generation Selective Inhibitor of Nuclear Export (SINE) Compound, KPT-8602, in Patients with Relapsed Refractory Multiple Myeloma. Blood, 2016, 128, 4509-4509.	0.6	10
15	Myeloma Canada Research Network (MCRN)-001 ASCT Study of Busulfan + Melphalan (BuMel) Conditioning Followed By Lenalidomide (Len) Maintenance: Updated Results Including Serial Minimal Residual Disease (MRD) and Involved Serum Hevyliteâ,,¢ Chain (HLC) Ratio Assessments. Blood, 2016, 128, 4632-4632.	0.6	1
16	Selinexor in Combination with Bortezomib and Dexamethasone (SdB) Demonstrates Significant Activity in Patients with Refractory Multiple Myeloma (MM) Including Proteasome-Inhibitor Refractory Patients: Results of the Phase I Stomp Trial. Blood, 2016, 128, 977-977.	0.6	15
17	Single Centre Experience in Treating Newly Diagnosed t(4;14) Multiple Myeloma with and without Planned Front-Line Autologous Stem Cell Transplant. Blood, 2016, 128, 3457-3457.	0.6	O
18	Is Prior Rituximab Exposure Associated with a Difference in Outcomes in Relapsed Low-Grade Lymphomas Retreated with Rituximab in Comparison to Rituximab NaÃ'Ve Patients?. Blood, 2016, 128, 1795-1795.	0.6	0

#	Article	IF	CITATIONS
19	Cost Analysis of Stored Autologous Peripheral Blood Stem Cells for a Second Autologous Transplantation in Multiple Myeloma Patients: A Markov Model. Blood, 2016, 128, 1184-1184.	0.6	0
20	Absolute lymphocyte count as predictor of overall survival for patients with multiple myeloma treated with single autologous stem cell transplant. Leukemia and Lymphoma, 2015, 56, 2668-2673.	0.6	27
21	Tolerability of Velcade (Bortezomib) subcutaneous administration using a maximum volume of 3 mL per injection site. Journal of Oncology Pharmacy Practice, 2015, 21, 285-292.	0.5	9
22	Myeloma Canada Research Network (MCRN)-001 Trial Utilizing Bortezomib (btz)-Based Induction, Enhanced Conditioning with IV Busulfan + Melphalan (BuMel) and Lenalidomide (len) Maintenance in Multiple Myeloma Patients Eligible for Autologous Stem Cell Transplant (ASCT): A National Canadian Study Evaluating Achievement of Minimal Residual Disease (MRD) Negativity and Involved Serum	0.6	0
23	HevyliteTMÃ, chain (HLC) Normalization. Blood, 2015, 126, 1982-1982.  Pomalidomide plus low-dose dexamethasone versus high-dose dexamethasone alone for patients with relapsed and refractory multiple myeloma (MM-003): a randomised, open-label, phase 3 trial. Lancet Oncology, The, 2013, 14, 1055-1066.	5.1	710