Jonathan Mant

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6955830/jonathan-mant-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

211 8,382 37 87 g-index

228 10,266 ext. papers ext. citations avg, IF 5.76

L-index

#	Paper	IF	Citations
211	The development of a self-management intervention for stroke survivors - My Life After Stroke (MLAS) <i>Disability and Rehabilitation</i> , 2022 , 1-9	2.4	
210	Factors influencing follow-up care post-TIA and minor stroke: a qualitative study using the theoretical domains framework <i>BMC Health Services Research</i> , 2022 , 22, 235	2.9	
209	Cost-Effectiveness of Antihypertensive Deprescribing in Primary Care: a Markov Modelling Study Using Data From the OPTiMISE Trial <i>Hypertension</i> , 2022 , HYPERTENSIONAHA12118726	8.5	O
208	Why do people take part in atrial fibrillation screening? Qualitative interview study in English primary care <i>BMJ Open</i> , 2022 , 12, e051703	3	0
207	Wearable Photoplethysmography for Cardiovascular Monitoring <i>Proceedings of the IEEE</i> , 2022 , 110, 355-381	14.3	4
206	Statin use is associated with lower risk of dementia in stroke patients: a community-based cohort study with inverse probability weighted marginal structural model analysis <i>European Journal of Epidemiology</i> , 2022 , 1	12.1	O
205	Statistical reanalysis of vascular event outcomes in primary and secondary vascular prevention trials. <i>BMC Medical Research Methodology</i> , 2021 , 21, 218	4.7	O
204	Impact of changes to national guidelines on hypertension-related workload: an interrupted time series analysis in English primary care. <i>British Journal of General Practice</i> , 2021 , 71, e296-e302	1.6	1
203	GPsRmindlines on deprescribing antihypertensives in older patients with multimorbidity: a qualitative study in English general practice. <i>British Journal of General Practice</i> , 2021 , 71, e498-e507	1.6	O
202	Blood Pressure Changes Following Antihypertensive Medication Reduction, by Drug Class and Dose Chosen for Withdrawal: Exploratory Analysis of Data From the OPTiMISE Trial. <i>Frontiers in Pharmacology</i> , 2021 , 12, 619088	5.6	0
201	Stroke risk following traumatic brain injury: Systematic review and meta-analysis. <i>International Journal of Stroke</i> , 2021 , 16, 370-384	6.3	1
200	Delivering screening programmes in primary care: protocol for a scoping and systematic mixed studies review. <i>BMJ Open</i> , 2021 , 11, e046331	3	
199	The feasibility of the PAM intervention to support treatment-adherence in people with hypertension in primary care: a randomised clinical controlled trial. <i>Scientific Reports</i> , 2021 , 11, 8897	4.9	1
198	Home blood pressure monitoring: methodology, clinical relevance and practical application: a 2021 position paper by the Working Group on Blood Pressure Monitoring and Cardiovascular Variability of the European Society of Hypertension. <i>Journal of Hypertension</i> , 2021 , 39, 1742-1767	1.9	15
197	What underlies the difference between self-reported health and disability after stroke? A qualitative study in the UK. <i>BMC Neurology</i> , 2021 , 21, 315	3.1	1
196	Frailty and cerebrovascular disease: Concepts and clinical implications for stroke medicine. <i>International Journal of Stroke</i> , 2021 , 17474930211034331	6.3	3
195	Anticoagulation trends in adults aged 65 years and over with atrial fibrillation: a cohort study. <i>Open Heart</i> , 2021 , 8,	3	6

194	Polypills with or without aspirin for primary prevention of cardiovascular disease. <i>Lancet, The</i> , 2021 , 398, 1106-1107	40	
193	Accuracy of blood-pressure monitors owned by patients with hypertension (ACCU-RATE study): a cross-sectional, observational study in central England. <i>British Journal of General Practice</i> , 2020 , 70, e54	18 ^{1.6} 554	1 ¹¹
192	How primary care can help survivors of transient ischaemic attack and stroke return to work: focus groups with stakeholders from a UK community. <i>British Journal of General Practice</i> , 2020 , 70, e294-e30.	2 ^{1.6}	2
191	Acquiring wearable photoplethysmography data in daily life: the PPG Diary Pilot Study 2020 , 2, 80		5
190	Screening for Atrial Fibrillation: Improving Efficiency of Manual Review of Handheld Electrocardiograms. <i>Engineering Proceedings</i> , 2020 , 2, 78	0.5	O
189	Temporal trends in incidence of atrial fibrillation in primary care records: a population-based cohort study. <i>BMJ Open</i> , 2020 , 10, e042518	3	3
188	Screening for Atrial Fibrillation: Improving Efficiency of Manual Review of Handheld Electrocardiograms. <i>Engineering Proceedings</i> , 2020 , 2, 78	0.5	
187	Acquiring Wearable Photoplethysmography Data in Daily Life: The PPG Diary Pilot Study. <i>Engineering Proceedings</i> , 2020 , 2, 80	0.5	4
186	Home Blood Pressure Monitoring for Treatment Titration. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2020 , 63-72	0.1	
185	"It& Difficult, There& No Formula": Qualitative Study of Stroke Related Communication Between Primary and Secondary Healthcare Professionals. <i>International Journal of Integrated Care</i> , 2020 , 20, 11	2	2
184	Infographic. Developing home-based cardiac rehabilitation for people post-transient ischaemic attack (TIA) and ischaemic stroke. <i>British Journal of Sports Medicine</i> , 2020 , 54, 487-488	10.3	
183	Impact of Changes to National Hypertension Guidelines on Hypertension Management and Outcomes in the United Kingdom. <i>Hypertension</i> , 2020 , 75, 356-364	8.5	7
182	Generalizability of Blood Pressure Lowering Trials to Older Patients: Cross-Sectional Analysis. <i>Journal of the American Geriatrics Society</i> , 2020 , 68, 2508-2515	5.6	16
181	Improving Primary Care After Stroke (IPCAS) randomised controlled trial: protocol for a multidimensional process evaluation. <i>BMJ Open</i> , 2020 , 10, e036879	3	О
180	Usability and Acceptability of a Novel Secondary Prevention Initiative Targeting Physical Activity for Individuals after a Transient Ischaemic Attack or "Minor" Stroke: A Qualitative Study. International Journal of Environmental Research and Public Health, 2020, 17,	4.6	2
179	Deprescribing Antihypertensive Medication in Elderly Adults-Reply. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 324, 1682-1683	27.4	3
178	Association of Prior Atherosclerotic Cardiovascular Disease with Dementia After Stroke: A Retrospective Cohort Study. <i>Journal of Alzheimerm Disease</i> , 2020 , 77, 1157-1167	4.3	4
177	Association of blood lipids, atherosclerosis and statin use with dementia and cognitive impairment after stroke: A systematic review and meta-analysis. <i>Ageing Research Reviews</i> , 2020 , 57, 100962	12	18

176	When is referral from primary care to specialist services appropriate for survivors of stroke? A modified RAND-appropriateness consensus study. <i>BMC Family Practice</i> , 2020 , 21, 66	2.6	1
175	Characteristics, service use and mortality of clusters of multimorbid patients in England: a population-based study. <i>BMC Medicine</i> , 2020 , 18, 78	11.4	18
174	Effect of Antihypertensive Medication Reduction vs Usual Care on Short-term Blood Pressure Control in Patients With Hypertension Aged 80 Years and Older: The OPTIMISE Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 323, 2039-2051	27.4	40
173	Estimated stroke risk, yield, and number needed to screen for atrial fibrillation detected through single time screening: a multicountry patient-level meta-analysis of 141,220 screened individuals. <i>PLoS Medicine</i> , 2019 , 16, e1002903	11.6	56
172	Schedules for Self-monitoring Blood Pressure: A Systematic Review. <i>American Journal of Hypertension</i> , 2019 , 32, 350-364	2.3	5
171	The role of the GP in managing suspected transient ischaemic attack: a qualitative study. <i>BMC Family Practice</i> , 2019 , 20, 67	2.6	1
170	Cost-Effectiveness of Telemonitoring and Self-Monitoring of Blood Pressure for Antihypertensive Titration in Primary Care (TASMINH4). <i>Hypertension</i> , 2019 , 73, 1231-1239	8.5	41
169	Opportunities and challenges of a novel cardiac output response to stress (CORS) test to enhance diagnosis of heart failure in primary care: qualitative study. <i>BMJ Open</i> , 2019 , 9, e028122	3	1
168	Older People Might Be at Most Serious Risk in Antihypertensive Treatment-Reply. <i>JAMA Internal Medicine</i> , 2019 , 179, 590-591	11.5	
167	Longer-term efficiency and safety of increasing the frequency of whole blood donation (INTERVAL): extension study of a randomised trial of 20 757 blood donors. <i>Lancet Haematology,the</i> , 2019 , 6, e510-e520	14.6	10
166	Statin use and high-dose statin use after ischemic stroke in the UK: a retrospective cohort study. <i>Clinical Epidemiology</i> , 2019 , 11, 495-508	5.9	10
165	Does characterising patterns of multimorbidity in stroke matter for developing collaborative care approaches in primary care?. <i>Primary Health Care Research and Development</i> , 2019 , 20, e110	1.6	1
164	Optimising Management of Patients with Heart Failure with Preserved Ejection Fraction in Primary Care (OPTIMISE-HFpEF): rationale and protocol for a multi-method study. <i>BJGP Open</i> , 2019 ,	3.1	4
163	How is poststroke fatigue understood by stroke survivors and carers? A thematic analysis of an online discussion forum. <i>BMJ Open</i> , 2019 , 9, e028958	3	4
162	Conceptualising post-stroke fatigue: a cross-sectional survey of UK-based physiotherapists and occupational therapists. <i>BMJ Open</i> , 2019 , 9, e033066	3	8
161	Rehabilitation of patients after transient ischaemic attack or minor stroke: pilot feasibility randomised trial of a home-based prevention programme. <i>British Journal of General Practice</i> , 2019 , 69, e706-e714	1.6	9
160	Improving Primary Care After Stroke (IPCAS) trial: protocol of a randomised controlled trial to evaluate a novel model of care for stroke survivors living in the community. <i>BMJ Open</i> , 2019 , 9, e03028.	53	5
159	TIA and minor stroke: a qualitative study of long-term impact and experiences of follow-up care. <i>BMC Family Practice</i> , 2019 , 20, 176	2.6	18

158	Using a checklist to facilitate management of long-term care needs after stroke: insights from focus groups and a feasibility study. <i>BMC Family Practice</i> , 2019 , 20, 2	2.6	13
157	Functional and emotional outcomes after transient ischemic attack: A 12-month prospective controlled cohort study. <i>International Journal of Stroke</i> , 2019 , 14, 522-529	6.3	8
156	Efficacy of self-monitored blood pressure, with or without telemonitoring, for titration of antihypertensive medication (TASMINH4): an unmasked randomised controlled trial. <i>Lancet, The</i> , 2018 , 391, 949-959	40	179
155	Rivaroxaban plus aspirin, compared with aspirin alone, reduced cardiovascular events in patients with stable peripheral or carotid artery disease, but increased the risk of major bleeding. <i>BMJ Evidence-Based Medicine</i> , 2018 , 23, 191-192	2.7	O
154	Online stroke forum as source of data for qualitative research: insights from a comparison with patients Rinterviews. <i>BMJ Open</i> , 2018 , 8, e020133	3	19
153	Evaluating practical support stroke survivors get with medicines and unmet needs in primary care: a survey. <i>BMJ Open</i> , 2018 , 8, e019874	3	6
152	Do patients actually do what we ask: patient fidelity and persistence to the Targets and Self-Management for the Control of Blood Pressure in Stroke and at Risk Groups blood pressure self-management intervention. <i>Journal of Hypertension</i> , 2018 , 36, 1753-1761	1.9	10
151	What are the implications for practice that arise from studies of medication taking? A systematic review of qualitative research. <i>PLoS ONE</i> , 2018 , 13, e0195076	3.7	4
150	Delivery and impact of the NHS Health Check in the first 8 years: a systematic review. <i>British Journal of General Practice</i> , 2018 , 68, e449-e459	1.6	21
149	Stroke survivorsRand informal caregiversRexperiences of primary care and community healthcare services - A systematic review and meta-ethnography. <i>PLoS ONE</i> , 2018 , 13, e0192533	3.7	87
148	Predicting Out-of-Office Blood Pressure in the Clinic for the Diagnosis of Hypertension in Primary Care: An Economic Evaluation. <i>Hypertension</i> , 2018 , 71, 250-261	8.5	7
147	Advanced heart failure: setting the research agenda. British Journal of Cardiac Nursing, 2018, 13, 243-2	435.2	
146	OPtimising Treatment for MIld Systolic hypertension in the Elderly (OPTiMISE): protocol for a randomised controlled non-inferiority trial. <i>BMJ Open</i> , 2018 , 8, e022930	3	8
145	Reasons why people do not attend NHS Health Checks: a systematic review and qualitative synthesis. <i>British Journal of General Practice</i> , 2018 , 68, e28-e35	1.6	32
144	Benefits and Harms of Antihypertensive Treatment in Low-Risk Patients With Mild Hypertension. JAMA Internal Medicine, 2018 , 178, 1626-1634	11.5	42
143	Association of guideline and policy changes with incidence of lifestyle advice and treatment for uncomplicated mild hypertension in primary care: a longitudinal cohort study in the Clinical Practice Research Datalink. <i>BMJ Open</i> , 2018 , 8, e021827	3	7
142	A novel cardiac output response to stress test developed to improve diagnosis and monitoring of heart failure in primary care. <i>ESC Heart Failure</i> , 2018 , 5, 703-712	3.7	5
141	Prospective external validation of the Predicting Out-of-OFfice Blood Pressure (PROOF-BP) strategy for triaging ambulatory monitoring in the diagnosis and management of hypertension: observational cohort study. <i>BMJ</i> , <i>The</i> , 2018 , 361, k2478	5.9	6

140	Training in positivity for stroke? A qualitative study of acceptability of use of Positive Mental Training (PosMT) as a tool to assist stroke survivors with post-stroke psychological problems and in coping with rehabilitation. <i>NeuroRehabilitation</i> , 2017 , 40, 259-270	2	9
139	Measurement of blood pressure for the diagnosis and management of hypertension in different ethnic groups: one size fits all. <i>BMC Cardiovascular Disorders</i> , 2017 , 17, 55	2.3	7
138	How do stroke survivors and their carers use practitioners Radvice on secondary prevention medications? Qualitative study of an online forum. <i>Family Practice</i> , 2017 , 34, 612-620	1.9	8
137	Interarm Difference in Systolic Blood Pressure in Different Ethnic Groups and Relationship to the "White Coat Effect": A Cross-Sectional Study. <i>American Journal of Hypertension</i> , 2017 , 30, 884-891	2.3	8
136	MICE or NICE? An economic evaluation of clinical decision rules in the diagnosis of heart failure in primary care. <i>International Journal of Cardiology</i> , 2017 , 241, 255-261	3.2	5
135	Intensive versus Guideline Blood Pressure and Lipid Lowering in Patients with Previous Stroke: Main Results from the Pilot Prevention of Decline in Cognition after Stroke TrialR(PODCAST) Randomised Controlled Trial. <i>PLoS ONE</i> , 2017 , 12, e0164608	3.7	41
134	Cost-effectiveness analysis of use of a polypill versus usual care or best practice for primary prevention in people at high risk of cardiovascular disease. <i>PLoS ONE</i> , 2017 , 12, e0182625	3.7	11
133	Stroke Prevention Rehabilitation Intervention Trial of Exercise (SPRITE) - a randomised feasibility study. <i>BMC Cardiovascular Disorders</i> , 2017 , 17, 290	2.3	20
132	Self-monitoring of blood pressure in hypertension: A systematic review and individual patient data meta-analysis. <i>PLoS Medicine</i> , 2017 , 14, e1002389	11.6	222
131	Efficiency and safety of varying the frequency of whole blood donation (INTERVAL): a randomised trial of 45 000 donors. <i>Lancet, The</i> , 2017 , 390, 2360-2371	40	140
130	Association Between More Intensive vs Less Intensive Blood Pressure Lowering and Risk of Mortality in Chronic Kidney Disease Stages 3 to 5: A Systematic Review and Meta-analysis. <i>JAMA Internal Medicine</i> , 2017 , 177, 1498-1505	11.5	98
129	Barriers and facilitators to adherence to secondary stroke prevention medications after stroke: analysis of survivors and caregivers views from an online stroke forum. <i>BMJ Open</i> , 2017 , 7, e016814	3	22
128	Predictive values of referrals for transient ischaemic attack from first-contact health care: a systematic review. <i>British Journal of General Practice</i> , 2017 , 67, e871-e880	1.6	5
127	When has service provision for transient ischaemic attack improved enough? A discrete event simulation economic modelling study. <i>BMJ Open</i> , 2017 , 7, e018189	3	
126	Cost-Effectiveness Analysis of Natriuretic Peptide Testing and Specialist Management in Patients with Suspected Acute Heart Failure. <i>Value in Health</i> , 2017 , 20, 1025-1033	3.3	6
125	Telemonitoring and/or self-monitoring of blood pressure in hypertension (TASMINH4): protocol for a randomised controlled trial. <i>BMC Cardiovascular Disorders</i> , 2017 , 17, 58	2.3	13
124	Screening for atrial fibrillation: a European Heart Rhythm Association (EHRA) consensus document endorsed by the Heart Rhythm Society (HRS), Asia Pacific Heart Rhythm Society (APHRS), and Sociedad Latinoamericana de Estimulaci Cardica y Electrofisiologi (SOLAECE). Europace, 2017,	3.9	135
123	19, 1589-1623 Patient experience of NHS health checks: a systematic review and qualitative synthesis. <i>BMJ Open</i> , 2017 , 7, e017169	3	19

122	Views of commissioners, managers and healthcare professionals on the NHS Health Check programme: a systematic review. <i>BMJ Open</i> , 2017 , 7, e018606	3	8
121	Accuracy of monitors used for blood pressure checks in English retail pharmacies: a cross-sectional observational study. <i>British Journal of General Practice</i> , 2016 , 66, e309-14	1.6	7
120	Self-monitoring blood pressure in patients with hypertension: an internet-based survey of UK GPs. British Journal of General Practice, 2016 , 66, e831-e837	1.6	23
119	Primary care interventions and current service innovations in modifying long-term outcomes after stroke: a protocol for a scoping review. <i>BMJ Open</i> , 2016 , 6, e012840	3	2
118	Impact of socioeconomic deprivation on screening for cardiovascular disease risk in a primary prevention population: a cross-sectional study. <i>BMJ Open</i> , 2016 , 6, e009984	3	19
117	Barriers and facilitators to staying in work after stroke: insight from an online forum. <i>BMJ Open</i> , 2016 , 6, e009974	3	35
116	Influence of ethnicity on acceptability of method of blood pressure monitoring: a cross-sectional study in primary care. <i>British Journal of General Practice</i> , 2016 , 66, e577-86	1.6	20
115	Stroke survivors and their families receive information and support on an individual basis from an online forum: descriptive analysis of a population of 2348 patients and qualitative study of a sample of participants. <i>BMJ Open</i> , 2016 , 6, e010501	3	34
114	Stroke survivorsRcaregiversRand GPsRattitudes towards a polypill for the secondary prevention of stroke: a qualitative interview study. <i>BMJ Open</i> , 2016 , 6, e010458	3	8
113	Cost-effectiveness analysis of different systolic blood pressure targets for people with a history of stroke or transient ischaemic attack: Economic analysis of the PAST-BP study. <i>European Journal of Preventive Cardiology</i> , 2016 , 23, 1590-8	3.9	5
112	Assessment of cardiovascular risk in primary care. <i>BMJ, The</i> , 2016 , 353, i3022	5.9	
111	Predicting Out-of-Office Blood Pressure in the Clinic (PROOF-BP): Derivation and Validation of a Tool to Improve the Accuracy of Blood Pressure Measurement in Clinical Practice. <i>Hypertension</i> , 2016 , 67, 941-50	8.5	24
110	Different systolic blood pressure targets for people with history of stroke or transient ischaemic attack: PAST-BP (Prevention After StrokeBlood Pressure) randomised controlled trial. <i>BMJ, The</i> , 2016 , 352, i708	5.9	37
109	Cost-effectiveness of self-management of blood pressure in hypertensive patients over 70 years with suboptimal control and established cardiovascular disease or additional cardiovascular risk diseases (TASMIN-SR). <i>European Journal of Preventive Cardiology</i> , 2016 , 23, 902-12	3.9	25
108	Effects of intensive blood pressure lowering on cardiovascular and renal outcomes: updated systematic review and meta-analysis. <i>Lancet, The</i> , 2016 , 387, 435-43	40	591
107	An Occupational Therapy intervention for residents with stroke-related disabilities in UK Care Homes (OTCH): cluster randomised controlled trial with economic evaluation. <i>Health Technology Assessment</i> , 2016 , 20, 1-138	4.4	8
106	The Stroke Prevention Programme: a programme of research to inform optimal stroke prevention in primary care. <i>Programme Grants for Applied Research</i> , 2016 , 4, 1-290	1.5	1
105	Relationship of Self-Rated Health to Stroke Incidence and Mortality in Older Individuals with and without a History of Stroke: A Longitudinal Study of the MRC Cognitive Function and Ageing (CFAS) Population. <i>PLoS ONE</i> , 2016 , 11, e0150178	3.7	9

104	PatientsRand cliniciansRviews on the optimum schedules for self-monitoring of blood pressure: a qualitative focus group and interview study. <i>British Journal of General Practice</i> , 2016 , 66, e819-e830	1.6	19
103	Barriers to medication adherence for the secondary prevention of stroke: a qualitative interview study in primary care. <i>British Journal of General Practice</i> , 2016 , 66, e568-76	1.6	31
102	The diagnostic accuracy of the natriuretic peptides in heart failure: systematic review and diagnostic meta-analysis in the acute care setting. <i>BMJ, The</i> , 2015 , 350, h910	5.9	196
101	Self-Screening and Non-Physician Screening for Hypertension in Communities: A Systematic Review. <i>American Journal of Hypertension</i> , 2015 , 28, 1316-24	2.3	36
100	Ethnicity and differences between clinic and ambulatory blood pressure measurements. <i>American Journal of Hypertension</i> , 2015 , 28, 729-38	2.3	15
99	The association between prehospital care and in-hospital treatment decisions in acute stroke: a cohort study. <i>Emergency Medicine Journal</i> , 2015 , 32, 93-9	1.5	34
98	Individual patient data meta-analysis of self-monitoring of blood pressure (BP-SMART): a protocol. <i>BMJ Open</i> , 2015 , 5, e008532	3	9
97	Baseline characteristics, analysis plan and report on feasibility for the Prevention Of Decline in Cognition After Stroke Trial (PODCAST). <i>Trials</i> , 2015 , 16, 509	2.8	5
96	The associations of "fatness," "fitness," and physical activity with all-cause mortality in older adults: A systematic review. <i>Obesity</i> , 2015 , 23, 1944-56	8	22
95	An occupational therapy intervention for residents with stroke related disabilities in UK care homes (OTCH): cluster randomised controlled trial. <i>BMJ, The</i> , 2015 , 350, h468	5.9	28
94	PatientsRviews about taking a polypill to manage cardiovascular risk: a qualitative study in primary care. <i>British Journal of General Practice</i> , 2015 , 65, e447-53	1.6	11
93	Adherence to medication in stroke survivors dependent on caregivers. <i>British Journal of General Practice</i> , 2015 , 65, e789-91	1.6	7
92	Decisions and delays within stroke patients Proute to the hospital: a qualitative study. <i>Annals of Emergency Medicine</i> , 2015 , 65, 279-287.e3	2.1	21
91	Warfarin versus aspirin for prevention of cognitive decline in atrial fibrillation: randomized controlled trial (Birmingham Atrial Fibrillation Treatment of the Aged Study). <i>Stroke</i> , 2014 , 45, 1381-6	6.7	45
90	Diagnosing and managing acute heart failure in adults: summary of NICE guidance. <i>BMJ, The</i> , 2014 , 349, g5695	5.9	14
89	Cost-effectiveness of optimizing acute stroke care services for thrombolysis. <i>Stroke</i> , 2014 , 45, 553-62	6.7	27
88	The drugs do work: blood pressure improvement in England. <i>Lancet, The</i> , 2014 , 383, 1868-9	40	1
87	Missed opportunities in prevention of cardiovascular disease in primary care: a cross-sectional study. <i>British Journal of General Practice</i> , 2014 , 64, e38-46	1.6	26

(2013-2014)

86	Making sense of patientsRinternet forums: a systematic method using discourse analysis. <i>British Journal of General Practice</i> , 2014 , 64, e178-80	1.6	16
85	Relationship of self-rated health with fatal and non-fatal outcomes in cardiovascular disease: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2014 , 9, e103509	3.7	88
84	Telemonitoring and self-management in the control of hypertension (TASMINH2): a cost-effectiveness analysis. <i>European Journal of Preventive Cardiology</i> , 2014 , 21, 1517-30	3.9	63
83	Self-monitoring in hypertension: a web-based survey of primary care physicians. <i>Journal of Human Hypertension</i> , 2014 , 28, 123-7	2.6	29
82	Prevalence and costs of treating uncomplicated stage 1 hypertension in primary care: a cross-sectional analysis. <i>British Journal of General Practice</i> , 2014 , 64, e641-8	1.6	8
81	Medication taking in coronary artery disease: a systematic review and qualitative synthesis. <i>Annals of Family Medicine</i> , 2014 , 12, 224-32	2.9	33
80	Point-of-care cluster randomized trial in stroke secondary prevention using electronic health records. <i>Stroke</i> , 2014 , 45, 2066-71	6.7	20
79	Effect of self-monitoring and medication self-titration on systolic blood pressure in hypertensive patients at high risk of cardiovascular disease: the TASMIN-SR randomized clinical trial. <i>JAMA - Journal of the American Medical Association</i> , 2014 , 312, 799-808	27.4	227
78	Response to letter regarding article, "warfarin versus aspirin for prevention of cognitive decline in atrial fibrillation: randomized controlled trial (birmingham atrial fibrillation treatment of the aged study)". <i>Stroke</i> , 2014 , 45, e192	6.7	1
77	Predicting out-of-office blood pressure level using repeated measurements in the clinic: an observational cohort study. <i>Journal of Hypertension</i> , 2014 , 32, 2171-8; discussion 2178	1.9	20
76	AuthorsRreply to Dalzell and Cannon and Satchithananda and colleagues. <i>BMJ, The</i> , 2014 , 349, g6703	5.9	
75	RApid Primary care Initiation of Drug treatment for Transient Ischaemic Attack (RAPID-TIA): study protocol for a pilot randomised controlled trial. <i>Trials</i> , 2013 , 14, 194	2.8	5
74	Targets and self-management for the control of blood pressure in stroke and at risk groups (TASMIN-SR): protocol for a randomised controlled trial. <i>BMC Cardiovascular Disorders</i> , 2013 , 13, 21	2.3	14
73	What determines the self-rated health of older individuals with stroke compared to other older individuals? A cross-sectional analysis of the Medical Research Council Cognitive Function and Aging Study. <i>BMC Geriatrics</i> , 2013 , 13, 85	4.1	13
72	A systematic evaluation of stroke surveillance studies in low- and middle-income countries. <i>Neurology</i> , 2013 , 80, 677-84	6.5	28
71	Potential impact of new oral anticoagulants on the management of atrial fibrillation-related stroke in primary care. <i>International Journal of Clinical Practice</i> , 2013 , 67, 647-55	2.9	3
70	Prevention of Decline in Cognition after Stroke Trial (PODCAST): a study protocol for a factorial randomised controlled trial of intensive versus guideline lowering of blood pressure and lipids. <i>Trials</i> , 2013 , 14, 401	2.8	13
69	Would primary healthcare professionals prescribe a polypill to manage cardiovascular risk? A qualitative interview study. <i>BMJ Open</i> , 2013 , 3,	3	18

68	Transient neurological symptoms in the older population: report of a prospective cohort studythe Medical Research Council Cognitive Function and Ageing Study (CFAS). <i>BMJ Open</i> , 2013 , 3,	3	9
67	Thromboprophylaxis of elderly patients with AF in the UK: an analysis using the General Practice Research Database (GPRD) 2000-2009. <i>Heart</i> , 2013 , 99, 127-32	5.1	66
66	Cost-effectiveness analysis of serial measurement of circulating natriuretic peptide concentration in chronic heart failure. <i>Heart</i> , 2013 , 99, 267-71	5.1	11
65	Trials to improve blood pressure through adherence to antihypertensives in stroke/TIA: systematic review and meta-analysis. <i>Journal of the American Heart Association</i> , 2013 , 2, e000251	6	17
64	Something for the weekend?. JAMA Neurology, 2013, 70, 130	17.2	1
63	Adherence to cardiovascular therapy: a meta-analysis of prevalence and clinical consequences. <i>European Heart Journal</i> , 2013 , 34, 2940-8	9.5	489
62	Self-monitoring for improving control of blood pressure in patients with hypertension. <i>The Cochrane Library</i> , 2013 ,	5.2	2
61	Patient self-monitoring of blood pressure and self-titration of medication in primary care: the TASMINH2 trial qualitative study of health professionalsRexperiences. <i>British Journal of General Practice</i> , 2013 , 63, e378-85	1.6	29
60	Accuracy of ambulatory blood pressure monitors: a systematic review of validation studies. <i>Journal of Hypertension</i> , 2013 , 31, 239-50	1.9	36
59	Do differences in blood pressure between arms matter?. <i>Lancet, The</i> , 2012 , 379, 872-873	40	9
58	A cross-sectional study of quality of life in an elderly population (75 years and over) with atrial fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the Aged study. <i>Europace</i> , 2012 , 14, 1420-7	3.9	21
58 57	fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the	3.9	21
	fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the Aged study. <i>Europace</i> , 2012 , 14, 1420-7 The REFER (REFer for EchocaRdiogram) protocol: a prospective validation of a clinical decision rule, NT-proBNP, or their combination, in the diagnosis of heart failure in primary care. Rationale and		
57	fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the Aged study. <i>Europace</i> , 2012 , 14, 1420-7 The REFER (REFer for EchocaRdiogram) protocol: a prospective validation of a clinical decision rule, NT-proBNP, or their combination, in the diagnosis of heart failure in primary care. Rationale and design. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 97 A cluster randomised controlled trial of an occupational therapy intervention for residents with	2.3	13
57 56	fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the Aged study. <i>Europace</i> , 2012 , 14, 1420-7 The REFER (REFer for EchocaRdiogram) protocol: a prospective validation of a clinical decision rule, NT-proBNP, or their combination, in the diagnosis of heart failure in primary care. Rationale and design. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 97 A cluster randomised controlled trial of an occupational therapy intervention for residents with stroke living in UK care homes (OTCH): study protocol. <i>BMC Neurology</i> , 2012 , 12, 52 PatientsRexperiences of self-monitoring blood pressure and self-titration of medication: the	2.3	13
57 56 55	fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the Aged study. <i>Europace</i> , 2012 , 14, 1420-7 The REFER (REFer for EchocaRdiogram) protocol: a prospective validation of a clinical decision rule, NT-proBNP, or their combination, in the diagnosis of heart failure in primary care. Rationale and design. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 97 A cluster randomised controlled trial of an occupational therapy intervention for residents with stroke living in UK care homes (OTCH): study protocol. <i>BMC Neurology</i> , 2012 , 12, 52 PatientsRexperiences of self-monitoring blood pressure and self-titration of medication: the TASMINH2 trial qualitative study. <i>British Journal of General Practice</i> , 2012 , 62, e135-42 Development and initial validation of a simple clinical decision tool to predict the presence of heart failure in primary care: the MICE (Male, Infarction, Crepitations, Edema) rule. <i>European Journal of</i>	2.3 3.1 1.6	13 14 65
57 56 55 54	fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation Treatment of the Aged study. <i>Europace</i> , 2012 , 14, 1420-7 The REFER (REFer for EchocaRdiogram) protocol: a prospective validation of a clinical decision rule, NT-proBNP, or their combination, in the diagnosis of heart failure in primary care. Rationale and design. <i>BMC Cardiovascular Disorders</i> , 2012 , 12, 97 A cluster randomised controlled trial of an occupational therapy intervention for residents with stroke living in UK care homes (OTCH): study protocol. <i>BMC Neurology</i> , 2012 , 12, 52 PatientsRexperiences of self-monitoring blood pressure and self-titration of medication: the TASMINH2 trial qualitative study. <i>British Journal of General Practice</i> , 2012 , 62, e135-42 Development and initial validation of a simple clinical decision tool to predict the presence of heart failure in primary care: the MICE (Male, Infarction, Crepitations, Edema) rule. <i>European Journal of Heart Failure</i> , 2012 , 14, 1000-8 Protocol for an observation and implementation study investigating optimisation of the	2.3 3.1 1.6	13 14 65 23

(2009-2011)

50	Republished technology and guidelines: the diagnosis and management of chronic heart failure: review following the publication of the NICE guidelines. <i>Postgraduate Medical Journal</i> , 2011 , 87, 841-6	2	О
49	The diagnosis and management of chronic heart failure: review following the publication of the NICE guidelines. <i>Heart</i> , 2011 , 97, 411-6	5.1	25
48	Cost-effectiveness of options for the diagnosis of high blood pressure in primary care: a modelling study. <i>Lancet, The</i> , 2011 , 378, 1219-30	40	237
47	Management of chronic heart failure in adults: synopsis of the National Institute For Health and clinical excellence guideline. <i>Annals of Internal Medicine</i> , 2011 , 155, 252-9	8	38
46	Measurement of blood pressure in primary care. <i>BMJ, The</i> , 2011 , 342, d382	5.9	3
45	Cost effectiveness of warfarin versus aspirin in patients older than 75 years with atrial fibrillation. <i>Stroke</i> , 2011 , 42, 1717-21	6.7	19
44	Impact of study design on recruitment of patients to a primary care trial: an observational time series analysis of the Birmingham Atrial Fibrillation Treatment of the Aged (BAFTA) study. <i>Family Practice</i> , 2010 , 27, 691-7	1.9	7
43	Primary care research and clinical practice: cardiovascular disease. <i>Postgraduate Medical Journal</i> , 2010 , 86, 696-703	2	6
42	Telemonitoring and self-management in the control of hypertension (TASMINH2): a randomised controlled trial. <i>Lancet, The</i> , 2010 , 376, 163-72	40	381
41	TelemonitoringBr better follow-up? [AuthorsRreply. Lancet, The, 2010, 376, 1737-1738	40	
41	TelemonitoringBr better follow-up? [AuthorsRreply. Lancet, The, 2010, 376, 1737-1738] Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized controlled trials. Annals of Medicine, 2010, 42, 371-86	1.5	175
	Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized		175 6
40	Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized controlled trials. <i>Annals of Medicine</i> , 2010 , 42, 371-86 Stroke prevention in atrial fibrillation: putting the guidelines into practice. <i>Drugs and Aging</i> , 2010 ,	1.5	
40	Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized controlled trials. <i>Annals of Medicine</i> , 2010 , 42, 371-86 Stroke prevention in atrial fibrillation: putting the guidelines into practice. <i>Drugs and Aging</i> , 2010 , 27, 859-70 Diagnosis and management of adults with chronic heart failure: summary of updated NICE	1.5 4·7	6
40 39 38	Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized controlled trials. <i>Annals of Medicine</i> , 2010 , 42, 371-86 Stroke prevention in atrial fibrillation: putting the guidelines into practice. <i>Drugs and Aging</i> , 2010 , 27, 859-70 Diagnosis and management of adults with chronic heart failure: summary of updated NICE guidance. <i>BMJ</i> , <i>The</i> , 2010 , 341, c4130 Protocol for Past BP: a randomised controlled trial of different blood pressure targets for people with a history of stroke of transient ischaemic attack (TIA) in primary care. <i>BMC Cardiovascular</i>	1.5 4.7 5.9	27
40 39 38 37	Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized controlled trials. <i>Annals of Medicine</i> , 2010 , 42, 371-86 Stroke prevention in atrial fibrillation: putting the guidelines into practice. <i>Drugs and Aging</i> , 2010 , 27, 859-70 Diagnosis and management of adults with chronic heart failure: summary of updated NICE guidance. <i>BMJ</i> , <i>The</i> , 2010 , 341, c4130 Protocol for Past BP: a randomised controlled trial of different blood pressure targets for people with a history of stroke of transient ischaemic attack (TIA) in primary care. <i>BMC Cardiovascular Disorders</i> , 2010 , 10, 37 Effect of age on stroke prevention therapy in patients with atrial fibrillation: the atrial fibrillation	1.5 4·7 5·9	6 27 11
40 39 38 37 36	Does self-monitoring reduce blood pressure? Meta-analysis with meta-regression of randomized controlled trials. <i>Annals of Medicine</i> , 2010 , 42, 371-86 Stroke prevention in atrial fibrillation: putting the guidelines into practice. <i>Drugs and Aging</i> , 2010 , 27, 859-70 Diagnosis and management of adults with chronic heart failure: summary of updated NICE guidance. <i>BMJ</i> , <i>The</i> , 2010 , 341, c4130 Protocol for Past BP: a randomised controlled trial of different blood pressure targets for people with a history of stroke of transient ischaemic attack (TIA) in primary care. <i>BMC Cardiovascular Disorders</i> , 2010 , 10, 37 Effect of age on stroke prevention therapy in patients with atrial fibrillation: the atrial fibrillation investigators. <i>Stroke</i> , 2009 , 40, 1410-6	1.5 4·7 5·9 2.3 6.7	6 27 11 259

32	Commentary: point. Blood pressure-lowering treatment should be offered to all patients regardless of blood pressure. <i>Journal of Clinical Hypertension</i> , 2009 , 11, 698-701	2.3	
31	A before and after study of the impact of Specialist Workers for Older People. <i>Journal of Evaluation in Clinical Practice</i> , 2009 , 15, 335-40	2.5	4
30	Management of blood pressure in primary care. <i>BMJ, The</i> , 2009 , 338, b940	5.9	12
29	A survey of tuberculosis clinic provision in England and Wales. <i>Public Health</i> , 2008 , 122, 602-12	4	4
28	Management of type 2 diabetes: summary of updated NICE guidance. <i>BMJ, The</i> , 2008 , 336, 1306-8	5.9	79
27	Short-term effects on mobility, activities of daily living and health-related quality of life of a Conductive Education programme for adults with multiple sclerosis, Parkinsonß disease and stroke. <i>Clinical Rehabilitation</i> , 2008 , 22, 329-37	3.3	12
26	Pro: PWarfarin should be the drug of choice for thromboprophylaxis in elderly patients with atrial fibrillation Why warfarin should really be the drug of choice for stroke prevention in elderly patients with atrial fibrillation. <i>Thrombosis and Haemostasis</i> , 2008 , 100, 14-5	7	17
25	Blood pressure self monitoring: questions and answers from a national conference. <i>BMJ, The</i> , 2008 , 337, a2732	5.9	25
24	Accuracy of diagnosing atrial fibrillation on electrocardiogram by primary care practitioners and interpretative diagnostic software: analysis of data from screening for atrial fibrillation in the elderly (SAFE) trial. <i>BMJ</i> , <i>The</i> , 2007 , 335, 380	5.9	97
23	An analysis of factors that predict patient consent to take part in a randomized controlled trial. <i>Family Practice</i> , 2007 , 24, 388-94	1.9	12
22	A cross-sectional study of hypertension in an elderly population (75 years and over) with atrial fibrillation: secondary analysis of data from the Birmingham Atrial Fibrillation in the Aged (BAFTA) randomised controlled trial. <i>International Journal of Cardiology</i> , 2007 , 117, 152-6	3.2	10
21	Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial. <i>Lancet, The</i> , 2007 , 370, 493-503	40	1175
20	Warfarin versus aspirin for stroke prevention (BAFTA) [Authors Reply. Lancet, The, 2007, 370, 1606-160]	7 40	2
19	Process of care and mortality of stroke patients with and without a do not resuscitate order in the West Midlands, UK. <i>International Journal for Quality in Health Care</i> , 2006 , 18, 102-6	1.9	17
18	Clinicians didnR reliably distinguish between different causes of cardiac death using case histories. Journal of Clinical Epidemiology, 2006, 59, 862-7	5.7	34
17	Applicability to primary care of national clinical guidelines on blood pressure lowering for people with stroke: cross sectional study. <i>BMJ, The</i> , 2006 , 332, 635-7	5.9	36
16	Does it matter whether patients take their antihypertensive medication as prescribed? The complex relationship between adherence and blood pressure control. <i>Journal of Human Hypertension</i> , 2006 , 20, 551-3	2.6	23
15	Comparing processes of stroke care in high- and low-mortality hospitals in the West Midlands, UK. International Journal for Quality in Health Care, 2005, 17, 31-6	1.9	17

LIST OF PUBLICATIONS

14	A randomised controlled trial and cost effectiveness study of systematic screening (targeted and total population screening) versus routine practice for the detection of atrial fibrillation in the over 65s: (SAFE) [ISRCTN19633732]. <i>BMC Cardiovascular Disorders</i> , 2004 , 4, 12	2.3	12
13	Does changing from mercury to electronic blood pressure measurement influence recorded blood pressure? An observational study. <i>British Journal of General Practice</i> , 2003 , 53, 953-6	1.6	27
12	Inter-rater reliability of the Frenchay activities index in patients with stroke and their careers. <i>Clinical Rehabilitation</i> , 2000 , 14, 433-40	3.3	63
11	Family support for stroke: a randomised controlled trial. <i>Lancet, The</i> , 2000 , 356, 808-13	40	187
10	Uptake of research findings into clinical practice: a controlled study of the impact of a brief external intervention on the use of corticosteroids in preterm delivery. <i>Journal of Evaluation in Clinical Practice</i> , 1999 , 5, 73-9	2.5	5
9	Oral contraception and other factors in relation to back disorders in women: findings in a large cohort study. <i>Contraception</i> , 1999 , 60, 331-5	2.5	9
8	Risk of myocardial infarction, angina and stroke in users of oral contraceptives: an updated analysis of a cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1998 , 105, 890-6	3.7	48
7	Oral contraception and other factors in relation to hospital referral for fracture. Findings in a large cohort study. <i>Contraception</i> , 1998 , 57, 231-5	2.5	62
6	The impact of an information pack on patients with stroke and their carers: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 1998 , 12, 465-76	3.3	95
5	Comparison of postal version of the Frenchay Activities Index with interviewer-administered version for use in people with stroke. <i>Clinical Rehabilitation</i> , 1997 , 11, 131-8	3.3	32
4	Using the evidence: putting the research into practice. British Journal of Midwifery, 1997, 5, 396-399	0.3	1
3	Combined oral contraceptives and liver disease. <i>Contraception</i> , 1997 , 55, 145-51	2.5	37
2	Epidemiology of genital prolapse: observations from the Oxford Family Planning Association Study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1997 , 104, 579-85	3.7	514
1	Assessing quality of care: what are the implications of the potential lack of sensitivity of outcome measures to differences in quality?. <i>Journal of Evaluation in Clinical Practice</i> , 1996 , 2, 243-8	2.5	10