

Jing Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6957248/publications.pdf>

Version: 2024-02-01

12
papers

125
citations

1307594

7
h-index

1281871

11
g-index

12
all docs

12
docs citations

12
times ranked

181
citing authors

#	ARTICLE	IF	CITATIONS
1	Citrus alkaline extracts prevent fibroblast senescence to ameliorate pulmonary fibrosis via activation of COX-2. <i>Biomedicine and Pharmacotherapy</i> , 2019, 112, 108669.	5.6	29
2	Fluorescent Probes for Subcellular Localization during Osteoclast Formation. <i>ACS Medicinal Chemistry Letters</i> , 2014, 5, 911-914.	2.8	15
3	A Bioreductive Prodrug of Cucurbitacin B Significantly Inhibits Tumor Growth in the 4T1 Xenograft Mice Model. <i>ACS Medicinal Chemistry Letters</i> , 2019, 10, 1400-1406.	2.8	13
4	An oleanolic acid derivative reduces denervation-induced muscle atrophy via activation of CNTF-mediated JAK2/STAT3 signaling pathway. <i>European Journal of Pharmacology</i> , 2019, 861, 172612.	3.5	12
5	A novel oleanolic acid derivative HA-19 ameliorates muscle atrophy via promoting protein synthesis and preventing protein degradation. <i>Toxicology and Applied Pharmacology</i> , 2019, 378, 114625.	2.8	12
6	Synthesis and evaluation of tetrahydroquinolin-2(1H)-one derivatives as novel anti-pancreatic cancer agents via targeting autophagy. <i>European Journal of Medicinal Chemistry</i> , 2019, 170, 28-44.	5.5	11
7	Bone Marrow Derived Mesenchymal Stem Cells Involve in the Lymphangiogenesis of Lung Cancer and Jinfukang Inhibits the Involvement In Vivo. <i>Journal of Cancer</i> , 2017, 8, 1786-1794.	2.5	9
8	Modified Quinoxaline- α -Fused Oleanolic Acid Derivatives as Inhibitors of Osteoclastogenesis and Potential Agent in Anti- α -Osteoporosis. <i>ChemistrySelect</i> , 2020, 5, 1526-1533.	1.5	7
9	Oleanolic acid derivative HA-20 inhibits adipogenesis in a manner involving PPAR γ -FABP4/aP2 pathway. <i>Journal of Molecular Endocrinology</i> , 2021, 66, 245-258.	2.5	6
10	Novel heterocyclic ring-fused oleanolic acid derivatives as osteoclast inhibitors for osteoporosis. <i>MedChemComm</i> , 2016, 7, 371-377.	3.4	5
11	Dual roles of QOA-8a in antiosteoporosis: a combination of bone anabolic and anti-resorptive effects. <i>Acta Pharmacologica Sinica</i> , 2018, 39, 230-242.	6.1	5
12	A Unique Skeletal Rearrangement of a Bicyclo[3.3.1]nonanetrione to a Tetrahydroquinolin-2(1H)-one System. <i>Synlett</i> , 2018, 29, 1711-1716.	1.8	1