


List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6957536/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Lipotoxicity-induced mtDNA release promotes diabetic cardiomyopathy by activating the cGAS-STING pathway in obesity-related diabetes. Cell Biology and Toxicology, 2023, 39, 277-299.	5.3	46
2	Comparison of bovine serum albumin glycation by ribose and fructose in vitro and in vivo. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2022, 1868, 166283.	3.8	18
3	Maresin 1 Alleviates Diabetic Kidney Disease via LGR6-Mediated cAMP-SOD2-ROS Pathway. Oxidative Medicine and Cellular Longevity, 2022, 2022, 1-15.	4.0	4
4	Function and Mechanism of Novel Histone Posttranslational Modifications in Health and Disease. BioMed Research International, 2021, 2021, 1-13.	1.9	21
5	Innate Immunity in Diabetic Wound Healing: Focus on the Mastermind Hidden in Chronic Inflammatory. Frontiers in Pharmacology, 2021, 12, 653940.	3.5	48
6	PCB118 Induces Inflammation of Islet Beta Cells via Activating ROS-NLRP3 Inflammasome Signaling. BioMed Research International, 2021, 2021, 1-8.	1.9	5
7	Butyrate ameliorates alcoholic fatty liver disease via reducing endotoxemia and inhibiting liver gasdermin D-mediated pyroptosis. Annals of Translational Medicine, 2021, 9, 873-873.	1.7	22
8	G-quadruplex DNA: a novel target for drug design. Cellular and Molecular Life Sciences, 2021, 78, 6557-6583.	5.4	57
9	Low serum Maresin-1 levels are associated with non-alcoholic fatty liver disease: a cross-sectional study. Lipids in Health and Disease, 2021, 20, 96.	3.0	9
10	Serum albumin was negatively associated with diabetic peripheral neuropathy inÂChineseÂpopulation: a cross-sectional study. Diabetology and Metabolic Syndrome, 2021, 13, 100.	2.7	8
11	Decreased Serum Maresin 1 Concentration Is Associated With Postmenopausal Osteoporosis: A Cross-Sectional Study. Frontiers in Medicine, 2021, 8, 759825.	2.6	4
12	The efficacy and safety of combinations of SGLT2 inhibitors and GLP-1 receptor agonists in the treatment of type 2 diabetes or obese adults: a systematic review and meta-analysis. Endocrine, 2020, 67, 294-304.	2.3	23
13	Skipping breakfast is associated with overweight and obesity: A systematic review and meta-analysis. Obesity Research and Clinical Practice, 2020, 14, 1-8.	1.8	144
14	ROS-induced NLRP3 inflammasome priming and activation mediate PCB 118- induced pyroptosis in endothelial cells. Ecotoxicology and Environmental Safety, 2020, 189, 109937.	6.0	69
15	Short-Chain Fatty Acids Ameliorate Diabetic Nephropathy via GPR43-Mediated Inhibition of Oxidative Stress and NF- <i>ΰ</i> B Signaling. Oxidative Medicine and Cellular Longevity, 2020, 2020, 1-21.	4.0	102
16	Association between Circulating B-Type Natriuretic Peptide and Diabetic Peripheral Neuropathy: A Cross-Sectional Study of a Chinese Type 2 Diabetic Population. Journal of Diabetes Research, 2020, 2020, 1-10.	2.3	6
17	Association of Circulating Omentin-1 with Osteoporosis in a Chinese Type 2 Diabetic Population. Mediators of Inflammation, 2020, 2020, 1-16.	3.0	6
18	A Toolbox for Site-Specific Labeling of RecQ Helicase With a Single Fluorophore Used in the Single-Molecule Assay. Frontiers in Molecular Biosciences, 2020, 7, 586450.	3.5	3

Yong Xu

#	Article	IF	CITATIONS
19	Decreased Plasma Maresin 1 Concentration Is Associated with Diabetic Foot Ulcer. Mediators of Inflammation, 2020, 2020, 1-7.	3.0	16
20	Sodium butyrate alleviates high-glucose-induced renal glomerular endothelial cells damage via inhibiting pyroptosis. International Immunopharmacology, 2019, 75, 105832.	3.8	64
21	RIPK2-Mediated Autophagy and Negatively Regulated ROS-NLRP3 Inflammasome Signaling in GMCs Stimulated with High Glucose. Mediators of Inflammation, 2019, 2019, 1-13.	3.0	19
22	FBW7 Regulates the Autophagy Signal in Mesangial Cells Induced by High Glucose. BioMed Research International, 2019, 2019, 1-9.	1.9	12
23	Sodium Butyrate Improves Liver Glycogen Metabolism in Type 2 Diabetes Mellitus. Journal of Agricultural and Food Chemistry, 2019, 67, 7694-7705.	5.2	70
24	Physiological serum total bilirubin concentrations were inversely associated with diabetic peripheral neuropathy in Chinese patients with type 2 diabetes: a cross-sectional study. Diabetology and Metabolic Syndrome, 2019, 11, 100.	2.7	16
25	Decreased plasma neuregulin 4 levels are associated with peripheral neuropathy in Chinese patients with newly diagnosed type 2 diabetes: A cross-sectional study. Cytokine, 2019, 113, 356-364.	3.2	27
26	SGLT2 inhibitors and risk of stroke in patients with type 2 diabetes: A systematic review and metaâ€analysis. Diabetes, Obesity and Metabolism, 2018, 20, 1977-1982.	4.4	53
27	Comprehensive analysis of lysine crotonylation in proteome of maintenance hemodialysis patients. Medicine (United States), 2018, 97, e12035.	1.0	24
28	Sodium butyrate supplementation ameliorates diabetic inflammation in db/db mice. Journal of Endocrinology, 2018, 238, 231-244.	2.6	107
29	Maresins: Specialized Proresolving Lipid Mediators and Their Potential Role in Inflammatory-Related Diseases. Mediators of Inflammation, 2018, 2018, 1-8.	3.0	61
30	Plasma Neuregulin 4 Levels Are Associated with Metabolic Syndrome in Patients Newly Diagnosed with Type 2 Diabetes Mellitus. Disease Markers, 2018, 2018, 1-11.	1.3	36
31	Sweet Taste Receptors Mediated ROS-NLRP3 Inflammasome Signaling Activation: Implications for Diabetic Nephropathy. Journal of Diabetes Research, 2018, 2018, 1-15.	2.3	27
32	Association of serum uric acid with bone mineral density and clinical fractures in Chinese type 2 diabetes mellitus patients: A cross-sectional study. Clinica Chimica Acta, 2018, 486, 76-85.	1.1	22
33	Short-Chain Fatty Acids Inhibit Oxidative Stress and Inflammation in Mesangial Cells Induced by High Glucose and Lipopolysaccharide. Experimental and Clinical Endocrinology and Diabetes, 2017, 125, 98-105.	1.2	122
34	DsbA-L prevents obesity-induced inflammation and insulin resistance by suppressing the mtDNA release-activated cGAS-cGAMP-STING pathway. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, 12196-12201.	7.1	185
35	d -Ribose as a Contributor to Glycated Haemoglobin. EBioMedicine, 2017, 25, 143-153.	6.1	36
36	The role of short-chain fatty acids in kidney injury induced by gut-derived inflammatory response. Metabolism: Clinical and Experimental, 2017, 68, 20-30.	3.4	81

Үолс Хи

#	Article	IF	CITATIONS
37	Maresin 1 Mitigates High Glucose-Induced Mouse Glomerular Mesangial Cell Injury by Inhibiting Inflammation and Fibrosis. Mediators of Inflammation, 2017, 2017, 1-11.	3.0	41
38	SUMO E3 Ligase PIASy Mediates High Clucose-Induced Activation of NF- <i>κ</i> B Inflammatory Signaling in Rat Mesangial Cells. Mediators of Inflammation, 2017, 2017, 1-9.	3.0	9
39	CYLD Deubiquitinase Negatively Regulates High Glucose-Induced NF- <i>β</i> B Inflammatory Signaling in Mesangial Cells. BioMed Research International, 2017, 2017, 1-9.	1.9	7
40	High Glucose and Lipopolysaccharide Prime NLRP3 Inflammasome via ROS/TXNIP Pathway in Mesangial Cells. Journal of Diabetes Research, 2016, 2016, 1-11.	2.3	89
41	High Glucose Induces Sumoylation of Smad4 via SUMO2/3 in Mesangial Cells. BioMed Research International, 2014, 2014, 1-10.	1.9	15
42	The Role of Ubiquitination and Sumoylation in Diabetic Nephropathy. BioMed Research International, 2014, 2014, 1-11.	1.9	51
43	The Proteasome Inhibitor, MG132, Attenuates Diabetic Nephropathy by Inhibiting SnoN Degradation <i>In Vivo</i> and <i>In Vitro</i> . BioMed Research International, 2014, 2014, 1-11.	1.9	24
44	MG132 Ameliorates Kidney Lesions by Inhibiting the Degradation of Smad7 in Streptozotocin-Induced Diabetic Nephropathy. Journal of Diabetes Research, 2014, 2014, 1-8.	2.3	19
45	High glucose induces activation of NF-κB inflammatory signaling through IκBα sumoylation in rat mesangial cells. Biochemical and Biophysical Research Communications, 2013, 438, 568-574.	2.1	44
46	Notch Signaling Molecules Activate TGF- <i>β</i> in Rat Mesangial Cells under High Glucose Conditions. Journal of Diabetes Research, 2013, 2013, 1-8.	2.3	21
47	Impact of High Glucose and Proteasome Inhibitor MG132 on Histone H2A and H2B Ubiquitination in Rat Glomerular Mesangial Cells. Journal of Diabetes Research, 2013, 2013, 1-10.	2.3	30