

Yasuyuki Arakane

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6959213/publications.pdf>

Version: 2024-02-01

64
papers

5,587
citations

101384

36
h-index

149479

56
g-index

64
all docs

64
docs citations

64
times ranked

4508
citing authors

#	ARTICLE	IF	CITATIONS
1	Unveiling characteristic proteins for the structural development of beetle elytra. <i>Acta Biomaterialia</i> , 2022, 140, 467-480.	4.1	6
2	Chitin in insect cuticle. <i>Advances in Insect Physiology</i> , 2022, , 1-110.	1.1	5
3	Superoxide dismutase 6 is required during metamorphosis for the development of properly movable legs in <i>Tribolium castaneum</i> . <i>Scientific Reports</i> , 2022, 12, 6900.	1.6	0
4	Ultrastructural analysis of beetle larva cuticles during infection with the entomopathogenic fungus, <i>Beauveria bassiana</i> . <i>Pest Management Science</i> , 2022, 78, 3356-3364.	1.7	9
5	AA15 lytic polysaccharide monooxygenase is required for efficient chitinous cuticle turnover during insect molting. <i>Communications Biology</i> , 2022, 5, .	2.0	10
6	Chitin deacetylases are necessary for insect femur muscle attachment and mobility. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, .	3.3	2
7	Yellow-y Functions in Egg Melanization and Chorion Morphology of the Asian Tiger Mosquito, <i>Aedes albopictus</i> . <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 769788.	1.8	10
8	Gene functions in adult cuticle pigmentation of the yellow mealworm, <i>Tenebrio molitor</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2020, 117, 103291.	1.2	37
9	Yellow-g and Yellow-g2 proteins are required for egg desiccation resistance and temporal pigmentation in the Asian tiger mosquito, <i>Aedes albopictus</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2020, 122, 103386.	1.2	46
10	Insect Cuticular Chitin Contributes to Form and Function. <i>Current Pharmaceutical Design</i> , 2020, 26, 3530-3545.	0.9	43
11	Chitin Organizing and Modifying Enzymes and Proteins Involved In Remodeling of the Insect Cuticle. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1142, 83-114.	0.8	37
12	Future questions in insect chitin biology: A microreview. <i>Archives of Insect Biochemistry and Physiology</i> , 2018, 98, e21454.	0.6	14
13	Group I chitin deacetylases are essential for higher order organization of chitin fibers in beetle cuticle. <i>Journal of Biological Chemistry</i> , 2018, 293, 6985-6995.	1.6	34
14	A chitinase with two catalytic domains is required for organization of the cuticular extracellular matrix of a beetle. <i>PLoS Genetics</i> , 2018, 14, e1007307.	1.5	46
15	Development and ultrastructure of the rigid dorsal and flexible ventral cuticles of the elytron of the red flour beetle, <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2017, 91, 21-33.	1.2	36
16	Cuticle formation and pigmentation in beetles. <i>Current Opinion in Insect Science</i> , 2016, 17, 1-9.	2.2	125
17	Chitin Metabolic Pathways in Insects and Their Regulation. , 2016, , 31-65.		12
18	Tyrosine Metabolism for Insect Cuticle Pigmentation and Sclerotization. , 2016, , 165-220.		20

#	ARTICLE	IF	CITATIONS
19	Arylalkylamine N-acetyltransferase 1 gene (TcAANAT1) is required for cuticle morphology and pigmentation of the adult red flour beetle, <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2016, 79, 119-129.	1.2	39
20	Cuticular protein with a low complexity sequence becomes cross-linked during insect cuticle sclerotization and is required for the adult molt. <i>Scientific Reports</i> , 2015, 5, 10484.	1.6	67
21	<i>Tribolium castaneum</i> RR-1 Cuticular Protein TcCPR4 Is Required for Formation of Pore Canals in Rigid Cuticle. <i>PLoS Genetics</i> , 2015, 11, e1004963.	1.5	69
22	Loss of function of the yellow-e gene causes dehydration-induced mortality of adult <i>Tribolium castaneum</i> . <i>Developmental Biology</i> , 2015, 399, 315-324.	0.9	53
23	Knickkopf and retroactive proteins are required for formation of laminar serosal procuticle during embryonic development of <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2015, 60, 1-6.	1.2	22
24	S2-2 Development, Ultrastructure and Morphology of Cuticle of a Beetle. <i>Bulletin of Applied Glycoscience</i> , 2015, 5, B32.	0.0	0
25	A Multicopper Oxidase-Related Protein Is Essential for Insect Viability, Longevity and Ovary Development. <i>PLoS ONE</i> , 2014, 9, e111344.	1.1	14
26	Functional Specialization Among Members Of Knickkopf Family Of Proteins In Insect Cuticle Organization. <i>PLoS Genetics</i> , 2014, 10, e1004537.	1.5	19
27	A Major Facilitator Superfamily protein encoded by TcMuck gene is not required for cuticle pigmentation, growth and development in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2014, 49, 43-48.	1.2	2
28	Two major cuticular proteins are required for assembly of horizontal laminae and vertical pore canals in rigid cuticle of <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2014, 53, 22-29.	1.2	76
29	Retroactive Maintains Cuticle Integrity by Promoting the Trafficking of Knickkopf into the Procuticle of <i>Tribolium castaneum</i> . <i>PLoS Genetics</i> , 2013, 9, e1003268.	1.5	34
30	Formation of Rigid, Non-Flight Forewings (Elytra) of a Beetle Requires Two Major Cuticular Proteins. <i>PLoS Genetics</i> , 2012, 8, e1002682.	1.5	68
31	Chitin-Related Enzymes in Agro-Biosciences. <i>Current Drug Targets</i> , 2012, 13, 442-470.	1.0	43
32	Chitin Metabolism in Insects. , 2012, , 193-235.		99
33	RNAi-based functional genomics in <i>Tribolium castaneum</i> and possible application for controlling insect pests. <i>Entomological Research</i> , 2012, 42, 1-10.	0.6	19
34	Mechanical Properties of the Beetle Elytron, a Biological Composite Material. <i>Biomacromolecules</i> , 2011, 12, 321-335.	2.6	68
35	Knickkopf protein protects and organizes chitin in the newly synthesized insect exoskeleton. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 17028-17033.	3.3	106
36	Both UDP N-acetylglucosamine pyrophosphorylases of <i>Tribolium castaneum</i> are critical for molting, survival and fecundity. <i>Insect Biochemistry and Molecular Biology</i> , 2011, 41, 42-50.	1.2	69

#	ARTICLE	IF	CITATIONS
37	Cuticle tanning in <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 293-293.	0.6	1
38	Expression Profiles and Functional Analysis of Genes Encoding Chitin Deacetylases, Extracellular Matrix-Modifying Proteins in <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 294-294.	0.6	1
39	Functional Analysis of Genes of Chitin Metabolism in <i>Tribolium castaneum</i> by RNA interference. <i>Entomological Research</i> , 2011, 41, 295-295.	0.6	0
40	Two Major Structural Proteins Are Required for Rigid Adult Cuticle Formation in the Red Flour Beetle, <i>Tribolium castaneum</i> . <i>Entomological Research</i> , 2011, 41, 297-297.	0.6	0
41	Comparative Genomic Analysis of Chitinase and Chitinase-Like Genes in the African Malaria Mosquito (<i>Anopheles gambiae</i>). <i>PLoS ONE</i> , 2011, 6, e19899.	1.1	77
42	Insect chitinase and chitinase-like proteins. <i>Cellular and Molecular Life Sciences</i> , 2010, 67, 201-216.	2.4	278
43	Mechanical properties of elytra from <i>Tribolium castaneum</i> wild-type and body color mutant strains. <i>Journal of Insect Physiology</i> , 2010, 56, 1901-1906.	0.9	29
44	Chymotrypsin-like peptidases from <i>Tribolium castaneum</i> : A role in molting revealed by RNA interference. <i>Insect Biochemistry and Molecular Biology</i> , 2010, 40, 274-283.	1.2	49
45	Tyrosine hydroxylase is required for cuticle sclerotization and pigmentation in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2010, 40, 267-273.	1.2	104
46	Genes encoding proteins with peritrophin A-type chitin-binding domains in <i>Tribolium castaneum</i> are grouped into three distinct families based on phylogeny, expression and function. <i>Insect Biochemistry and Molecular Biology</i> , 2010, 40, 214-227.	1.2	141
47	Identification, mRNA expression and functional analysis of several yellow family genes in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2010, 40, 259-266.	1.2	72
48	Characterization of Multicopper Oxidase Related Protein (MCORP) in Two Insect Species. <i>FASEB Journal</i> , 2010, 24, 854.6.	0.2	0
49	Molecular and Functional Analyses of Amino Acid Decarboxylases Involved in Cuticle Tanning in <i>Tribolium castaneum</i> . <i>Journal of Biological Chemistry</i> , 2009, 284, 16584-16594.	1.6	181
50	Repeated Co-options of Exoskeleton Formation during Wing-to-Elytron Evolution in Beetles. <i>Current Biology</i> , 2009, 19, 2057-2065.	1.8	122
51	Analysis of functions of the chitin deacetylase gene family in <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2009, 39, 355-365.	1.2	145
52	The genome of the model beetle and pest <i>Tribolium castaneum</i> . <i>Nature</i> , 2008, 452, 949-955.	18.7	1,255
53	Functional analysis of four neuropeptides, EH, ETH, CCAP and bursicon, and their receptors in adult ecdysis behavior of the red flour beetle, <i>Tribolium castaneum</i> . <i>Mechanisms of Development</i> , 2008, 125, 984-995.	1.7	168
54	Domain organization and phylogenetic analysis of the chitinase-like family of proteins in three species of insects. <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 452-466.	1.2	129

#	ARTICLE	IF	CITATIONS
55	Characterization of recombinant chitinase-like proteins of <i>Drosophila melanogaster</i> and <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 467-477.	1.2	68
56	Characterization and expression of the β -N-acetylhexosaminidase gene family of <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 478-489.	1.2	84
57	Domain organization and phylogenetic analysis of proteins from the chitin deacetylase gene family of <i>Tribolium castaneum</i> and three other species of insects. <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 440-451.	1.2	130
58	Chitin synthases are required for survival, fecundity and egg hatch in the red flour beetle, <i>Tribolium castaneum</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2008, 38, 959-962.	1.2	145
59	Functional specialization among insect chitinase family genes revealed by RNA interference. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 6650-6655.	3.3	221
60	Laccase 2 is the phenoloxidase gene required for beetle cuticle tanning. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 11337-11342.	3.3	342
61	Chitin synthase genes in <i>Manduca sexta</i> : characterization of a gut-specific transcript and differential tissue expression of alternately spliced mRNAs during development. <i>Insect Biochemistry and Molecular Biology</i> , 2005, 35, 529-540.	1.2	110
62	Sequences of cDNAs and expression of genes encoding chitin synthase and chitinase in the midgut of <i>Spodoptera frugiperda</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2005, 35, 1249-1259.	1.2	89
63	Characterization of two chitin synthase genes of the red flour beetle, <i>Tribolium castaneum</i> , and alternate exon usage in one of the genes during development. <i>Insect Biochemistry and Molecular Biology</i> , 2004, 34, 291-304.	1.2	167
64	Properties of catalytic, linker and chitin-binding domains of insect chitinase. <i>Insect Biochemistry and Molecular Biology</i> , 2003, 33, 631-648.	1.2	120