Susanne C Schneider

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6962606/publications.pdf

Version: 2024-02-01

43 papers

1,710 citations

20 h-index 289244 40 g-index

43 all docs

43 docs citations

43 times ranked 1892 citing authors

#	Article	IF	CITATIONS
1	Ecological correlates of riverine diatom and macroinvertebrate alpha and beta diversity across Arctic Fennoscandia. Freshwater Biology, 2022, 67, 49-63.	2.4	17
2	Biodiversity patterns of Arctic diatom assemblages in lakes and streams: Current reference conditions and historical context for biomonitoring. Freshwater Biology, 2022, 67, 116-140.	2.4	18
3	Relating environmental pressures to littoral biological water quality indicators in Western Balkan lakes: Can we fill the largest gaps?. Science of the Total Environment, 2022, 804, 150160.	8.0	3
4	Charophytes in warm springs on Svalbard (Spitsbergen): DNA barcoding identifies Chara aspera and Chara canescens with unusual morphological traits. Botany Letters, 2020, 167, 179-186.	1.4	4
5	Littoral eutrophication indicators are more closely related to nearshore land use than to water nutrient concentrations: A critical evaluation of stressor-response relationships. Science of the Total Environment, 2020, 748, 141193.	8.0	15
6	Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems. Nature Ecology and Evolution, 2020, 4, 1060-1068.	7.8	336
7	Nutrient retention by the littoral vegetation of a large lake: Can Lake Ohrid cope with current and future loading?. Limnology and Oceanography, 2020, 65, 2390-2402.	3.1	7
8	Genetic and morphological variation in Chara contraria and a taxon morphologically resembling Chara connivens. Botany Letters, 2020, 167, 187-200.	1.4	9
9	Do benthic algae provide important information over and above that provided by macrophytes and phytoplankton in lake status assessment? $\hat{a}\in$ Results from a case study in Norway. Limnologica, 2019, 76, 28-40.	1.5	8
10	Morphological and molecular features of a <i>Chara vulgaris</i> population from desert springs on the Sinai Peninsula (Springs of Moses, Egypt). Botany Letters, 2018, 165, 77-89.	1.4	3
11	Recovery of benthic algal assemblages from acidification: how long does it take, and is there a link to eutrophication?. Hydrobiologia, 2018, 805, 33-47.	2.0	8
12	Unravelling the effect of flow regime on macroinvertebrates and benthic algae in regulated versus unregulated streams. Ecohydrology, 2018, 11, e1996.	2.4	13
13	Effects of flow regime on benthic algae and macroinvertebrates - A comparison between regulated and unregulated rivers. Science of the Total Environment, 2017, 579, 1059-1072.	8.0	50
14	Effects of flow events and nutrient addition on stream periphyton and macroinvertebrates: an experimental study using flumes. Knowledge and Management of Aquatic Ecosystems, 2017, , 47.	1.1	6
15	Species differentiation in the genus <i>Chara</i> (Charophyceae): considerable phenotypic plasticity occurs within homogenous genetic groups. European Journal of Phycology, 2016, 51, 282-293.	2.0	32
16	The "Forgotten―Ecology Behind Ecological Status Evaluation: Re-Assessing the Roles of Aquatic Plants and Benthic Algae in Ecosystem Functioning. Progress in Botany Fortschritte Der Botanik, 2016, , 285-304.	0.3	6
17	Greener rivers in a changing climate?—Effects of climate and hydrological regime on benthic algal assemblages in pristine streams. Limnologica, 2015, 55, 21-32.	1.5	14
18	Customs, habits, and traditions: the role of nonscientific factors in the development of ecological assessment methods. Wiley Interdisciplinary Reviews: Water, 2015, 2, 159-165.	6.5	17

#	Article	IF	CITATIONS
19	<scp>DNA</scp> barcoding the genus <i>Chara</i> molecular evidence recovers fewer taxa than the classical morphological approach. Journal of Phycology, 2015, 51, 367-380.	2.3	25
20	The role of charophytes (Charales) in past and present environments: An overview. Aquatic Botany, 2015, 120, 2-6.	1.6	50
21	Light acclimation in submerged macrophytes: The roles of plant elongation, pigmentation and branch orientation differ among Chara species. Aquatic Botany, 2015, 120, 121-128.	1.6	32
22	Eutrophication impacts littoral biota in Lake Ohrid while water phosphorus concentrations are low. Limnologica, 2014, 44, 90-97.	1.5	39
23	Assessment of littoral eutrophication in Lake Ohrid by submerged macrophytes. Biologia (Poland), 2014, 69, 756-764.	1.5	8
24	Juncus bulbosus nuisance growth in oligotrophic freshwater ecosystems: Different triggers for the same phenomenon in rivers and lakes?. Aquatic Botany, 2013, 104, 15-24.	1.6	21
25	Interactions between pH and nutrients on benthic algae in streams and consequences for ecological status assessment and species richness patterns. Science of the Total Environment, 2013, 444, 73-84.	8.0	68
26	Nuisance growth of <i>Juncus bulbosus</i> : the roles of genetics and environmental drivers tested in a largeâ€scale survey. Freshwater Biology, 2013, 58, 114-127.	2.4	12
27	Bioconcentration and Intracellular Storage of Hexachlorobenzene in Charophytes and Their Potential Role in Monitoring and Remediation Actions. Environmental Science & Environ	10.0	12
28	Establishing expectations for pan-European diatom based ecological status assessments. Ecological Indicators, 2012, 20, 177-186.	6.3	55
29	Do macrophytes, diatoms and non-diatom benthic algae give redundant information? Results from a case study in Poland. Limnologica, 2012, 42, 204-211.	1.5	54
30	Impact of calcium and TOC on biological acidification assessment in Norwegian rivers. Science of the Total Environment, 2011, 409, 1164-1171.	8.0	13
31	The periphyton index of trophic status PIT: a new eutrophication metric based on non-diatomaceous benthic algae in Nordic rivers. Hydrobiologia, 2011, 665, 143-155.	2.0	78
32	Chara baltica Bruzelius 1824 and Chara intermedia A. Braun 1859â€"Distinct species or habitat specific modifications?. Aquatic Botany, 2010, 93, 195-201.	1.6	19
33	Physiological acclimation to light in Chara intermedia nodes. Aquatic Botany, 2009, 91, 151-156.	1.6	11
34	Macrophyte trophic indicator values from a European perspective. Limnologica, 2007, 37, 281-289.	1.5	64
35	Differentiation of Chara intermedia and C. baltica compared to C. hispida based on morphology and amplified fragment length polymorphism. Hydrobiologia, 2007, 586, 155-166.	2.0	20
36	Growth towards light as an adaptation to high light conditions in Chara branches. New Phytologist, 2006, 172, 83-91.	7.3	49

#	Article	lF	CITATIONS
37	Macrophyte-based assessment of lakes - a contribution to the implementation of the European Water Framework Directive in Germany. International Review of Hydrobiology, 2005, 90, 223-237.	0.9	76
38	Sediment and Water Nutrient Characteristics in Patches of Submerged Macrophytes in Running Waters. Hydrobiologia, 2004, 527, 195-207.	2.0	35
39	Ecological classification of macrophytes and phytobenthos for rivers in Germany according to the water framework directive. Limnologica, 2004, 34, 283-301.	1.5	133
40	Macrophytes and phytobenthos as indicators of ecological status in German lakes $\hat{a} \in a$ contribution to the implementation of the water framework directive. Limnologica, 2004, 34, 302-314.	1.5	119
41	The Trophic Index of Macrophytes (TIM)– a New Tool for Indicating the Trophic State of Running Waters. International Review of Hydrobiology, 2003, 88, 49-67.	0.9	130
42	The importance of submerged macrophytes as indicators for the nutrient concentration in a small stream (Rotbach, Bavaria). Limnologica, 2000, 30, 351-358.	1.5	11
43	Indicating the trophic state of running waters by submersed macrophytes and epilithic diatoms. Limnologica, 2000, 30, 1-8.	1.5	10