

GÃ¼lÅah Akdemir

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6970014/publications.pdf>

Version: 2024-02-01

14
papers

185
citations

1478505

6
h-index

1281871

11
g-index

14
all docs

14
docs citations

14
times ranked

662
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinical and radiological outcomes of 5-year drug-free remission-steered treatment in patients with early arthritis: IMPROVED study. <i>Annals of the Rheumatic Diseases</i> , 2018, 77, 111-118.	0.9	51
2	Incidence and risk factors for adalimumab and infliximab anti-drug antibodies in rheumatoid arthritis: A European retrospective multicohort analysis. <i>Seminars in Arthritis and Rheumatism</i> , 2019, 48, 967-975.	3.4	46
3	Two-year results of disease activity score (DAS)-remission-steered treatment strategies aiming at drug-free remission in early arthritis patients (the IMPROVED-study). <i>Arthritis Research and Therapy</i> , 2016, 18, 23.	3.5	37
4	Predictive factors of radiological progression after 2 years of remission-steered treatment in early arthritis patients: a post hoc analysis of the IMPROVED study. <i>RMD Open</i> , 2016, 2, e000172.	3.8	13
5	Effectiveness of four dynamic treatment strategies in patients with anticitrullinated protein antibody-negative rheumatoid arthritis: a randomised trial. <i>RMD Open</i> , 2016, 2, e000143.	3.8	12
6	Rheumatologists' adherence to a disease activity score steered treatment protocol in early arthritis patients is less if the target is remission. <i>Clinical Rheumatology</i> , 2017, 36, 317-326.	2.2	10
7	Comparison between low disease activity or DAS remission as treatment target in patients with early active rheumatoid arthritis. <i>RMD Open</i> , 2018, 4, e000649.	3.8	7
8	Further Treatment Intensification in Undifferentiated and Rheumatoid Arthritis Patients Already in Low Disease Activity has Limited Benefit towards Physical Functioning. <i>Arthritis Research and Therapy</i> , 2017, 19, 220.	3.5	4
9	Erosions in the foot at baseline are predictive of orthopaedic shoe use after 10 years of treat to target therapy in patients with recent onset rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2016, 35, 2101-2107.	2.2	2
10	SAT0071 Predictive Factors of Radiological Progression After Two Years of Remission Steered Treatment in Early Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 674.1-674.	0.9	1
11	Age affects joint space narrowing in patients with early active rheumatoid arthritis. <i>RMD Open</i> , 2016, 2, e000338.	3.8	1
12	FRI0097 Repair of joint damage in newly diagnosed rheumatoid arthritis patients occurs but does not relate to previous suppression of inflammation; an 8-years sub analysis in the best-cohort. , 2017, , .		1
13	SAT0091 Age-Specific Risk Factors for Joint Space Narrowing Progression in Early Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 681.2-682.	0.9	0
14	SAT0610 Magnetic Resonance Imaging Outcomes in Patients with Chronic or Recurrent Gonarthrosis Treated with Intra-Articular Infliximab or Corticosteroid Injections. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 882.3-883.	0.9	0