
GülÅÄh Akdemir

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6970014/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Incidence and risk factors for adalimumab and infliximab anti-drug antibodies in rheumatoid arthritis: A European retrospective multicohort analysis. Seminars in Arthritis and Rheumatism, 2019, 48, 967-975.	3.4	46
2	Clinical and radiological outcomes of 5-year drug-free remission-steered treatment in patients with early arthritis: IMPROVED study. Annals of the Rheumatic Diseases, 2018, 77, 111-118.	0.9	51
3	Comparison between low disease activity or DAS remission as treatment target in patients with early active rheumatoid arthritis. RMD Open, 2018, 4, e000649.	3.8	7
4	Rheumatologists' adherence to a disease activity score steered treatment protocol in early arthritis patients is less if the target is remission. Clinical Rheumatology, 2017, 36, 317-326.	2.2	10
5	FRI0097â€Repair of joint damage in newly diagnosed rheumatoid arthritis patients occurs but does not relate to previous suppression of inflammation; an 8-years sub analysis in the best-cohort. , 2017, , .		1
6	Further Treatment Intensification in Undifferentiated and Rheumatoid Arthritis Patients Already in Low Disease Activity has Limited Benefit towards Physical Functioning. Arthritis Research and Therapy, 2017, 19, 220.	3.5	4
7	Predictive factors of radiological progression after 2â€years of remission-steered treatment in early arthritis patients: a post hoc analysis of the IMPROVED study. RMD Open, 2016, 2, e000172.	3.8	13
8	Effectiveness of four dynamic treatment strategies in patients with anticitrullinated protein antibody-negative rheumatoid arthritis: a randomised trial. RMD Open, 2016, 2, e000143.	3.8	12
9	Age affects joint space narrowing in patients with early active rheumatoid arthritis. RMD Open, 2016, 2, e000338.	3.8	1
10	Two-year results of disease activity score (DAS)-remission-steered treatment strategies aiming at drug-free remission in early arthritis patients (the IMPROVED-study). Arthritis Research and Therapy, 2016, 18, 23.	3.5	37
11	Erosions in the foot at baseline are predictive of orthopaedic shoe use after 10Âyears of treat to target therapy in patients with recent onset rheumatoid arthritis. Clinical Rheumatology, 2016, 35, 2101-2107.	2.2	2
12	SAT0091â€Age-Specific Risk Factors for Joint Space Narrowing Progression in Early Rheumatoid Arthritis Patients. Annals of the Rheumatic Diseases, 2015, 74, 681.2-682.	0.9	0
13	SAT0071â€Predictive Factors of Radiological Progression After Two Years of Remission Steered Treatment in Early Arthritis Patients. Annals of the Rheumatic Diseases, 2015, 74, 674.1-674.	0.9	1
14	SAT0610â€Magnetic Resonance Imaging Outcomes in Patients with Chronic or Recurrent Gonarthritis Treated with Intra-Articular Infliximab or Corticosteroid Injections. Annals of the Rheumatic Diseases, 2015, 74, 882.3-883.	0.9	0