Moyra E Mortby

List of Publications by Citations

Source: https://exaly.com/author-pdf/6973826/moyra-e-mortby-publications-by-citations.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40 1,114 16 33 g-index

52 1,500 4 4.4 L-index

#	Paper	IF	Citations
40	Prevalence of Depression in Patients With Mild Cognitive Impairment: A Systematic Review and Meta-analysis. <i>JAMA Psychiatry</i> , 2017 , 74, 58-67	14.5	203
39	The Mild Behavioral Impairment Checklist[(MBI-C): A Rating Scale for[Neuropsychiatric Symptoms in[Pre-Dementia Populations. <i>Journal of Alzheimerts Disease</i> , 2017 , 56, 929-938	4.3	182
38	Apathy associated with neurocognitive disorders: Recent progress and future directions. <i>Alzheimert</i> s and Dementia, 2017 , 13, 84-100	1.2	127
37	Affective and emotional dysregulation as pre-dementia risk markers: exploring the mild behavioral impairment symptoms of depression, anxiety, irritability, and euphoria. <i>International Psychogeriatrics</i> , 2018 , 30, 185-196	3.4	86
36	Prevalence of mild behavioral impairment in mild cognitive impairment and subjective cognitive decline, and its association with caregiver burden. <i>International Psychogeriatrics</i> , 2018 , 30, 233-244	3.4	65
35	Prevalence estimates of mild behavioral impairment in a population-based sample of pre-dementia states and cognitively healthy older adults. <i>International Psychogeriatrics</i> , 2018 , 30, 221-232	3.4	61
34	Caregiver rating bias in mild cognitive impairment and mild Alzheimer's disease: impact of caregiver burden and depression on dyadic rating discrepancy across domains. <i>International Psychogeriatrics</i> , 2013 , 25, 1345-55	3.4	43
33	Apathy: a separate syndrome from depression in dementia? A critical review. <i>Aging Clinical and Experimental Research</i> , 2012 , 24, 305-16	4.8	40
32	Dementia clinical trial implications of mild behavioral impairment. <i>International Psychogeriatrics</i> , 2018 , 30, 171-175	3.4	39
31	High "normal" blood glucose is associated with decreased brain volume and cognitive performance in the 60s: the PATH through life study. <i>PLoS ONE</i> , 2013 , 8, e73697	3.7	37
30	Association of Alzheimer's genetic loci with mild behavioral impairment. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2018 , 177, 727-735	3.5	31
29	Neuropsychiatric Symptoms and Cognitive Impairment: Understanding the Importance of Co-Morbid Symptoms. <i>Journal of Alzheimerts Disease</i> , 2017 , 59, 141-153	4.3	24
28	Evaluation of a research diagnostic algorithm for DSM-5 neurocognitive disorders in a population-based cohort of older adults. <i>Alzheimerts Research and Therapy</i> , 2017 , 9, 15	9	21
27	Diagnostic criteria for apathy in neurocognitive disorders. Alzheimerts and Dementia, 2021,	1.2	18
26	Blood pressure, brain structure, and cognition: opposite associations in men and women. <i>American Journal of Hypertension</i> , 2015 , 28, 225-31	2.3	17
25	Using sulcal and gyral measures of brain structure to investigate benefits of an active lifestyle. <i>NeuroImage</i> , 2014 , 91, 353-9	7.9	16
24	Association of sex differences in dementia risk factors with sex differences in memory decline in a population-based cohort spanning 20-76lyears. <i>Scientific Reports</i> , 2021 , 11, 7710	4.9	15

(2020-2018)

23	Depressive symptoms as a barrier to engagement in physical activity in older adults with and without Alzheimer's disease. <i>PLoS ONE</i> , 2018 , 13, e0208581	3.7	12
22	Agitation and impulsivity in mid and late life as possible risk markers for incident dementia. <i>Alzheimerts and Dementia: Translational Research and Clinical Interventions</i> , 2020 , 6, e12016	6	9
21	Cognitive and Neuropsychiatric Screening Tests in Older Adults. <i>Mental Health and Illness Worldwide</i> , 2017 , 343-368	0.1	7
20	Differential associations between sensory loss and neuropsychiatric symptoms in adults with and without a neurocognitive disorder. <i>International Psychogeriatrics</i> , 2018 , 30, 261-272	3.4	7
19	Depressive symptoms in later life: differential impact of social support and motivational processes on depression in individuals with and without cognitive impairment. <i>European Journal of Ageing</i> , 2014 , 11, 321-332	3.6	5
18	Midlife motivational abilities predict apathy and depression in Alzheimer disease: the aging, demographics, and memory study. <i>Journal of Geriatric Psychiatry and Neurology</i> , 2011 , 24, 151-60	3.8	5
17	Cohort Profile Update: The PATH Through Life Project. <i>International Journal of Epidemiology</i> , 2021 , 50, 35-36	7.8	5
16	Apathy as a Treatment Target in Alzheimer's Disease: Implications for Clinical Trials. <i>American Journal of Geriatric Psychiatry</i> , 2021 ,	6.5	5
15	Relating education, brain structure, and cognition: the role of cardiovascular disease risk factors. <i>BioMed Research International</i> , 2014 , 2014, 271487	3	4
14	Sex differences in neuropsychiatric symptoms in Alzheimer's disease dementia: a meta-analysis <i>Alzheimerts Research and Therapy</i> , 2022 , 14, 48	9	4
13	Non-pharmacological interventions for neuropsychiatric symptoms of dementia in residential aged care settings: An umbrella review <i>International Journal of Nursing Studies</i> , 2022 , 128, 104187	5.8	2
12	[P1🛮40]: ALZHEIMER'S GENETIC RISK SCORE LINKED TO MILD BEHAVIORAL IMPAIRMENT 2017 , 13, P296-P296		1
11	Cognitive and Neuropsychiatric Screening Tests in Older Adults. <i>Mental Health and Illness Worldwide</i> , 2016 , 1-26	0.1	1
10	Mild behavioral impairment: challenges facing a quickly developing evidence base. <i>International Psychogeriatrics</i> , 2021 , 33, 209-212	3.4	O
9	Gender differences in cognitive development in cohorts of young, middle, and older adulthood over 12 years. <i>Developmental Psychology</i> , 2021 , 57, 1403-1410	3.7	О
8	Diastolic Blood Pressure Variability in Later Life May Be a Key Risk Marker for Cognitive Decline <i>Hypertension</i> , 2022 , HYPERTENSIONAHA12118799	8.5	O
7	Who displays dementia-related stigma and what does the general public know about dementia? Findings from a nationally representative survey <i>Aging and Mental Health</i> , 2022 , 1-9	3.5	0
6	The BPSDPLUS Program: Efficacy of a novel person-centred care approach to reduce neuropsychiatric symptoms in residents with dementia. <i>Alzheimerts and Dementia</i> , 2020 , 16, e038866	1.2	

5	Third-Party Impact of Dual Sensory Loss on Neuropsychiatric Symptom-Related Distress among Friends and Family. <i>Gerontology</i> , 2020 , 66, 351-361	5.5
4	Greater cognitive decline relative to normal ageing occurs at least 10 years prior to clinical dementia diagnosis. <i>BMJ Evidence-Based Medicine</i> , 2018 , 23, 79	2.7
3	Residential aged care staff perceptions and responses towards neuropsychiatric symptoms: a mixed methods analysis of electronic healthcare records <i>Aging and Mental Health</i> , 2022 , 1-8	3.5
2	303 - Residential aged care staffd perception of and response to behavioral and psychological symptoms of dementia de qualitative analysis of clinical care notes. <i>International Psychogeriatrics</i> , 2020 , 32, 61-61	3.4
1	Association Between Anxiety and Cognitive Decline Over 12 Years in a Population-Based Cohort. Journal of Alzheimerts Disease, 2021, 84, 409-418	4.3