

# Mads Svart

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6976441/publications.pdf>

Version: 2024-02-01

12  
papers

534  
citations

1040056

9  
h-index

1199594

12  
g-index

13  
all docs

13  
docs citations

13  
times ranked

855  
citing authors

#	ARTICLE	IF	CITATIONS
1	A randomized placebo-controlled clinical trial of nicotinamide riboside in obese men: safety, insulin-sensitivity, and lipid-mobilizing effects. <i>American Journal of Clinical Nutrition</i> , 2018, 108, 343-353.	4.7	195
2	Ketone Body Infusion With 3- <sup>11</sup> C-Hydroxybutyrate Reduces Myocardial Glucose Uptake and Increases Blood Flow in Humans: A Positron Emission Tomography Study. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	144
3	Regional cerebral effects of ketone body infusion with 3-hydroxybutyrate in humans: Reduced glucose uptake, unchanged oxygen consumption and increased blood flow by positron emission tomography. A randomized, controlled trial. <i>PLoS ONE</i> , 2018, 13, e0190556.	2.5	59
4	A macrophage-hepatocyte glucocorticoid receptor axis coordinates fasting ketogenesis. <i>Cell Metabolism</i> , 2022, 34, 473-486.e9.	16.2	34
5	Effects of ̢-hydroxybutyrate on cognition in patients with type 2 diabetes. <i>European Journal of Endocrinology</i> , 2020, 182, 233-242.	3.7	23
6	Oral D/L-3-Hydroxybutyrate Stimulates Cholecystokinin and Insulin Secretion and Slows Gastric Emptying in Healthy Males. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e3597-e3605.	3.6	18
7	Macrophage activation marker sCD163 correlates with accelerated lipolysis following LPS exposure: a human-randomised clinical trial. <i>Endocrine Connections</i> , 2018, 7, 107-114.	1.9	16
8	Combined Insulin Deficiency and Endotoxin Exposure Stimulate Lipid Mobilization and Alter Adipose Tissue Signaling in an Experimental Model of Ketoacidosis in Subjects With Type 1 Diabetes: A Randomized Controlled Crossover Trial. <i>Diabetes</i> , 2016, 65, 1380-1386.	0.6	13
9	Ketone Body Infusion Increases Circulating Erythropoietin and Bone Marrow Glucose Uptake. <i>Diabetes Care</i> , 2018, 41, e152-e154.	8.6	11
10	Oral 3- <sup>11</sup> C-hydroxybutyrate ingestion decreases endogenous glucose production, lipolysis, and hormone-sensitive lipase phosphorylation in adipose tissue in men: a human randomized, controlled, crossover trial. <i>Diabetic Medicine</i> , 2021, 38, e14385.	2.3	11
11	Soluble CD163 correlates with lipid metabolic adaptations in type 1 diabetes patients during ketoacidosis. <i>Journal of Diabetes Investigation</i> , 2019, 10, 67-72.	2.4	9
12	3-Hydroxybutyrate administration elevates plasma parathyroid hormone in a pilot human randomized, controlled, cross over trial. <i>Bone</i> , 2021, 153, 116166.	2.9	1