

Tadeusz Przewlocki

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6977226/tadeusz-przewlocki-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

76
papers

1,009
citations

18
h-index

28
g-index

79
ext. papers

1,145
ext. citations

2
avg, IF

3.61
L-index

#	Paper	IF	Citations
76	Clinical Significance of Carotid Intima-Media Complex and Carotid Plaque Assessment by Ultrasound for the Prediction of Adverse Cardiovascular Events in Primary and Secondary Care Patients. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	4
75	Importance of Increased Arterial Resistance in Risk Prediction in Patients with Cardiovascular Risk Factors and Degenerative Aortic Stenosis. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1
74	Expression of miR-1-3p, miR-16-5p and miR-122-5p as Possible Risk Factors of Secondary Cardiovascular Events. <i>Biomedicines</i> , 2021 , 9,	4.8	4
73	Long-term outcomes and determinants of stenosis recurrence after renal artery angioplasty in hypertensive patients with renovascular disease. <i>Postepy W Kardiologii Interwencyjnej</i> , 2020 , 16, 65-75	0.4	
72	Percutaneous closure of atrial septal defect: a consensus document of the joint group of experts from the Association of Cardiovascular Interventions and the Grown-Up Congenital Heart Disease Section of the Polish Cardiac Society. <i>Kardiologia Polska</i> , 2020 , 78, 1066-1083	0.9	
71	One-day, sequential carotid artery stenting followed by cardiac surgery in patients with severe carotid and cardiac disease. <i>Vascular Medicine</i> , 2019 , 24, 431-438	3.3	1
70	The role of serial carotid intima-media thickness assessment as a surrogate marker of atherosclerosis control in patients with recent myocardial infarction. <i>Postepy W Kardiologii Interwencyjnej</i> , 2019 , 15, 74-80	0.4	3
69	Comparison of drug-eluting and bare metal stents for extracranial vertebral artery stenting. <i>Postepy W Kardiologii Interwencyjnej</i> , 2019 , 15, 328-337	0.4	2
68	Transcatheter closure of atrial septal communication: Impact on quality of life in mid-term follow-up. <i>Advances in Clinical and Experimental Medicine</i> , 2019 , 28, 1079-1085	1.8	2
67	Prospective study on the prognostic value of repeated carotid intima-media thickness assessment in patients with coronary and extra coronary steno-occlusive arterial disease. <i>Polish Archives of Internal Medicine</i> , 2019 , 129, 12-21	1.9	6
66	Determinants of long-term outcome in patients after percutaneous stent-assisted intervention for renal artery steno-occlusive atherosclerotic disease. <i>Polish Archives of Internal Medicine</i> , 2019 , 129, 747-760	1.8	4
65	Interventional closure of patent foramen ovale in prevention of thromboembolic events. Consensus document of the Association of Cardiovascular Interventions and the Section of Grown-up Congenital Heart Disease of the Polish Cardiac Society. <i>Kardiologia Polska</i> , 2019 , 77, 1094-1105	0.9	2
64	Severe, recurrent in-stent carotid restenosis: endovascular approach, risk factors. Results from a prospective academic registry of 2637 consecutive carotid artery stenting procedures (TARGET-CAS). <i>Postepy W Kardiologii Interwencyjnej</i> , 2019 , 15, 465-471	0.4	8
63	Circulating miRNA levels differ with respect to carotid plaque characteristics and symptom occurrence in patients with carotid artery stenosis and provide information on future cardiovascular events. <i>Postepy W Kardiologii Interwencyjnej</i> , 2018 , 14, 75-84	0.4	11
62	Improvement of physical capacity in patients undergoing transcatheter closure of atrial septal defects. <i>Postepy W Kardiologii Interwencyjnej</i> , 2018 , 14, 90-94	0.4	2
61	Diagnostic and prognostic micro-RNAs in ischaemic stroke due to carotid artery stenosis and in acute coronary syndrome: a four-year prospective study. <i>Kardiologia Polska</i> , 2018 , 76, 362-369	0.9	22
60	Temporal changes in the pattern of invasive angiography use and its outcome in suspected coronary artery disease: implications for patient management and healthcare resources utilization. <i>Postepy W Kardiologii Interwencyjnej</i> , 2018 , 14, 247-257	0.4	2

59	Increased risk profile in the treatment of patients with symptomatic degenerative aortic valve stenosis over the last 10 years. <i>Postepy W Kardiologii Interwencyjnej</i> , 2018 , 14, 276-284	0.4	2
58	Magnetic resonance imaging and clinical outcome in patients with symptomatic carotid artery stenosis after carotid artery revascularization. <i>Postepy W Kardiologii Interwencyjnej</i> , 2017 , 13, 225-232	0.4	4
57	Simultaneous vertebral and subclavian artery stenting. <i>Postepy W Kardiologii Interwencyjnej</i> , 2017 , 13, 142-149	0.4	2
56	Determinants of long-term outcome in patients after percutaneous stent-assisted management of symptomatic subclavian or innominate artery stenosis or occlusion. <i>EuroIntervention</i> , 2017 , 13, 1355-1364 ^{3,1}	3.1	9
55	Resolution of ischemic symptoms after percutaneous angioplasty for a symptomatic subclavian artery stenosis. <i>Journal of Vascular Surgery</i> , 2016 , 64, 684-91	3.5	5
54	Decision-making microRNAs (miR-124, -133a/b, -34a and -134) in patients with occluded target vessel in acute coronary syndrome. <i>Kardiologia Polska</i> , 2016 , 74, 280-8	0.9	27
53	Triple anticoagulation therapy in patients with atrial fibrillation undergoing percutaneous coronary intervention - real life assessment. <i>Postepy W Kardiologii Interwencyjnej</i> , 2016 , 12, 303-313	0.4	5
52	Multiterritory Atherosclerosis and Carotid Intima-Media Thickness as Cardiovascular Risk Predictors After Percutaneous Angioplasty of Symptomatic Subclavian Artery Stenosis. <i>Journal of Ultrasound in Medicine</i> , 2016 , 35, 1977-84	2.9	7
51	Low prevalence of collateral cerebral circulation in the circle of Willis in patients with severe carotid artery stenosis and recent ischemic stroke. <i>Postepy W Kardiologii Interwencyjnej</i> , 2015 , 11, 312-7	0.4	6
50	Elevated level of plasma endothelin-1 in patients with atrial septal defect. <i>Cardiovascular Ultrasound</i> , 2014 , 12, 31	2.4	2
49	Transcranial Doppler ultrasonography should it be the first choice for persistent foramen ovale screening?. <i>Cardiovascular Ultrasound</i> , 2014 , 12, 16	2.4	35
48	Conduction Abnormality and Arrhythmia After Transcatheter Closure of Atrial Septal Defect. <i>Circulation Journal</i> , 2014 , 78, 2415-2421	2.9	16
47	The benefit of atrial septal defect closure in elderly patients. <i>Clinical Interventions in Aging</i> , 2014 , 9, 1101-7	1.7	27
46	Is it worth closing the atrial septal defect in patients with insignificant shunt?. <i>Postepy W Kardiologii Interwencyjnej</i> , 2014 , 10, 78-83	0.4	5
45	Bioresorbable vascular scaffolds in patients with acute coronary syndromes: the POLAR ACS study. <i>Polish Archives of Internal Medicine</i> , 2014 , 124, 669-77	1.9	18
44	Carotid artery stenting according to the "tailored CAS" algorithm performed in the very elderly patients: the thirty day outcome. <i>Catheterization and Cardiovascular Interventions</i> , 2013 , 82, 681-8	2.7	8
43	Multimarker approach in discriminating patients with symptomatic and asymptomatic atherosclerotic carotid artery stenosis. <i>Journal of Clinical Neurology (Korea)</i> , 2013 , 9, 165-75	1.7	28
42	Impact of renal artery stenting on cytokine levels, left ventricle mass and diastolic function. <i>Kardiologia Polska</i> , 2013 , 71, 121-8	0.9	3

41	Microparticles, not only markers but also a therapeutic target in the early stage of diabetic retinopathy and vascular aging. <i>Expert Opinion on Therapeutic Targets</i> , 2012 , 16, 677-88	6.4	18
40	Fibrinogen and carotid intima media thickness determine fibrin density in different atherosclerosis extents. <i>International Journal of Cardiology</i> , 2012 , 157, 411-3	3.2	6
39	Control of local haemostasis with the AngioSeal [®] vascular closure device in peripheral endovascular interventions via 6-9 F femoral artery access. <i>Postepy W Kardiologii Interwencyjnej</i> , 2012 , 1, 1-7	0.4	1
38	Temporary vertebral flow reversal by Mono Mo.Ma: a new proximal protection system in the treatment of recent vertebral artery occlusion. <i>Postepy W Kardiologii Interwencyjnej</i> , 2012 , 1, 66-69	0.4	
37	Zotarolimus-eluting stent for the treatment of recurrent, severe carotid artery in-stent stenosis in the TARGET-CAS population. <i>Journal of Endovascular Therapy</i> , 2012 , 19, 316-24	2.5	29
36	Safety of embolic protection device-assisted and unprotected intravascular ultrasound in evaluating carotid artery atherosclerotic lesions. <i>Medical Science Monitor</i> , 2012 , 18, MT7-18	3.2	17
35	Evolution of myocardial perfusion during primary angioplasty in spontaneously reperfused infarct-related artery: impact on long-term clinical outcomes and left ventricular function recovery. <i>International Journal of Cardiology</i> , 2011 , 147, 25-31	3.2	10
34	Effect of renal artery revascularization on left ventricular hypertrophy, diastolic function, blood pressure, and the one-year outcome. <i>Journal of Vascular Surgery</i> , 2011 , 53, 692-7	3.5	15
33	Carotid intima-media thickness, hs-CRP and TNF- α are independently associated with cardiovascular event risk in patients with atherosclerotic occlusive disease. <i>Atherosclerosis</i> , 2011 , 214, 185-90	3.1	55
32	Commentary: Carotid artery stenting drawbacks: microembolic ischemic cerebral lesions--do they matter?. <i>Journal of Endovascular Therapy</i> , 2011 , 18, 527-30	2.5	2
31	Conservative strategy in patients with non-ST-segment elevation acute coronary syndromes. <i>Postepy W Kardiologii Interwencyjnej</i> , 2010 , 4, 147-153	0.4	
30	Conservative strategy in patients with ST-segment elevation myocardial infarction. <i>Postepy W Kardiologii Interwencyjnej</i> , 2010 , 3, 97-103	0.4	
29	Predictors of cerebral reperfusion injury after carotid stenting: the role of transcranial color-coded Doppler ultrasonography. <i>Journal of Endovascular Therapy</i> , 2010 , 17, 556-63	2.5	18
28	The role of carotid intima-media thickness assessment in cardiovascular risk evaluation in patients with polyvascular atherosclerosis. <i>Atherosclerosis</i> , 2010 , 209, 125-30	3.1	18
27	Altered fibrin clot structure/function in patients with cryptogenic ischemic stroke. <i>Stroke</i> , 2009 , 40, 1499-501	6.7	99
26	Flow reversal for proximal neuroprotection during endovascular management of critical symptomatic carotid artery stenosis coexisting with ipsilateral external carotid artery occlusion. <i>Journal of Endovascular Therapy</i> , 2009 , 16, 744-51	2.5	10
25	Polyvascular extracoronary atherosclerotic disease in patients with coronary artery disease. <i>Kardiologia Polska</i> , 2009 , 67, 978-84	0.9	12
24	Supra-aortic extracranial artery atherosclerotic lesions in patients diagnosed for coronary artery disease: prevalence and predictors. <i>Kardiologia Polska</i> , 2009 , 67, 985-91	0.9	7

23	Carotid artery stenting with patient- and lesion-tailored selection of the neuroprotection system and stent type: early and 5-year results from a prospective academic registry of 535 consecutive procedures (TARGET-CAS). <i>Journal of Endovascular Therapy</i> , 2008 , 15, 249-62	2.5	62
22	Carotid intima-media thickness in pre- and postmenopausal women with suspected coronary artery disease. <i>Heart and Vessels</i> , 2008 , 23, 295-300	2.1	16
21	Novel paclitaxel-eluting, biodegradable polymer coated stent in the treatment of de novo coronary lesions: a prospective multicenter registry. <i>Catheterization and Cardiovascular Interventions</i> , 2008 , 71, 51-7	2.7	29
20	IVRT/IVRT index is a useful tool for detection of elevated left ventricular filling pressure in patients with preserved ejection fraction. <i>Echocardiography</i> , 2008 , 25, 473-81	1.5	8
19	Prevalence and prediction of renal artery stenosis in patients with coronary and supraaortic artery atherosclerotic disease. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 580-5	4.3	21
18	Doppler ultrasonography in suspected subclavian artery obstruction and in patient monitoring after subclavian stenting. <i>CardioVascular and Interventional Radiology</i> , 2007 , 30, 894-900	2.7	8
17	Diagnostic value of carotid intima-media thickness in indicating multi-level atherosclerosis. <i>Atherosclerosis</i> , 2007 , 193, 395-400	3.1	28
16	Subarachnoid hemorrhage after carotid artery stenting. <i>Journal of Vascular Surgery</i> , 2007 , 45, 1072-5	3.5	6
15	Determinants of immediate and long-term results of subclavian and innominate artery angioplasty. <i>Catheterization and Cardiovascular Interventions</i> , 2006 , 67, 519-26	2.7	60
14	Assessment of flow changes in the circle of Willis after stenting for severe internal carotid artery stenosis. <i>Journal of Endovascular Therapy</i> , 2006 , 13, 205-13	2.5	20
13	Prognostic value of carotid intima-media thickness in detection of coronary atherosclerosis in patients with calcified aortic valve stenosis. <i>Journal of Ultrasound in Medicine</i> , 2005 , 24, 461-7	2.9	12
12	Gender differences in carotid intima-media thickness in patients with suspected coronary artery disease. <i>American Journal of Cardiology</i> , 2005 , 96, 1217-22	3	32
11	Use of the Parodi Anti-Emboli System and transient subclavian steal for cerebral protection during emergent vertebral artery recanalization. <i>Journal of Endovascular Therapy</i> , 2004 , 11, 511-6	2.5	18
10	Detection of microvascular injury by evaluating epicardial blood flow in early reperfusion following primary angioplasty. <i>International Journal of Cardiology</i> , 2004 , 96, 389-96	3.2	13
9	Prognostic value of nitrate enhanced Tc99m MIBI SPECT study in detecting viable myocardium in patients with coronary artery disease. <i>International Journal of Cardiovascular Imaging</i> , 2003 , 19, 129-35		8
8	Facilitated percutaneous coronary intervention in patients with acute myocardial infarction transferred from remote hospitals. <i>American Journal of Cardiology</i> , 2003 , 91, 227-9	3	14
7	Assessment of myocardial perfusion in patients with coronary artery disease. Comparison of myocardial contrast echocardiography and 99mTc MIBI single photon emission computed tomography. <i>International Journal of Cardiology</i> , 2003 , 90, 49-55	3.2	24
6	Clinical and angiographic performance of a new-generation modular stent design for treatment of de novo coronary lesions. <i>Catheterization and Cardiovascular Interventions</i> , 2001 , 54, 276-82	2.7	2

5	Predictive potential of noninvasive methods, inclusive of exercise SPECT Tc99m MIBI imaging, in recognition of high-risk patients with left main coronary artery stenosis. <i>International Journal of Cardiovascular Imaging</i> , 2001 , 17, 347-52		5
4	Long-term outcome in patients with unstable angina treated by coronary balloon angioplasty. <i>International Journal of Cardiology</i> , 2001 , 77, 13-24	3.2	3
3	Balloon positioning difficulties during nonsurgical septal reduction therapy in a patient with hypertrophic obstructive cardiomyopathy. <i>Catheterization and Cardiovascular Interventions</i> , 2000 , 49, 314-7	2.7	2
2	Long-term outcome of coronary balloon angioplasty in diabetic patients. <i>International Journal of Cardiology</i> , 2000 , 76, 7-16	3.2	11
1	Thallium-201 myocardial perfusion imaging in patients before and after successful percutaneous transluminal coronary angioplasty. <i>International Journal of Cardiology</i> , 1996 , 53, 299-304	3.2	5