Qing-Nan Tang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6995474/publications.pdf

Version: 2024-02-01

218381 214527 2,826 105 26 47 citations g-index h-index papers 107 107 107 2312 docs citations times ranked citing authors all docs

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Establishment and Validation of Prognostic Nomograms for Endemic Nasopharyngeal Carcinoma. Journal of the National Cancer Institute, 2016, 108, djv291. | 3.0 | 281 |
| 2 | Neoadjuvant chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: A phase III multicentre randomised controlled trial. European Journal of Cancer, 2017, 75, 14-23. | 1.3 | 226 |
| 3 | Prospective Study of Tailoring Whole-Body Dual-Modality [¹⁸ F]Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography With Plasma Epstein-Barr Virus DNA for Detecting Distant Metastasis in Endemic Nasopharyngeal Carcinoma at Initial Staging. Journal of Clinical Oncology, 2013, 31, 2861-2869. | 0.8 | 171 |
| 4 | Induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in locoregionally advanced nasopharyngeal carcinoma: long-term results of a phase III multicentre randomised controlled trial. European Journal of Cancer, 2019, 119, 87-96. | 1.3 | 150 |
| 5 | Concurrent chemoradiotherapy with nedaplatin versus cisplatin in stage Il–IVB nasopharyngeal carcinoma: an open-label, non-inferiority, randomised phase 3 trial. Lancet Oncology, The, 2018, 19, 461-473. | 5.1 | 118 |
| 6 | Single-cell transcriptomic analysis defines the interplay between tumor cells, viral infection, and the microenvironment in nasopharyngeal carcinoma. Cell Research, 2020, 30, 950-965. | 5.7 | 111 |
| 7 | The Prognostic Value of Plasma Epstein-Barr Viral DNA and Tumor Response to Neoadjuvant Chemotherapy in Advanced-Stage Nasopharyngeal Carcinoma. International Journal of Radiation Oncology Biology Physics, 2015, 93, 862-869. | 0.4 | 110 |
| 8 | The Association Between the Development of Radiation Therapy, Image Technology, and Chemotherapy, and the Survival of Patients With Nasopharyngeal Carcinoma: A Cohort Study From 1990 to 2012. International Journal of Radiation Oncology Biology Physics, 2019, 105, 581-590. | 0.4 | 80 |
| 9 | Tumor CTLA-4 overexpression predicts poor survival in patients with nasopharyngeal carcinoma. Oncotarget, 2016, 7, 13060-13068. | 0.8 | 80 |
| 10 | Phase I trial of adoptively transferred tumor-infiltrating lymphocyte immunotherapy following concurrent chemoradiotherapy in patients with locoregionally advanced nasopharyngeal carcinoma. Oncolmmunology, 2015, 4, e976507. | 2.1 | 61 |
| 11 | The Prognostic Value of Treatment-Related Lymphopenia in Nasopharyngeal Carcinoma Patients. Cancer Research and Treatment, 2018, 50, 19-29. | 1.3 | 56 |
| 12 | EBV infection-induced GPX4 promotes chemoresistance and tumor progression in nasopharyngeal carcinoma. Cell Death and Differentiation, 2022, 29, 1513-1527. | 5.0 | 45 |
| 13 | A deep survival analysis method based on ranking. Artificial Intelligence in Medicine, 2019, 98, 1-9. | 3.8 | 44 |
| 14 | Optimal cumulative cisplatin dose in nasopharyngeal carcinoma patients based on induction chemotherapy response. Radiotherapy and Oncology, 2019, 137, 83-94. | 0.3 | 44 |
| 15 | High-Sensitivity C-Reactive Protein Complements Plasma Epstein-Barr Virus Deoxyribonucleic Acid Prognostication in Nasopharyngeal Carcinoma: A Large-Scale Retrospective and Prospective Cohort Study. International Journal of Radiation Oncology Biology Physics, 2015, 91, 325-336. | 0.4 | 41 |
| 16 | Ten-year outcomes of survival and toxicity for a phase III randomised trial of concurrent chemoradiotherapy versus radiotherapy alone in stage II nasopharyngeal carcinoma. European Journal of Cancer, 2019, 110, 24-31. | 1.3 | 40 |
| 17 | Combination of Tumor Volume and Epstein-Barr Virus DNA Improved Prognostic Stratification of Stage II Nasopharyngeal Carcinoma in the Intensity Modulated Radiotherapy Era: A Large-Scale Cohort Study. Cancer Research and Treatment, 2018, 50, 861-871. | 1.3 | 38 |
| 18 | Plasma Epstein-Barr viral DNA complements TNM classification of nasopharyngeal carcinoma in the era of intensity-modulated radiotherapy. Oncotarget, 2016, 7, 6221-6230. | 0.8 | 37 |

| # | Article | IF | CITATIONS |
|----|--|-------------|-----------|
| 19 | New surgical staging system for patients with recurrent nasopharyngeal carcinoma based on the AJCC/UICC rTNM classification system. European Journal of Cancer, 2015, 51, 1771-1779. | 1.3 | 36 |
| 20 | In-cell infection: a novel pathway for Epstein-Barr virus infection mediated by cell-in-cell structures. Cell Research, 2015, 25, 785-800. | 5. 7 | 36 |
| 21 | Identifying optimal candidates for local treatment of the primary tumor among patients with de novo metastatic nasopharyngeal carcinoma: a retrospective cohort study based on Epstein–Barr virus DNA level and tumor response to palliative chemotherapy. BMC Cancer, 2019, 19, 92. | 1.1 | 33 |
| 22 | High-density lipoprotein cholesterol as a predictor of poor survival in patients with nasopharyngeal carcinoma. Oncotarget, 2016, 7, 42978-42987. | 0.8 | 32 |
| 23 | Prognostic Value of Plasma Epstein–Barr Virus DNA for Local and Regionally Advanced Nasopharyngeal Carcinoma Treated With Cisplatin-Based Concurrent Chemoradiotherapy in Intensity-Modulated Radiotherapy Era. Medicine (United States), 2016, 95, e2642. | 0.4 | 29 |
| 24 | Symptom clusters in patients with nasopharyngeal carcinoma during radiotherapy. European Journal of Oncology Nursing, 2017, 28, 7-13. | 0.9 | 29 |
| 25 | Concurrent chemoradiotherapy with or without cetuximab for stage II to IVb nasopharyngeal carcinoma: a case–control study. BMC Cancer, 2017, 17, 567. | 1.1 | 29 |
| 26 | Is Hemoglobin Level in Patients with Nasopharyngeal Carcinoma Still a Significant Prognostic Factor in the Era of Intensity-Modulated Radiotherapy Technology?. PLoS ONE, 2015, 10, e0136033. | 1.1 | 28 |
| 27 | Different Prognostic Values of Plasma Epstein-Barr Virus DNA and Maximal Standardized Uptake Value of 18F-FDG PET/CT for Nasopharyngeal Carcinoma Patients with Recurrence. PLoS ONE, 2015, 10, e0122756. | 1.1 | 27 |
| 28 | Development and validation of the immune signature to predict distant metastasis in patients with nasopharyngeal carcinoma., 2020, 8, e000205. | | 26 |
| 29 | Deintensified Chemoradiotherapy for Pretreatment Epstein-Barr Virus DNA-Selected Low-Risk Locoregionally Advanced Nasopharyngeal Carcinoma: A Phase II Randomized Noninferiority Trial. Journal of Clinical Oncology, 2022, 40, 1163-1173. | 0.8 | 25 |
| 30 | FMNL1 mediates nasopharyngeal carcinoma cell aggressiveness by epigenetically upregulating MTA1. Oncogene, 2018, 37, 6243-6258. | 2.6 | 24 |
| 31 | The diagnostic and prognostic values of plasma Epsteinâ€Barr virus DNA for residual cervical lymphadenopathy in nasopharyngeal carcinoma patients: a retrospective study. Cancer Communications, 2019, 39, 1-13. | 3.7 | 24 |
| 32 | Combining pretreatment plasma Epsteinâ€Barr virus DNA level and cervical node necrosis improves prognostic stratification in patients with nasopharyngeal carcinoma: A cohort study. Cancer Medicine, 2019, 8, 6841-6852. | 1.3 | 22 |
| 33 | Effect of Induction Chemotherapy With Paclitaxel, Cisplatin, and Capecitabine vs Cisplatin and Fluorouracil on Failure-Free Survival for Patients With Stage IVA to IVB Nasopharyngeal Carcinoma. JAMA Oncology, 2022, 8, 706. | 3.4 | 22 |
| 34 | The impact of the cumulative dose of cisplatin during concurrent chemoradiotherapy on the clinical outcomes of patients with advanced-stage nasopharyngeal carcinoma in an era of intensity-modulated radiotherapy. BMC Cancer, 2015, 15, 977. | 1.1 | 21 |
| 35 | Famitinib in combination with concurrent chemoradiotherapy in patients with locoregionally advanced nasopharyngeal carcinoma: a phase 1, openâ€label, doseâ€escalation Study. Cancer Communications, 2018, 38, 1-13. | 3.7 | 20 |
| 36 | Targeting the IRAK1–S100A9 Axis Overcomes Resistance to Paclitaxel in Nasopharyngeal Carcinoma. Cancer Research, 2021, 81, 1413-1425. | 0.4 | 19 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Induction Chemotherapy Plus Concurrent Chemoradiotherapy Versus Concurrent Chemoradiotherapy Alone in Locoregionally Advanced Nasopharyngeal Carcinoma in Children and Adolescents: A Matched Cohort Analysis. Cancer Research and Treatment, 2018, 50, 1304-1315. | 1.3 | 19 |
| 38 | Efficacy of controlled-release oxycodone for reducing pain due to oral mucositis in nasopharyngeal carcinoma patients treated with concurrent chemoradiotherapy: a prospective clinical trial. Supportive Care in Cancer, 2019, 27, 3759-3767. | 1.0 | 18 |
| 39 | Subdivision of Nasopharyngeal Carcinoma Patients with Bone-Only Metastasis at Diagnosis for Prediction of Survival and Treatment Guidance. Cancer Research and Treatment, 2019, 51, 1259-1268. | 1.3 | 18 |
| 40 | With or without reirradiation in advanced local recurrent nasopharyngeal carcinoma: a caseâ€"control study. BMC Cancer, 2016, 16, 774. | 1.1 | 17 |
| 41 | Individualized concurrent chemotherapy by pretreatment plasma Epsteinâ€Barr viral DNA in Ilâ€II stage nasopharyngeal carcinoma: A propensity score matching analysis using a large cohort. Cancer Medicine, 2019, 8, 4214-4225. | 1.3 | 17 |
| 42 | Effect of local treatment for metastasis and its sequence with chemotherapy on prognosis of post-treatment metastatic nasopharyngeal carcinoma patients. Oral Oncology, 2019, 92, 40-45. | 0.8 | 17 |
| 43 | Deep learning signatures reveal multiscale intratumor heterogeneity associated with biological functions and survival in recurrent nasopharyngeal carcinoma. European Journal of Nuclear Medicine and Molecular Imaging, 2022, 49, 2972-2982. | 3.3 | 17 |
| 44 | The role of capecitabine as maintenance therapy in <i>de novo</i> metastatic nasopharyngeal carcinoma: A propensity score matching study. Cancer Communications, 2020, 40, 32-42. | 3.7 | 16 |
| 45 | The impact of smoking on the clinical outcome of locoregionally advanced nasopharyngeal carcinoma after chemoradiotherapy. Radiation Oncology, 2014, 9, 246. | 1.2 | 15 |
| 46 | Induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in stage III-IVb nasopharyngeal carcinoma patients with Epstein-Barr virus DNA ≥4000 copies/ml: a matched study. Oncotarget, 2016, 7, 29739-29748. | 0.8 | 15 |
| 47 | Relationship Between the Comprehensive Nutritional Index and the EORTC QLQ-H&N35 in Nasopharyngeal Carcinoma Patients Treated with Intensity-Modulated Radiation Therapy. Nutrition and Cancer, 2017, 69, 436-443. | 0.9 | 15 |
| 48 | A Randomized Controlled Trial Comparing Two Different Schedules for Cisplatin Treatment in Patients with Locoregionally Advanced Nasopharyngeal Cancer. Clinical Cancer Research, 2021, 27, 4186-4194. | 3.2 | 15 |
| 49 | Pretreatment Serum Amyloid A and C-reactive Protein Comparing with Epstein-Barr Virus DNA as Prognostic Indicators in Patients with Nasopharyngeal Carcinoma: A Prospective Study. Cancer Research and Treatment, 2018, 50, 701-711. | 1.3 | 14 |
| 50 | Liposomal paclitaxel versus docetaxel in induction chemotherapy using Taxanes, cisplatin and 5-fluorouracil for locally advanced nasopharyngeal carcinoma. BMC Cancer, 2018, 18, 1279. | 1.1 | 13 |
| 51 | Pretreatment quality of life as a predictor of survival for patients with nasopharyngeal carcinoma treated with IMRT. BMC Cancer, 2018, 18, 114. | 1.1 | 13 |
| 52 | <p>The development of a nomogram to predict post-radiation necrosis in nasopharyngeal carcinoma patients: a large-scale cohort study</p> . Cancer Management and Research, 2019, Volume 11, 6253-6263. | 0.9 | 13 |
| 53 | Establishment and validation of a nomogram for predicting survival in patients with de novo metastatic nasopharyngeal carcinoma. Oral Oncology, 2019, 94, 73-79. | 0.8 | 12 |
| 54 | Establishment and validation of a nomogram for predicting the benefit of concurrent chemotherapy in stage II nasopharyngeal carcinoma: A study based on a phase III randomized clinical trial with 10-year follow-up. Oral Oncology, 2020, 100, 104490. | 0.8 | 12 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Autocrine <scp>INSL</scp> 5 promotes tumor progression and glycolysis via activation of <scp>STAT</scp> 5 signaling. EMBO Molecular Medicine, 2020, 12, e12050. | 3.3 | 12 |
| 56 | Advanced-Stage Nasopharyngeal Carcinoma: Restaging System after Neoadjuvant Chemotherapy on the Basis of MR Imaging Determines Survival. Radiology, 2017, 282, 171-181. | 3.6 | 11 |
| 57 | Weekly versus triweekly cisplatin plus intensity-modulated radiotherapy in locally advanced nasopharyngeal carcinoma: A propensity score analysis with a large cohort. Journal of Cancer, 2018, 9, 3447-3455. | 1.2 | 11 |
| 58 | Patterns of Failure and Survival Trends Of 720 Patients with Stage I Nasopharyngeal Carcinoma Diagnosed from 1990-2012: A Large-scale Retrospective Cohort Study. Journal of Cancer, 2018, 9, 1308-1317. | 1.2 | 11 |
| 59 | The impact of Adult Comorbidity Evaluation-27 on the clinical outcome of elderly nasopharyngeal carcinoma patients treated with chemoradiotherapy or radiotherapy: a matched cohort analysis. Journal of Cancer, 2019, 10, 5614-5621. | 1.2 | 11 |
| 60 | Subdivision of de-novo metastatic nasopharyngeal carcinoma based on tumor burden and pretreatment EBV DNA for therapeutic guidance of locoregional radiotherapy. BMC Cancer, 2021, 21, 534. | 1.1 | 11 |
| 61 | Patterns of Failure and Survival Trends in 3,808 Patients with Stage II Nasopharyngeal Carcinoma Diagnosed from 1990 to 2012: A Large-Scale Retrospective Cohort Study. Cancer Research and Treatment, 2019, 51, 1449-1463. | 1.3 | 11 |
| 62 | Stratification of Candidates for Induction Chemotherapy in Stage III-IV Nasopharyngeal Carcinoma: A Large Cohort Study Based on a Comprehensive Prognostic Model. Frontiers in Oncology, 2020, 10, 255. | 1.3 | 10 |
| 63 | Establishment of a prognostic nomogram to identify optimal candidates for local treatment among patients with local recurrent nasopharyngeal carcinoma. Oral Oncology, 2020, 106, 104711. | 0.8 | 10 |
| 64 | Combining plasma Epstein-Barr virus DNA and nodal maximal standard uptake values of 18F-fluoro-2-deoxy-D-glucose positron emission tomography improved prognostic stratification to predict distant metastasis for locoregionally advanced nasopharyngeal carcinoma. Oncotarget, 2015, 6, 38296-38307. | 0.8 | 10 |
| 65 | Identifying distinct risks of treatment failure in nasopharyngeal carcinoma: A study based on the dynamic changes in peripheral blood lymphocytes, monocytes, N classification, and plasma Epsteinâ€Barr virus DNA. Head and Neck, 2021, , . | 0.9 | 10 |
| 66 | Development and validation of a transcriptomics-based gene signature to predict distant metastasis and guide induction chemotherapy in locoregionally advanced nasopharyngeal carcinoma. European Journal of Cancer, 2022, 163, 26-34. | 1.3 | 10 |
| 67 | Establishment of a prognostic scoring model for regional recurrent nasopharyngeal carcinoma after neck dissection. Cancer Biology and Medicine, 2020, 17, 227-236. | 1.4 | 9 |
| 68 | Effect of Concurrent Chemoradiotherapy With Nedaplatin vs Cisplatin on the Long-term Outcomes of Survival and Toxic Effects Among Patients With Stage II to IVB Nasopharyngeal Carcinoma. JAMA Network Open, 2021, 4, e2138470. | 2.8 | 9 |
| 69 | Maximal standard uptake values of 18F-fluoro-2-deoxy-D-glucose positron emission tomography compared with Epstein-Barr virus DNA as prognostic indicators in de novo metastatic nasopharyngeal carcinoma patients. BMC Cancer, 2019, 19, 908. | 1.1 | 8 |
| 70 | Establishment and validation of two nomograms to predict the benefit of concurrent chemotherapy in stage Ilâ€IVa nasopharyngeal carcinoma patients with different risk factors: Analysis based on a large cohort. Cancer Medicine, 2020, 9, 1661-1670. | 1.3 | 8 |
| 71 | Comparing three induction chemotherapy regimens for patients with locoregionally advanced nasopharyngeal carcinoma based on TNM stage and plasma Epstein–Barr virus DNA level. BMC Cancer, 2020, 20, 89. | 1.1 | 8 |
| 72 | Geriatric nutritional risk index as an independent prognostic factor in locally advanced nasopharyngeal carcinoma treated using radical concurrent chemoradiotherapy: a retrospective cohort study. Annals of Translational Medicine, 2021, 9, 532-532. | 0.7 | 8 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Irradiation-related longitudinal white matter atrophy underlies cognitive impairment in patients with nasopharyngeal carcinoma. Brain Imaging and Behavior, 2021, 15, 2426-2435. | 1.1 | 8 |
| 74 | Nomogram for the prediction of primary distant metastasis of nasopharyngeal carcinoma to guide individualized application of FDG PET/CT. European Journal of Nuclear Medicine and Molecular Imaging, 2021, 48, 2586-2598. | 3.3 | 8 |
| 75 | Optimal cumulative cisplatin dose in nasopharyngeal carcinoma patients based on plasma Epstein–Barr virus DNA level after induction chemotherapy. Aging, 2020, 12, 4931-4944. | 1.4 | 8 |
| 76 | Optimizing the Treatment Pattern for De Novo Metastatic Nasopharyngeal Carcinoma Patients: A Large-Scale Retrospective Cohort Study. Frontiers in Oncology, 2020, 10, 543646. | 1.3 | 7 |
| 77 | Intensive Local Radiotherapy Is Associated With Better Local Control and Prolonged Survival in Bone-Metastatic Nasopharyngeal Carcinoma Patients. Frontiers in Oncology, 2020, 10, 378. | 1.3 | 7 |
| 78 | Construction of a comprehensive nutritional index and comparison of its prognostic performance with the PNI and NRI for survival in older patients with nasopharyngeal carcinoma: a retrospective study. Supportive Care in Cancer, 2021, 29, 5371-5381. | 1.0 | 7 |
| 79 | Impact of smoking on survival in nasopharyngeal carcinoma: A cohort study with 23,325 patients diagnosed from 1990 to 2016. Radiotherapy and Oncology, 2021, 162, 7-17. | 0.3 | 7 |
| 80 | Epstein-Barr virus glycoprotein gH/gL antibodies complement IgA-viral capsid antigen for diagnosis of nasopharyngeal carcinoma. Oncotarget, 2016, 7, 16372-16383. | 0.8 | 7 |
| 81 | Utility of Epstein–Barr Virus DNA in Nasopharynx Swabs as a Reflex Test to Triage Seropositive Individuals in Nasopharyngeal Carcinoma Screening Programs. Clinical Chemistry, 2022, 68, 953-962. | 1.5 | 7 |
| 82 | Nomogram Predicting the Benefits of Adding Concurrent Chemotherapy to Intensity-Modulated Radiotherapy After Induction Chemotherapy in Stages II–IVb Nasopharyngeal Carcinoma. Frontiers in Oncology, 2020, 10, 539321. | 1.3 | 6 |
| 83 | Optimal sequencing of chemotherapy with chemoradiotherapy based on TNM stage classification and EBV DNA in locoregionally advanced nasopharyngeal carcinoma. Cancer Communications, 2019, 39, 1-3. | 3.7 | 5 |
| 84 | Induction chemotherapy followed by radiotherapy versus concurrent chemoradiotherapy in the treatment of different risk locoregionally advanced nasopharyngeal carcinoma. Therapeutic Advances in Medical Oncology, 2020, 12, 175883592092821. | 1.4 | 5 |
| 85 | Low value of whole-body dual-modality [18f]fluorodeoxyglucose positron emission tomography/computed tomography in primary staging of stage l–ll nasopharyngeal carcinoma: a nest case-control study. European Radiology, 2021, 31, 5222-5233. | 2.3 | 5 |
| 86 | Increased Angiogenin Expression Correlates With Radiation Resistance and Predicts Poor Survival for Patients With Nasopharyngeal Carcinoma. Frontiers in Pharmacology, 2021, 12, 627935. | 1.6 | 5 |
| 87 | Establishment and validation of a prognostic nomogram to predict early metastasis in nasopharyngeal carcinoma patients within six months after radiotherapy and to guide intensive treatment. Radiotherapy and Oncology, 2021, 162, 202-211. | 0.3 | 5 |
| 88 | The effect of adding concurrent chemotherapy to radiotherapy for stage II nasopharyngeal carcinoma with undetectable pretreatment Epstein-Barr virus DNA: Retrospective analysis with a large institutional-based cohort. Translational Oncology, 2021, 14, 100990. | 1.7 | 4 |
| 89 | Efficacy of Transnasal Endoscopic Fineâ€Needle Aspiration Biopsy in Diagnosing Submucosal Nasopharyngeal Carcinoma. Laryngoscope, 2021, 131, 1798-1804. | 1.1 | 4 |
| 90 | Do all patients with locoregionally advanced nasopharyngeal carcinoma benefit from the maintenance chemotherapy using S-1/capecitabine?. Oral Oncology, 2021, 122, 105539. | 0.8 | 4 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 91 | Prognostic effect of pregnancy on young female patients with nasopharyngeal carcinoma: results from a matched cohort analysis. Oncotarget, 2016, 7, 21913-21921. | 0.8 | 4 |
| 92 | Percent change in apparent diffusion coefficient and plasma EBV DNA after induction chemotherapy identifies distinct prognostic response phenotypes in advanced nasopharyngeal carcinoma. BMC Cancer, 2021, 21, 1320. | 1.1 | 4 |
| 93 | The prognosis of neck residue nasopharyngeal carcinoma (NPC) patients: results from a case-cohort study. Journal of Cancer, 2018, 9, 1765-1772. | 1.2 | 3 |
| 94 | Identifying optimal candidates for induction chemotherapy among stage II–IVa nasopharyngeal carcinoma based on pretreatment Epstein–Barr virus DNA and nodal maximal standard uptake values of [18 F]â€fluorodeoxyglucose positron emission tomography. Cancer Medicine, 2020, 9, 8852-8863. | 1.3 | 3 |
| 95 | Prognostic significance of a combined and controlled nutritional status score and EBV-DNA in patients with advanced nasopharyngeal carcinoma: a long-term follow-up study. Cancer Biology and Medicine, 2021, 19, 551-564. | 1.4 | 3 |
| 96 | Management of suboptimal response to induction chemotherapy in locoregionally advanced nasopharyngeal carcinoma: Re-induction therapy or direct to Radiotherapy?. Radiotherapy and Oncology, 2021, 163, 185-191. | 0.3 | 3 |
| 97 | Establishment and Validation of a Nomogram for Nasopharyngeal Carcinoma Patients Concerning the Prognostic Effect of Parotid Lymph Node Metastases. Cancer Research and Treatment, 2020, 52, 855-866. | 1.3 | 3 |
| 98 | Induction or adjuvant chemotherapy plus concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in paediatric nasopharyngeal carcinoma in the IMRT era: A recursive partitioning risk stratification analysis based on EBV DNA. European Journal of Cancer, 2021, 159, 133-143. | 1.3 | 3 |
| 99 | Role of zoledronic acid in nasopharyngeal carcinoma patients with bone-only metastasis at diagnosis. Oral Oncology, 2019, 97, 31-36. | 0.8 | 2 |
| 100 | Development and validation of a normal tissue complication probability model for acquired nasal cavity stenosis and atresia after radical radiotherapy for nasopharyngeal carcinoma. Radiotherapy and Oncology, 2021, 160, 9-17. | 0.3 | 2 |
| 101 | Impact of salvage radiotherapy on survival of patients with advanced locally recurrent nasopharyngeal carcinoma: Derivation and validation of a predictive model. Radiotherapy and Oncology, 2022, 167, 252-260. | 0.3 | 2 |
| 102 | Cost-Effectiveness analysis of combining plasma Epstein-Barr virus DNA testing and different surveillance imaging modalities for nasopharyngeal carcinoma patients in first remission. Oral Oncology, 2022, 128, 105851. | 0.8 | 2 |
| 103 | Longitudinal Trend of Health-Related Quality of Life During Concurrent Chemoradiotherapy and Survival in Patients With Stage Il–IVb Nasopharyngeal Carcinoma. Frontiers in Oncology, 2020, 10, 579292. | 1.3 | 1 |
| 104 | Divergent effects of irradiation on brain cortical morphology in patients with nasopharyngeal carcinoma: one-year follow-up study using structural magnetic resonance imaging. Quantitative Imaging in Medicine and Surgery, 2021, 11, 2307-2320. | 1.1 | 1 |
| 105 | Alpha-fetoprotein–producing recurrent nasopharyngeal carcinoma: A case report. SAGE Open Medical Case Reports, 2021, 9, 2050313X2110577. | 0.2 | 0 |