

# David G Mendoza-Cozatl

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6996641/publications.pdf>

Version: 2024-02-01

38  
papers

2,980  
citations

304602

22  
h-index

330025

37  
g-index

54  
all docs

54  
docs citations

54  
times ranked

3607  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Cadmium interference with iron sensing reveals transcriptional programs sensitive and insensitive to reactive oxygen species. <i>Journal of Experimental Botany</i> , 2022, 73, 324-338.   | 2.4 | 9         |
| 2  | Iron Availability within the Leaf Vasculature Determines the Magnitude of Iron Deficiency Responses in Source and Sink Tissues in <i>Arabidopsis</i> . <i>Plant and Cell Physiology</i> , 2022, 63, 829-841.                                 | 1.5 | 8         |
| 3  | Cross species multi-omics reveals cell wall sequestration and elevated global transcript abundance as mechanisms of boron tolerance in plants. <i>New Phytologist</i> , 2021, 230, 1985-2000.  | 3.5 | 25        |
| 4  | Exogenous 3,3'-Diindolylmethane Improves Vanadium Stress Tolerance in <i>Brassica napus</i> Seedling Shoots by Modulating Antioxidant Enzyme Activities. <i>Biomolecules</i> , 2021, 11, 436.  | 1.8 | 5         |
| 5  | Draft Genome Sequence of the Putative Endophytic Bacterium <i>Pantoea agglomerans</i> R6, Associated with <i>Lactuca serriola</i> from South Africa. <i>Microbiology Resource Announcements</i> , 2021, 10, .                                | 0.3 | 1         |
| 6  | Root-to-shoot iron partitioning in <i>Arabidopsis</i> requires IRON-REGULATED TRANSPORTER1 (IRT1) protein but not its iron(II) transport function. <i>Plant Journal</i> , 2021, , .  | 2.8 | 18        |
| 7  | Expression of a dominant-negative AtNEET-H89C protein disrupts iron-sulfur metabolism and iron homeostasis in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2020, 101, 1152-1169.  | 2.8 | 41        |
| 8  | Copper uptake mechanism of <i>Arabidopsis thaliana</i> high-affinity COPT transporters. <i>Protoplasma</i> , 2019, 256, 161-170.   | 1.0 | 31        |
| 9  | Keep talking: crosstalk between iron and sulfur networks fine-tunes growth and development to promote survival under iron limitation. <i>Journal of Experimental Botany</i> , 2019, 70, 4197-4210.   | 2.4 | 22        |
| 10 | Zinc uptake in the Basidiomycota: Characterization of zinc transporters in <i>Ustilago maydis</i> . <i>Molecular Membrane Biology</i> , 2019, 35, 39-50.   | 2.0 | 10        |
| 11 | Changes in iron availability in <i>Arabidopsis</i> are rapidly sensed in the leaf vasculature and impaired sensing leads to opposite transcriptional programs in leaves and roots. <i>Plant, Cell and Environment</i> , 2018, 41, 2263-2276. | 2.8 | 68        |
| 12 | Quantitative proteomics analysis of leaves from two <i>Sedum alfredii</i> ( <i>Crassulaceae</i> ) populations that differ in cadmium accumulation. <i>Proteomics</i> , 2017, 17, e1600456.   | 1.3 | 5         |
| 13 | Moderate to severe water limitation differentially affects the phenome and ionome of <i>Arabidopsis</i> . <i>Functional Plant Biology</i> , 2017, 44, 94.  | 1.1 | 35        |
| 14 | Common Bean: A Legume Model on the Rise for Unraveling Responses and Adaptations to Iron, Zinc, and Phosphate Deficiencies. <i>Frontiers in Plant Science</i> , 2016, 7, 600.  | 1.7 | 77        |
| 15 | Enhanced cadmium efflux and root-to-shoot translocation are conserved in the hyperaccumulator <i>Sedum alfredii</i> ( <i>Crassulaceae</i> family). <i>FEBS Letters</i> , 2016, 590, 1757-1764.   | 1.3 | 18        |
| 16 | Purification of Translating Ribosomes and Associated mRNAs from Soybean ( <i>Glycine max</i> ). <i>Current Protocols in Plant Biology</i> , 2016, 1, 185-196.  | 2.8 | 9         |
| 17 | Hydroponics: A Versatile System to Study Nutrient Allocation and Plant Responses to Nutrient Availability and Exposure to Toxic Elements. <i>Journal of Visualized Experiments</i> , 2016, , .   | 0.2 | 45        |
| 18 | Identification of AtOPT4 as a Plant Glutathione Transporter. <i>Molecular Plant</i> , 2016, 9, 481-484.  | 3.9 | 24        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Moving toward a precise nutrition: preferential loading of seeds with essential nutrients over non-essential toxic elements. <i>Frontiers in Plant Science</i> , 2014, 5, 51.   | 1.7 | 42        |
| 20 | Phytochelatinâ€‘metal(loid) transport into vacuoles shows different substrate preferences in barley and <i>Arabidopsis</i> . <i>Plant, Cell and Environment</i> , 2014, 37, 1192-1201.  | 2.8 | 134       |
| 21 | Zn-bis-glutathionate is the best co-substrate of the monomeric phytochelatin synthase from the photosynthetic heavy metal-hyperaccumulator <i>Euglena gracilis</i> . <i>Metallomics</i> , 2014, 6, 604.   | 1.0 | 13        |
| 22 | OPT3 Is a Component of the Iron-Signaling Network between Leaves and Roots and Misregulation of OPT3 Leads to an Over-Accumulation of Cadmium in Seeds. <i>Molecular Plant</i> , 2014, 7, 1455-1469.  | 3.9 | 135       |
| 23 | Elemental Concentrations in the Seed of Mutants and Natural Variants of <i>Arabidopsis thaliana</i> Grown under Varying Soil Conditions. <i>PLoS ONE</i> , 2013, 8, e63014.   | 1.1 | 19        |
| 24 | Feedback inhibition by thiols outranks glutathione depletion: a luciferaseâ€‘based screen reveals glutathioneâ€‘deficient $\gamma$ -ECS and glutathione synthetase mutants impaired in cadmiumâ€‘induced sulfate assimilation. <i>Plant Journal</i> , 2012, 70, 783-795.                | 2.8 | 60        |
| 25 | Long-distance transport, vacuolar sequestration, tolerance, and transcriptional responses induced by cadmium and arsenic. <i>Current Opinion in Plant Biology</i> , 2011, 14, 554-562.  | 3.5 | 366       |
| 26 | Tonoplast-localized Abc2 Transporter Mediates Phytochelatin Accumulation in Vacuoles and Confers Cadmium Tolerance. <i>Journal of Biological Chemistry</i> , 2010, 285, 40416-40426.  | 1.6 | 87        |
| 27 | Arsenic tolerance in <i>Arabidopsis</i> is mediated by two ABCC-type phytochelatin transporters. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 21187-21192.   | 3.3 | 555       |
| 28 | ARS5 is a component of the 26S proteasome complex, and negatively regulates thiol biosynthesis and arsenic tolerance in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2009, 59, 802-813.  | 2.8 | 64        |
| 29 | Identification of high levels of phytochelatin, glutathione and cadmium in the phloem sap of <i>Brassica napus</i> . A role for thiolâ€‘peptides in the longâ€‘distance transport of cadmium and the effect of cadmium on iron translocation. <i>Plant Journal</i> , 2008, 54, 249-259. | 2.8 | 311       |
| 30 | Thiol peptides induction in the seagrass <i>Thalassia testudinum</i> (Banks ex K nig) in response to cadmium exposure. <i>Aquatic Toxicology</i> , 2008, 86, 12-19.   | 1.9 | 20        |
| 31 | Cell wall composition affects Cd <sup>2+</sup> accumulation and intracellular thiol peptides in marine red algae. <i>Aquatic Toxicology</i> , 2007, 81, 65-72.  | 1.9 | 46        |
| 32 | Phytochelatin-cadmium-sulfide high-molecular-mass complexes of <i>Euglena gracilis</i> . <i>FEBS Journal</i> , 2006, 273, 5703-5713.  | 2.2 | 34        |
| 33 | Control of glutathione and phytochelatin synthesis under cadmium stress. Pathway modeling for plants. <i>Journal of Theoretical Biology</i> , 2006, 238, 919-936.   | 0.8 | 111       |
| 34 | Time-course development of the Cd <sup>2+</sup> hyper-accumulating phenotype in <i>Euglena gracilis</i> . <i>Archives of Microbiology</i> , 2005, 184, 83-92.   | 1.0 | 16        |
| 35 | Sulfur assimilation and glutathione metabolism under cadmium stress in yeast, protists and plants. <i>FEMS Microbiology Reviews</i> , 2005, 29, 653-671.  | 3.9 | 364       |
| 36 | Cd <sup>2+</sup> transport and storage in the chloroplast of <i>Euglena gracilis</i> . <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2005, 1706, 88-97.  | 0.5 | 58        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | The bacterial-like lactate shuttle components from heterotrophic <i>Euglena gracilis</i> . <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2005, 1709, 181-190. | 0.5 | 18        |
| 38 | Cadmium accumulation in the chloroplast of <i>Euglena gracilis</i> . <i>Physiologia Plantarum</i> , 2002, 115, 276-283.  | 2.6 | 66        |