Elisa Polledri

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7001175/publications.pdf

Version: 2024-02-01

all docs

45 810 18 25 g-index

49 49 49 1055

times ranked

citing authors

docs citations

#	Article	IF	CITATIONS
1	The role of salivary cortisol measured by liquid chromatography–tandem mass spectrometry in the diagnosis of subclinical hypercortisolism. European Journal of Endocrinology, 2013, 168, 289-296.	1.9	49
2	Biological monitoring of exposure to tebuconazole in winegrowers. Journal of Exposure Science and Environmental Epidemiology, 2014, 24, 643-649.	1.8	43
3	Development and validation of a gas chromatography/mass spectrometry method for the assessment of genomic DNA methylation. Rapid Communications in Mass Spectrometry, 2009, 23, 2637-2646.	0.7	38
4	Identification and Quantification of Metabolites of the Fungicide Tebuconazole in Human Urine. Chemical Research in Toxicology, 2014, 27, 1943-1949.	1.7	37
5	Global DNA methylation and low-level exposure to benzene. Medicina Del Lavoro, 2012, 103, 84-95.	0.3	36
6	An LC-MS/MS method to profile urinary mercapturic acids, metabolites of electrophilic intermediates of occupational and environmental toxicants. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2019, 1117, 66-76.	1.2	34
7	Highâ€throughput determination of cortisol, cortisone, and melatonin in oral fluid by onâ€line turbulent flow liquid chromatography interfaced with liquid chromatography/tandem mass spectrometry. Rapid Communications in Mass Spectrometry, 2013, 27, 1450-1460.	0.7	33
8	Determinants of active and environmental exposure to tobacco smoke and upper reference value of urinary cotinine in not exposed individuals. Environmental Research, 2016, 148, 154-163.	3.7	30
9	Urinary biomonitoring of subjects with different smoking habits. Part I: Profiling mercapturic acids. Toxicology Letters, 2020, 327, 48-57.	0.4	27
10	In Postmenopausal Female Subjects With Type 2 Diabetes Mellitus, Vertebral Fractures Are Independently Associated With Cortisol Secretion and Sensitivity. Journal of Clinical Endocrinology and Metabolism, 2015, 100, 1417-1425.	1.8	23
11	Assessment of penconazole exposure in winegrowers using urinary biomarkers. Environmental Research, 2019, 168, 54-61.	3.7	23
12	Human biomonitoring of polycyclic aromatic hydrocarbonsand metals in the general population residing near the municipal solid waste incinerator of Modena, Italy. Chemosphere, 2017, 186, 546-557.	4.2	22
13	Epigenetic and Transcriptional Modifications in Repetitive Elements in Petrol Station Workers Exposed to Benzene and MTBE. International Journal of Environmental Research and Public Health, 2018, 15, 735.	1.2	22
14	A Validated Method for Urinary Cotinine Quantification Used to Classify Active and Environmental Tobacco Smoke Exposure. Current Analytical Chemistry, 2013, 9, 447-456.	0.6	22
15	Long-term occupational and environmental exposure to penconazole and tebuconazole by hair biomonitoring. Toxicology Letters, 2018, 298, 19-24.	0.4	21
16	Determination of tebuconazole and penconazole fungicides in rat and human hair by liquid chromatography/tandem mass spectrometry. Rapid Communications in Mass Spectrometry, 2018, 32, 1243-1249.	0.7	21
17	Association of Urinary and Dietary Selenium and of Serum Selenium Species with Serum Alanine Aminotransferase in a Healthy Italian Population. Antioxidants, 2021, 10, 1516.	2.2	21
18	Cortisol Secretion, Sensitivity, and Activity Are Associated With Hypertension in Postmenopausal Eucortisolemic Women. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 4441-4448.	1.8	18

#	Article	IF	Citations
19	Development and validation of an LC–MS/MS method for the quantitation of 30 legacy and emerging per- and polyfluoroalkyl substances (PFASs) in human plasma, including HFPO-DA, DONA, and cC6O4. Analytical and Bioanalytical Chemistry, 2022, 414, 1259-1278.	1.9	18
20	Biomonitoring short- and long-term exposure to the herbicide terbuthylazine in agriculture workers and in the general population using urine and hair specimens. Environment International, 2013, 60, 42-47.	4.8	17
21	Immunosuppressive drugs in whole blood: validation of a commercially available liquid chromatography/tandem mass spectrometry kit and comparison with immunochemical assays. Rapid Communications in Mass Spectrometry, 2017, 31, 1111-1120.	0.7	17
22	Associations between Urinary and Dietary Selenium and Blood Metabolic Parameters in a Healthy Northern Italy Population. Antioxidants, 2021, 10, 1193.	2.2	16
23	Biological Monitoring of Occupational Exposure to Polycyclic Aromatic Hydrocarbons at an Electric Steel Foundry in Tunisia. Annals of Occupational Hygiene, 2016, 60, 700-716.	1.9	15
24	Identification of Metabolites of the Fungicide Penconazole in Human Urine. Chemical Research in Toxicology, 2016, 29, 1179-1186.	1.7	15
25	Associations of urinary and dietary cadmium with urinary 8-oxo-7,8-dihydro-2′-deoxyguanosine and blood biochemical parameters. Environmental Research, 2022, 210, 112912.	3.7	14
26	Development of a method to profile 2- to 4-ring polycyclic aromatic hydrocarbons in saliva samples from smokers and non-smokers by headspace-solid-phase microextraction-gas chromatography-triple quadrupole tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2020, 1152, 122273.	1.2	13
27	Prediction of hypertension, diabetes and fractures in eucortisolemic women by measuring parameters of cortisol milieu. Endocrine, 2020, 68, 411-419.	1.1	13
28	Preoperative treatment with metyrapone in patients with Cushing's syndrome due to adrenal adenoma: a pilot prospective study. Endocrine Connections, 2018, 7, 1227-1235.	0.8	13
29	Environmental and biological monitoring of personal exposure to air pollutants of adult people living in a metropolitan area. Science of the Total Environment, 2021, 767, 144916.	3.9	12
30	Terbuthylazine in hair as a biomarker of exposure. Toxicology Letters, 2012, 210, 169-173.	0.4	11
31	Biological Monitoring of Occupational Exposure to Metals in Electric Steel Foundry Workers and Its Contribution to 8-Oxo-7,8-Dihydro-2′-Deoxyguanosine Levels. International Journal of Environmental Research and Public Health, 2020, 17, 1811.	1.2	11
32	Dermal exposure and risk assessment of tebuconazole applicators in vineyards. Medicina Del Lavoro, 2015, 106, 294-315.	0.3	11
33	Adrenalectomy Improves Blood Pressure and Metabolic Control in Patients With Possible Autonomous Cortisol Secretion: Results of a RCT. Frontiers in Endocrinology, 2022, 13, .	1.5	11
34	Urinary biomonitoring of subjects with different smoking habits. Part II: an untargeted metabolomic approach and the comparison with the targeted measurement of mercapturic acids. Toxicology Letters, 2020, 329, 56-66.	0.4	10
35	Development and validation of a liquid chromatography/tandem mass spectrometry method to quantify metabolites of phthalates, including diâ€2â€ethylhexyl terephthalate (DEHTP) and bisphenol A, in human urine. Rapid Communications in Mass Spectrometry, 2020, 34, e8796.	0.7	10
36	The activity of $11\hat{1}^2$ -hydroxysteroid dehydrogenase type 2 enzyme and cortisol secretion in patients with adrenal incidentalomas. Endocrine, 2016, 53, 809-815.	1.1	9

#	Article	IF	CITATIONS
37	Cumulative Pesticides Exposure of Children and Their Parents Living near Vineyards by Hair Analysis. International Journal of Environmental Research and Public Health, 2021, 18, 3723.	1.2	8
38	Determination of terbuthylazine and desethylterbuthylazine in human urine and hair samples by eletrospray ionization-liquid chromatography/triple quadrupole mass spectrometry. Analytical and Bioanalytical Chemistry, 2012, 404, 875-886.	1.9	7
39	Urinary 8-Oxo-7,8-Dihydro-2′-Deoxyguanosine in Tunisian Electric Steel Foundry Workers Exposed to Polycyclic Aromatic Hydrocarbons. Annals of Work Exposures and Health, 2017, 61, 333-343.	0.6	7
40	Oral Vitamin D supplementation impacts gene expression in granulosa cells in women undergoing IVF. Human Reproduction, 2020, 36, 130-144.	0.4	7
41	A liquid chromatography tandem mass spectrometry method to assess 41 pesticides in human hair. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2020, 1159, 122389.	1.2	5
42	Simultaneous Quantification of Bisphenol A, Its Glucuronide Metabolite, and Commercial Alternatives by LC-MS/MS for <i>In Vitro</i> Skin Absorption Evaluation. Chemical Research in Toxicology, 2020, 33, 2390-2400.	1.7	3
43	Dermal exposure to the fungicide tebuconazole during application in vineyards. Toxicology Letters, 2012, 211, S172.	0.4	2
44	ETS Exposure and PAH Body Burden in Nonsmoking Italian Adults. International Journal of Environmental Research and Public Health, $2018,15,1156.$	1.2	2
45	Biomonitoring pesticide exposure in nonconventional specimens. , 2021, , 245-281.		O