

Daniel T Lilly

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7004504/publications.pdf>

Version: 2024-02-01

23
papers

214
citations

1163117

8
h-index

1058476

14
g-index

24
all docs

24
docs citations

24
times ranked

317
citing authors

#	ARTICLE	IF	CITATIONS
1	An Assessment of Nonoperative Management Strategies in a Herniated Lumbar Disc Population: Successes Versus Failures. <i>Global Spine Journal</i> , 2021, 11, 1054-1063.	2.3	5
2	Years of potential life lost secondary to COVID-19: Cook County, Illinois. <i>Annals of Epidemiology</i> , 2021, 58, 124-127.	1.9	6
3	Regional Variation in Nonoperative Therapy Utilization for Symptomatic Lumbar Stenosis and Spondylolisthesis: A 2-Year Costs Analysis. <i>Global Spine Journal</i> , 2020, 10, 138-147.	2.3	4
4	Racial Differences in Perioperative Opioid Utilization in Lumbar Decompression and Fusion Surgery for Symptomatic Lumbar Stenosis or Spondylolisthesis. <i>Global Spine Journal</i> , 2020, 10, 160-168.	2.3	5
5	Total Health Care Expenditure in Patients With a Herniated Lumbar Disk That Ultimately Require Surgery. <i>Clinical Spine Surgery</i> , 2020, 33, E108-E115.	1.3	1
6	Adjunct Analgesia Reduces Pain and Opioid Consumption After Hip Arthroscopy: A Systematic Review of Randomized Controlled Trials. <i>American Journal of Sports Medicine</i> , 2020, 48, 3638-3651.	4.2	18
7	Gender differences in the 3-month utilization of nonoperative therapies prior to primary lumbar microdiscectomy. <i>Journal of Clinical Neuroscience</i> , 2020, 76, 107-113.	1.5	3
8	Gender differences in use of prolonged non-operative therapies prior to index ACDF surgery. <i>Journal of Clinical Neuroscience</i> , 2020, 78, 228-235.	1.5	1
9	Reduction in Narcotic Use After Lumbar Decompression and Fusion in Patients With Symptomatic Lumbar Stenosis or Spondylolisthesis. <i>Global Spine Journal</i> , 2019, 9, 598-606.	2.3	4
10	A 2-Year Cost Analysis of Maximum Nonoperative Treatments in Patients With Symptomatic Lumbar Stenosis or Spondylolisthesis That Ultimately Required Surgery. <i>Global Spine Journal</i> , 2019, 9, 424-433.	2.3	7
11	A Two-Year Cost Analysis of Maximum Nonoperative Treatments in Patients with Cervical Stenosis that Ultimately Required Surgery. <i>World Neurosurgery</i> , 2019, 124, e616-e625.	1.3	2
12	Sex Differences in Opioid Use in Patients With Symptomatic Lumbar Stenosis or Spondylolisthesis Undergoing Lumbar Decompression and Fusion. <i>Spine</i> , 2019, 44, E800-E807.	2.0	14
13	Regional Differences in the Cost and Utilization of Nonoperative Management Within 3 Months Prior to Lumbar Microdiscectomy. <i>Spine</i> , 2019, 44, 1571-1577.	2.0	4
14	Long-Term Costs of Maximum Nonoperative Treatments in Patients With Symptomatic Lumbar Stenosis or Spondylolisthesis that Ultimately Required Surgery. <i>Spine</i> , 2019, 44, 424-430.	2.0	25
15	Regional Variation in Opioid Use After Lumbar Spine Surgery. <i>World Neurosurgery</i> , 2019, 121, e691-e699.	1.3	22
16	Extended Length of Stay After Lumbar Spine Surgery: Sick Patients, Postoperative Complications, or Practice Style Differences Among Hospitals and Physicians?. <i>World Neurosurgery</i> , 2019, 123, e734-e739.	1.3	33
17	Opioid Prescribing Practices of Neurosurgeons: Analysis of Medicare Part D. <i>World Neurosurgery</i> , 2018, 112, e31-e38.	1.3	7
18	Extended Length of Stay in Elderly Patients after Anterior Cervical Discectomy and Fusion Is Not Attributable to Baseline Illness Severity or Postoperative Complications. <i>World Neurosurgery</i> , 2018, 115, e552-e557.	1.3	16

#	ARTICLE	IF	CITATIONS
19	Limited post-operative dexamethasone use does not affect lumbar fusion: a single institutional experience. <i>Journal of Spine Surgery</i> , 2018, 4, 254-259.	1.2	2
20	The use of subfascial drains after multi-level anterior cervical discectomy and fusion: does the data support its use?. <i>Journal of Spine Surgery</i> , 2018, 4, 227-232.	1.2	7
21	Gender Differences in Use of Prolonged Nonoperative Therapies Before Index Lumbar Surgery. <i>World Neurosurgery</i> , 2018, 120, e580-e592.	1.3	9
22	Extended Length of Stay in Elderly Patients After Lumbar Decompression and Fusion Surgery May Not Be Attributable to Baseline Illness Severity or Postoperative Complications. <i>World Neurosurgery</i> , 2018, 116, e996-e1001.	1.3	11
23	Does Nasal Carriage of <i>Staphylococcus aureus</i> Increase the Risk of Postoperative Infections After Elective Spine Surgery: Do Most Infections Occur in Carriers?. <i>World Neurosurgery</i> , 2018, 116, e519-e524.	1.3	6