

Slawomir Wojcik

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7005891/publications.pdf>

Version: 2024-02-01

35
papers

564
citations

566801

15
h-index

642321

23
g-index

35
all docs

35
docs citations

35
times ranked

808
citing authors

#	ARTICLE	IF	CITATIONS
1	Crosstalk between autophagy and proteasome protein degradation systems: possible implications for cancer therapy. <i>Folia Histochemica Et Cytobiologica</i> , 2014, 51, 249-264.	0.6	59
2	Endoplasmic reticulum stress induces myostatin precursor protein and NF- κ B in cultured human muscle fibers: Relevance to inclusion body myositis. <i>Experimental Neurology</i> , 2007, 204, 610-618.	2.0	50
3	Age related skeletal muscle atrophy and upregulation of autophagy in dogs. <i>Veterinary Journal</i> , 2015, 206, 54-60.	0.6	36
4	Myostatin is increased and complexes with amyloid- β^2 within sporadic inclusion-body myositis muscle fibers. <i>Acta Neuropathologica</i> , 2005, 110, 173-177.	3.9	35
5	NOGO is increased and binds to BACE1 in sporadic inclusion-body myositis and in A β 2PP-overexpressing cultured human muscle fibers. <i>Acta Neuropathologica</i> , 2007, 114, 517-526.	3.9	34
6	Canine inflammatory myopathy associated with <i>Leishmania Infantum</i> infection. <i>Neuromuscular Disorders</i> , 2009, 19, 124-130.	0.3	33
7	A β 2PP-overexpression and proteasome inhibition increase β -crystallin in cultured human muscle: Relevance to inclusion-body myositis. <i>Neuromuscular Disorders</i> , 2006, 16, 839-844.	0.3	29
8	Neuropeptide-containing neurons in the endopiriform region of the rat: morphology and colocalization with calcium-binding proteins and nitric oxide synthase. <i>Brain Research</i> , 2004, 996, 97-110.	1.1	27
9	In inclusion-body myositis muscle fibers Parkinson-associated DJ-1 is increased and oxidized. <i>Free Radical Biology and Medicine</i> , 2008, 45, 773-779.	1.3	24
10	Colocalization of neuropeptides with calcium-binding proteins in the claustral interneurons during postnatal development of the rat. <i>Brain Research Bulletin</i> , 2009, 80, 100-106.	1.4	22
11	Syrian hamster infected with <i>Leishmania infantum</i> : A new experimental model for inflammatory myopathies. <i>Muscle and Nerve</i> , 2010, 41, 355-361.	1.0	22
12	Expression of Nogo-66 in human muscle fibers is not specific for amyotrophic lateral sclerosis. <i>Annals of Neurology</i> , 2007, 62, 676-677.	2.8	18
13	Amyloid precursor protein, lipofuscin accumulation and expression of autophagy markers in aged bovine brain. <i>BMC Veterinary Research</i> , 2017, 13, 102.	0.7	18
14	Age-Related Changes in Skeletal Muscle of Cattle. <i>Veterinary Pathology</i> , 2016, 53, 436-446.	0.8	17
15	Myostatin precursor protein is increased and associates with amyloid- β precursor protein in inclusion-body myositis culture model. <i>Neuropathology and Applied Neurobiology</i> , 2007, 33, 238-242.	1.8	16
16	NPY-, SOM- and VIP-containing interneurons in postnatal development of the rat claustrum. <i>Brain Research Bulletin</i> , 2008, 76, 565-571.	1.4	16
17	Postnatal development of the basolateral complex of rabbit amygdala: a stereological and histochemical study. <i>Journal of Anatomy</i> , 2003, 203, 513-521.	0.9	13
18	Proteasome inhibitors against amelanotic melanoma. <i>Cell Biology and Toxicology</i> , 2017, 33, 557-573.	2.4	13

#	ARTICLE	IF	CITATIONS
19	Tobacco smoking alters the number of oral epithelial cells with apoptotic features. <i>Folia Histochemica Et Cytobiologica</i> , 2014, 52, 60-68.	0.6	13
20	Variations in popliteal fossa venous anatomy: implications for diagnosis of deep-vein thrombosis. <i>Folia Morphologica</i> , 2013, 72, 51-56.	0.4	13
21	Original article Transcription factor Pax6 is expressed by astroglia after transient brain ischemia in the rat model. <i>Folia Neuropathologica</i> , 2013, 3, 203-213.	0.5	10
22	Quantitative analysis of influence of sevoflurane on the reactivity of microglial cells in the course of the experimental model of intracerebral haemorrhage. <i>European Journal of Anaesthesiology</i> , 2006, 23, 874-881.	0.7	7
23	Noncovalent inhibitors of human 20S and 26S proteasome based on trypsin inhibitor SFTI-1. <i>Biopolymers</i> , 2016, 106, 685-696.	1.2	7
24	Expression of the calcium-binding proteins in the central, medial and cortical nuclei of the rabbit amygdaloid complex during postnatal development. <i>Acta Neurobiologiae Experimentalis</i> , 2013, 73, 260-79.	0.4	7
25	Original article Nigrostriatal pathway degeneration in rats after intraperitoneal administration of proteasome inhibitor MG-132. <i>Folia Neuropathologica</i> , 2014, 1, 41-55.	0.5	6
26	Qualitative and quantitative study of the postnatal development of the rabbit claustrum. <i>International Journal of Developmental Neuroscience</i> , 2002, 20, 113-123.	0.7	5
27	The dentin sialoprotein (DSP) expression in rat tooth germs following fluoride treatment: An immunohistochemical study. <i>Archives of Oral Biology</i> , 2006, 51, 252-261.	0.8	4
28	Morphological Changes within the Rat Lateral Ventricle after the Administration of Proteasome Inhibitors. <i>PLoS ONE</i> , 2015, 10, e0140536.	1.1	4
29	G.P.2 01 Alpha-synuclein and parkin are novel proteins accumulated in ragged red fibers. <i>Neuromuscular Disorders</i> , 2006, 16, 657.	0.3	2
30	G.P.2 02 In cultured human muscle fibers (CHMFs) amyloid- β precursor protein (A β PP) and proteasome inhibition increase I β -crystallin (I β BC). Relevance to sporadic inclusion-body myositis (s-IBM). <i>Neuromuscular Disorders</i> , 2006, 16, 657.	0.3	1
31	G.P.13.15 In sporadic inclusion-body myositis muscle-fiber (s-IBM) cytoplasm, cytochrome C aggregates with I β -synuclein and amyloid- β precursor protein (A β PP), but does not activate caspase-3. <i>Neuromuscular Disorders</i> , 2007, 17, 853.	0.3	1
32	Reduced level of synapsin I protein in the rat striatum after intraventricular administration of proteasome inhibitors: preliminary studies. <i>Folia Morphologica</i> , 2015, 74, 428-433.	0.4	1
33	The anatomical relationships between the serotonergic afferents and the neurons containing calcium-binding proteins in the rat claustrum. <i>Acta Neurobiologiae Experimentalis</i> , 2006, 66, 33-42.	0.4	1
34	G.O.2 In sporadic inclusion-body myositis (s-IBM) muscle fibers, Parkinson-disease-associated DJ-1 is oxidized and might play a novel pathogenic role. <i>Neuromuscular Disorders</i> , 2007, 17, 766-767.	0.3	0
35	G.P.3.16 In skeletal muscle of patients with type-II fiber atrophy, myostatin (MSTN) and myostatin precursor protein (MSTN-PP) are both increased. <i>Neuromuscular Disorders</i> , 2007, 17, 787.	0.3	0