

Luiz Sergio F De Carvalho

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/701062/luiz-sergio-f-de-carvalho-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

53
papers

510
citations

11
h-index

21
g-index

65
ext. papers

683
ext. citations

3.9
avg, IF

4.07
L-index

#	Paper	IF	Citations
53	Rationale and design of the Brazilian Diabetes Study: a prospective cohort of type 2 diabetes.. <i>Current Medical Research and Opinion</i> , 2022 , 1-13	2.5	1
52	Dapagliflozin reduces adiposity and increases adiponectin in patients with type 2 diabetes and atherosclerotic disease at short-term: An active-controlled randomised trial. <i>Diabetes and Metabolism</i> , 2021 , 48, 101304	5.4	0
51	Dapagliflozin effect on endothelial dysfunction in diabetic patients with atherosclerotic disease: a randomized active-controlled trial. <i>Cardiovascular Diabetology</i> , 2021 , 20, 74	8.7	16
50	Intra-operative esmolol and pain following mastectomy: A randomised clinical trial. <i>European Journal of Anaesthesiology</i> , 2021 , 38, 735-743	2.3	4
49	Cardiovascular safety of naltrexone and bupropion therapy: Systematic review and meta-analyses. <i>Obesity Reviews</i> , 2021 , 22, e13224	10.6	0
48	Dapagliflozin increases the lean-to total mass ratio in type 2 diabetes mellitus. <i>Nutrition and Diabetes</i> , 2021 , 11, 17	4.7	1
47	Glucose-lowering Drugs and Hospitalization for Heart Failure: A Systematic Review and Additive-effects Network Meta-analysis With More Than 500 000 Patient-years. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, 3060-3067	5.6	1
46	Defective Allele of the Neuronal Nitric Oxide Synthase Gene Increases Insulin Resistance During Acute Phase of Myocardial Infarction. <i>International Journal of General Medicine</i> , 2021 , 14, 3669-3676	2.3	
45	Increased particle size of triacylglycerol-enriched remnant lipoproteins, but not their plasma concentration or lipid content, augments risk prediction of incident type 2 diabetes. <i>Diabetologia</i> , 2021 , 64, 385-396	10.3	9
44	Gensini Score and Thrombus Burden Add Predictive Value to the SYNTAX Score in Detecting No-Reflow after Myocardial Infarction. <i>Arquivos Brasileiros De Cardiologia</i> , 2021 , 116, 466-472	1.2	0
43	The Impact of Education on All-cause Mortality Following St-Segment Elevation Myocardial Infarction (STEMI): Results from the Brazilian Heart Study. <i>Arquivos Brasileiros De Cardiologia</i> , 2021 , 117, 5-12	1.2	0
42	The impact of low income on long-term mortality of myocardial infarction patients: results from the Brazilian Heart Study. <i>Current Medical Research and Opinion</i> , 2021 , 37, 1689-1695	2.5	
41	Dapagliflozin increases retinal thickness in type 2 diabetic patients as compared with glibenclamide: A randomized controlled trial. <i>Diabetes and Metabolism</i> , 2021 , 47, 101280	5.4	1
40	Letter to the Editor: "Cardiovascular Effects of Pioglitazone or Sulfonylureas According to Pretreatment Risk: Moving Toward Personalized Care". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	
39	Synergistic effect of the association between lidocaine and magnesium sulfate on peri-operative pain after mastectomy: A randomised, double-blind trial. <i>European Journal of Anaesthesiology</i> , 2020 , 37, 224-234	2.3	5
38	Statin Use in the Early Phase of ST-Segment Elevation Myocardial Infarction Is Associated With Decreased QTc Dispersion. <i>Journal of Cardiovascular Pharmacology and Therapeutics</i> , 2020 , 25, 226-231	2.6	0
37	1480-P: Increased Particle Size of Triglyceride Remnant Lipoproteins, but Not Plasma Concentration or Lipid Content, Boost Risk Prediction of Incident Type 2 Diabetes. <i>Diabetes</i> , 2020 , 69, 1480-P	0.9	

36	Lower bone mass is associated with subclinical atherosclerosis, endothelial dysfunction and carotid thickness in the very elderly. <i>Atherosclerosis</i> , 2020 , 292, 70-74	3.1	4
35	Excess weight mediates changes in HDL pool that reduce cholesterol efflux capacity and increase antioxidant activity. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 254-264	4.5	5
34	Machine Learning Improves the Identification of Individuals With Higher Morbidity and Avoidable Health Costs After Acute Coronary Syndromes. <i>Value in Health</i> , 2020 , 23, 1570-1579	3.3	4
33	Rationale and design of the expanded combination of evolocumab plus empagliflozin in diabetes: EXCEED-BHS3 trial. <i>Therapeutic Advances in Chronic Disease</i> , 2020 , 11, 2040622320959248	4.9	4
32	Reciprocal Multifaceted Interaction Between HDL (High-Density Lipoprotein) and Myocardial Infarction. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2019 , 39, 1550-1564	9.4	9
31	Omega-3 intake is associated with attenuated inflammatory response and cardiac remodeling after myocardial infarction. <i>Nutrition Journal</i> , 2019 , 18, 29	4.3	6
30	Intensive treatment of hyperglycemia in the acute phase of myocardial infarction: the tenuous balance between effectiveness and safety - a systematic review and meta-analysis of randomized clinical trials. <i>Revista Da Associação Médica Brasileira</i> , 2019 , 65, 24-32	1.4	1
29	Prevalence, treatment, and control of dyslipidemia in diabetic participants of two brazilian cohorts: a place far from heaven. <i>Revista Da Associação Médica Brasileira</i> , 2019 , 65, 3-8	1.4	1
28	Inhibition of the sodium-glucose co-transporter 2 in the elderly: clinical and mechanistic insights into safety and efficacy. <i>Revista Da Associação Médica Brasileira</i> , 2019 , 65, 70-86	1.4	9
27	Statin Short-term Inhibition of Insulin Sensitivity and Secretion During Acute Phase of ST-Elevation Myocardial Infarction. <i>Scientific Reports</i> , 2019 , 9, 16401	4.9	2
26	Updated Cardiovascular Prevention Guideline of the Brazilian Society of Cardiology - 2019. <i>Arquivos Brasileiros De Cardiologia</i> , 2019 , 113, 787-891	1.2	29
25	HDL-Targeted Therapies During Myocardial Infarction. <i>Cardiovascular Drugs and Therapy</i> , 2019 , 33, 371-384	3.1	10
24	Adverse interaction between HDL and the mass of myocardial infarction. <i>Atherosclerosis</i> , 2019 , 281, 9-16	3.1	4
23	Change of BNP between admission and discharge after ST-elevation myocardial infarction (Killip I) improves risk prediction of heart failure, death, and recurrent myocardial infarction compared to single isolated measurement in addition to the GRACE score. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2019 , 8, 643-651	4.3	9
22	Relevance of AND-ASPEN criteria of malnutrition to predict hospital mortality in critically ill patients: A prospective study. <i>Journal of Critical Care</i> , 2018 , 44, 398-403	4	19
21	Adverse outcome has a U-shaped relation with acute phase change in insulin sensitivity after ST-Elevation Myocardial Infarction. <i>International Journal of Cardiology</i> , 2018 , 254, 16-22	3.2	1
20	Adiponectin concentration data improve the estimation of atherosclerotic risk in normal and in overweight subjects. <i>Clinical Endocrinology</i> , 2018 , 88, 388-396	3.4	1
19	Response to Comment on de Carvalho et al. Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Inhibitors and Incident Type 2 Diabetes: A Systematic Review and Meta-analysis With Over 96,000 Patient-Years. <i>Diabetes Care</i> 2018;41:364-367. <i>Diabetes Care</i> , 2018 , 41, e70-e71	14.6	

18	Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Inhibitors and Incident Type 2 Diabetes: A Systematic Review and Meta-analysis With Over 96,000 Patient-Years. <i>Diabetes Care</i> , 2018 , 41, 364-367	14.6	58
17	GLP-1RAs in type 2 diabetes: mechanisms that underlie cardiovascular effects and overview of cardiovascular outcome data. <i>Cardiovascular Diabetology</i> , 2018 , 17, 157	8.7	60
16	Statin-associated muscle symptoms: position paper from the Luso-Latin American Consortium. <i>Current Medical Research and Opinion</i> , 2017 , 33, 239-251	2.5	11
15	TCF7L2 polymorphism is associated with low nitric oxide release, endothelial dysfunction and enhanced inflammatory response after myocardial infarction. <i>BBA Clinical</i> , 2016 , 5, 159-65		2
14	Vitamin D for the prevention of cardiovascular disease: Are we ready for that?. <i>Atherosclerosis</i> , 2015 , 241, 729-40	3.1	47
13	Glycosylated hemoglobin is associated with decreased endothelial function, high inflammatory response, and adverse clinical outcome in non-diabetic STEMI patients. <i>Atherosclerosis</i> , 2015 , 243, 124-30	3.1	14
12	Endothelial nitric oxide synthase genotypes modulate peripheral vasodilatory properties after myocardial infarction. <i>Gene</i> , 2015 , 568, 165-9	3.8	6
11	HDL levels and oxidizability during myocardial infarction are associated with reduced endothelial-mediated vasodilation and nitric oxide bioavailability. <i>Atherosclerosis</i> , 2014 , 237, 840-6	3.1	23
10	Elevated CETP activity during acute phase of myocardial infarction is independently associated with endothelial dysfunction and adverse clinical outcome. <i>Atherosclerosis</i> , 2014 , 237, 777-83	3.1	17
9	Validation of surrogate indexes of insulin sensitivity in acute phase of myocardial infarction based on euglycemic-hyperinsulinemic clamp. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2014 , 306, E399-403	6	10
8	High-density lipoprotein levels are strongly associated with the recovery rate of insulin sensitivity during the acute phase of myocardial infarction: a study by euglycemic hyperinsulinemic clamp. <i>Journal of Clinical Lipidology</i> , 2013 , 7, 24-8	4.9	4
7	Pericardial fat volume measured by non-contrast tomography improves risk prediction of subclinical coronary artery disease estimated by coronary artery calcium score and Framingham risk score. <i>European Heart Journal</i> , 2013 , 34, P3137-P3137	9.5	
6	High cholesteryl ester transfer protein (CETP) activity in acute phase of myocardial infarction predicts short-term mortality by inducing endothelial dysfunction and reduced microvascular perfusion. <i>European Heart Journal</i> , 2013 , 34, P475-P475	9.5	
5	Validation of HOMA2S in acute phase of myocardial infarction, based on euglycemic-hyperinsulinemic clamp. <i>European Heart Journal</i> , 2013 , 34, P1272-P1272	9.5	
4	High plasma HDL-C attenuates stress hyperglycemia during acute phase of myocardial infarction. <i>Atherosclerosis</i> , 2012 , 220, 231-6	3.1	7
3	Atherosclerotic disease in octogenarians: a challenge for science and clinical practice. <i>Atherosclerosis</i> , 2012 , 225, 281-9	3.1	28
2	The I405V and Taq1B polymorphisms of the CETP gene differentially affect sub-clinical carotid atherosclerosis. <i>Lipids in Health and Disease</i> , 2012 , 11, 130	4.4	4
1	Rebound inflammatory response during the acute phase of myocardial infarction after simvastatin withdrawal. <i>Atherosclerosis</i> , 2009 , 207, 191-4	3.1	52

