

# Niels de Fine Olivarius

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/701064/publications.pdf>

Version: 2024-02-01

87  
papers

2,699  
citations

218381

26  
h-index

197535

49  
g-index

89  
all docs

89  
docs citations

89  
times ranked

3838  
citing authors

#	ARTICLE	IF	CITATIONS
1	The impact of cancer on diabetes outcomes. BMC Endocrine Disorders, 2019, 19, 60.	0.9	8
2	Changes in Prescription Routines for Treating Hypothyroidism Between 2001 and 2015: An Observational Study of 929,684 Primary Care Patients in Copenhagen. Thyroid, 2019, 29, 910-919.	2.4	43
3	The effect of structured personal care on RNA oxidation: A 19-year follow-up of the randomized trial Diabetes Care in General Practice (DCGP). Journal of Diabetes and Its Complications, 2019, 33, 202-207.	1.2	3
4	Visual impairment and mortality in patients with type 2 diabetes. BMJ Open Diabetes Research and Care, 2019, 7, e000638.	1.2	8
5	The effect of structured personal care on diabetes symptoms and self-rated health over 14 years after diabetes diagnosis. Primary Care Diabetes, 2018, 12, 354-363.	0.9	4
6	Markers of DNA/RNA damage from oxidation as predictors of a registry-based diagnosis of psychiatric illness in type 2 diabetic patients. Psychiatry Research, 2018, 259, 370-376.	1.7	6
7	Multimorbidity and mortality. Journal of Comorbidity, 2018, 8, 2235042X1880406.	3.9	84
8	The development of multimorbidity during 16 years after diagnosis of type 2 diabetes. Journal of Comorbidity, 2018, 8, 2235042X1880165.	3.9	12
9	Brain Natriuretic Peptide in Plasma as Predictor of All-Cause Mortality in a Large Danish Primary Health Care Population Suspected of Heart Failure. Clinical Chemistry, 2018, 64, 1723-1731.	1.5	9
10	Interpretation of HbA <sub>1c</sub> in primary care and potential influence of anaemia and chronic kidney disease: an analysis from the Copenhagen Primary Care Laboratory (CopLab) Database. Diabetic Medicine, 2018, 35, 1700-1706.	1.2	19
11	Anemia is present years before myelodysplastic syndrome diagnosis: Results from the pre-diagnostic period. American Journal of Hematology, 2017, 92, E130-E132.	2.0	5
12	Urinary albumin and 8-oxo-7,8-dihydroguanosine as markers of mortality and cardiovascular disease during 19 years after diagnosis of type 2 diabetes – A comparative study of two markers to identify high risk patients. Redox Biology, 2017, 13, 363-369.	3.9	17
13	Socio-demographic determinants and effect of structured personal diabetes care: a 19-year follow-up of the randomized controlled study diabetes Care in General Practice (DCGP). BMC Endocrine Disorders, 2017, 17, 75.	0.9	0
14	The excess mortality of patients with diabetes and concurrent psychiatric illness is markedly reduced by structured personal diabetes care. General Hospital Psychiatry, 2016, 38, 42-52.	1.2	21
15	The role of diseases, risk factors and symptoms in the definition of multimorbidity – a systematic review. Scandinavian Journal of Primary Health Care, 2016, 34, 112-121.	0.6	144
16	The impact of gender on the long-term morbidity and mortality of patients with type 2 diabetes receiving structured personal care: a 13-year follow-up study. Diabetologia, 2016, 59, 275-285.	2.9	37
17	Intentional Weight Loss and Longevity in Overweight Patients with Type 2 Diabetes: A Population-Based Cohort Study. PLoS ONE, 2016, 11, e0146889.	1.1	32
18	Missing portion sizes in FFQ – alternatives to use of standard portions. Public Health Nutrition, 2015, 18, 1914-1921.	1.1	12

#	ARTICLE	IF	CITATIONS
19	Weight Changes following the Diagnosis of Type 2 Diabetes: The Impact of Recent and Past Weight History before Diagnosis. Results from the Danish Diabetes Care in General Practice (DCGP) Study. PLoS ONE, 2015, 10, e0122219.	1.1	20
20	Change in self-rated general health is associated with perceived illness burden: a 1-year follow up of patients newly diagnosed with type 2 diabetes. BMC Public Health, 2015, 15, 439.	1.2	7
21	Association of the blood eosinophil count with hematological malignancies and mortality. American Journal of Hematology, 2015, 90, 225-229.	2.0	20
22	Back on track—Smoking cessation and weight changes over 9 years in a community-based cohort study. Preventive Medicine, 2015, 81, 320-325.	1.6	4
23	Urinary markers of nucleic acid oxidation and cancer in type 2 diabetes. Redox Biology, 2015, 4, 34-39.	3.9	31
24	The impact of patients' involvement in cooking on their mortality and morbidity: A 19-year follow-up of patients diagnosed with type 2 diabetes mellitus. Scandinavian Journal of Primary Health Care, 2015, 33, 33-39.	0.6	6
25	Is thrombocytosis a valid indicator of advanced stage and high mortality of gynecological cancer?. Gynecologic Oncology, 2015, 139, 312-318.	0.6	10
26	Problems and challenges in relation to the treatment of patients with multimorbidity: General practitioners' views and attitudes. Scandinavian Journal of Primary Health Care, 2015, 33, 121-126.	0.6	52
27	The Copenhagen Primary Care Differential Count (CopDiff) database. Clinical Epidemiology, 2014, 6, 199.	1.5	16
28	Eosinophilia in routine blood samples as a biomarker for solid tumor development—A study based on The Copenhagen Primary Care Differential Count (CopDiff) Database. Acta Oncologica, 2014, 53, 1245-1250.	0.8	9
29	Motivation, effort and life circumstances as predictors of foot ulcers and amputations in people with Type 2 diabetes mellitus. Diabetic Medicine, 2014, 31, 1468-1476.	1.2	7
30	The effectiveness of structured personal care of type 2 diabetes on recurrent outcomes: a 19-year follow-up of the study Diabetes Care in General Practice (DCGP). Diabetologia, 2014, 57, 1119-23.	2.9	6
31	Risk of Lymphoma and Solid Cancer among Patients with Rheumatoid Arthritis in a Primary Care Setting. PLoS ONE, 2014, 9, e99388.	1.1	15
32	Amputations and foot ulcers in patients newly diagnosed with Type 2 diabetes mellitus and observed for 19 years. The role of age, gender and comorbidity. Diabetic Medicine, 2013, 30, 964-972.	1.2	83
33	Structured personal care of type 2 diabetes: a 19-year follow-up of the study Diabetes Care in General Practice (DCGP). Diabetologia, 2013, 56, 1243-1253.	2.9	41
34	Patients newly diagnosed with clinical type 2 diabetes mellitus but presenting with HbA1c within normal range: 19-Year mortality and clinical outcomes. Primary Care Diabetes, 2013, 7, 33-38.	0.9	3
35	Association Between Urinary Markers of Nucleic Acid Oxidation and Mortality in Type 2 Diabetes. Diabetes Care, 2013, 36, 669-676.	4.3	68
36	Eosinophilia in routine blood samples and the subsequent risk of hematological malignancies and death. American Journal of Hematology, 2013, 88, 843-847.	2.0	33

#	ARTICLE	IF	CITATIONS
37	FRIO113â€¦Rheumatoid arthritis and eosinophilia: the risk of lymphoproliferative malignancies and solid cancers. a study based on the copenhagen primary care differential count (copdiff) database. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A408.2-A408.	0.5	0
38	The astonishing hypothesis. <i>Primary Care Diabetes</i> , 2012, 6, 341-342.	0.9	0
39	Patients Newly Diagnosed with Clinical Type 2 Diabetes during Oral Glucocorticoid Treatment and Observed for 14 Years: Allâ€™Cause Mortality and Clinical Developments. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2011, 108, 285-288.	1.2	4
40	Prevalence and progression of visual impairment in patients newly diagnosed with clinical type 2 diabetes: a 6-year follow up study. <i>BMC Public Health</i> , 2011, 11, 80.	1.2	23
41	The Danish National Health Service Register. <i>Scandinavian Journal of Public Health</i> , 2011, 39, 34-37.	1.2	617
42	Urinary Markers of Nucleic Acid Oxidation and Long-Term Mortality of Newly Diagnosed Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2011, 34, 2594-2596.	4.3	92
43	Similar cardiovascular risk factor profile in screen-detected and known type 2 diabetic subjects. <i>Scandinavian Journal of Primary Health Care</i> , 2011, 29, 85-91.	0.6	4
44	The relationship between HbA<sub>1c</sub> level, symptoms and self-rated health in type 2 diabetic patients. <i>Scandinavian Journal of Primary Health Care</i> , 2011, 29, 157-164.	0.6	38
45	Fitness consultations in routine care of patients with type 2 diabetes in general practice: an 18-month non-randomised intervention study. <i>BMC Family Practice</i> , 2010, 11, 83.	2.9	14
46	Predictors of mortality of patients newly diagnosed with clinical type 2 diabetes: a 5-year follow up study. <i>BMC Endocrine Disorders</i> , 2010, 10, 14.	0.9	19
47	16-year excess all-cause mortality of newly diagnosed type 2 diabetic patients: a cohort study. <i>BMC Public Health</i> , 2009, 9, 400.	1.2	29
48	Changes in levels of haemoglobin A1cduring the first 6 years after diagnosis of clinical type 2 diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2009, 69, 851-857.	0.6	8
49	Randomised controlled trial of extraarticular gold bead implantation for treatment of knee osteoarthritis: a pilot study. <i>Clinical Rheumatology</i> , 2008, 27, 1363-1369.	1.0	16
50	Weight history of patients with newly diagnosed Type 2 diabetes. <i>Diabetic Medicine</i> , 2008, 25, 933-941.	1.2	15
51	Relationship of glucose concentrations with PAI-1 and t-PA in subjects with normal glucose tolerance. <i>Diabetic Medicine</i> , 2006, 23, 887-893.	1.2	10
52	Near-patient blood glucose measurements should only be used alone in the diagnosis of diabetes in cases of high glucose concentrations. <i>Diabetic Medicine</i> , 2006, 23, 1042-1042.	1.2	1
53	Changes in patient weight and the impact of antidiabetic therapy during the first 5Âˆyears after diagnosis of diabetes mellitus. <i>Diabetologia</i> , 2006, 49, 2058-2067.	2.9	28
54	Prevalence, severity and determinants of asthma in Danish five-year-olds. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2006, 95, 1182-1190.	0.7	17

#	ARTICLE	IF	CITATIONS
55	Urinary creatinine concentration is inversely related to glycaemic control and the presence of some diabetic complications in patients with newly diagnosed Type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2006, 20, 45-50.	1.2	9
56	General practitioners may diagnose type 2 diabetes mellitus at an early disease stage in patients they know well. <i>Family Practice</i> , 2006, 23, 192-197.	0.8	13
57	Structured personal diabetes care in primary health care affects only women's HbA1c. <i>Diabetes Care</i> , 2006, 29, 963-9.	4.3	11
58	Symptoms, signs and complications in newly diagnosed type 2 diabetic patients, and their relationship to glycaemia, blood pressure and weight. <i>Diabetologia</i> , 2005, 48, 210-214.	2.9	93
59	An educational model for improving diet counselling in primary care. <i>Patient Education and Counseling</i> , 2005, 58, 199-202.	1.0	10
60	General practitioners need to pay more attention to their poorly controlled type 2 diabetic patients. <i>European Journal of General Practice</i> , 2005, 11, 81-83.	0.9	0
61	Consequences Of Bias and Imprecision in Measurements of Glucose and Hba1c for the Diagnosis and Prognosis of Diabetes Mellitus. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2005, 65, 51-60.	0.6	36
62	Creation of a low-risk reference group and referenceinterval of fasting venous plasma glucose. <i>Clinical Chemistry and Laboratory Medicine</i> , 2004, 42, 817-23.	1.4	2
63	Diabetes care today: not everyone should have intensive multipharmacological treatment. <i>Scandinavian Journal of Primary Health Care</i> , 2004, 22, 67-70.	0.6	9
64	Individualised treatment goals in diabetes care. <i>Scandinavian Journal of Primary Health Care</i> , 2004, 22, 71-77.	0.6	9
65	Menstrual bleeding patterns in pre- and perimenopausal women: a population-based prospective diary study. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2004, 83, 197-202.	1.3	17
66	Peripheral vascular disease is associated with reduced glycosuria in newly diagnosed type 2 diabetic patients. <i>Diabetes and Metabolism</i> , 2004, 30, 269-274.	1.4	4
67	Menstrual bleeding patterns in pre- and perimenopausal women: a population-based prospective diary study. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2004, 83, 197-202.	1.3	9
68	Encouraging structured personalised diabetes care in general practice. <i>Scandinavian Journal of Primary Health Care</i> , 2003, 21, 89-95.	0.6	14
69	The Effect of the New ADA and WHO Guidelines on the Number of Diagnosed Cases of Diabetes Mellitus. <i>Clinical Chemistry and Laboratory Medicine</i> , 2003, 41, 1246-50.	1.4	13
70	Can capillary whole blood glucose and venous plasma glucose measurements be used interchangeably in diagnosis of diabetes mellitus?. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2002, 62, 159-166.	0.6	46
71	Upper reference limit, analytical quality specifications and clinical use of haemoglobin A1C. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2002, 62, 609-622.	0.6	26
72	Can general practitioners be randomised?. <i>Scandinavian Journal of Primary Health Care</i> , 2002, 20, 23-24.	0.6	0

#	ARTICLE	IF	CITATIONS
73	Optimization of preanalytical conditions and analysis of plasma glucose. 1. Impact of the new WHO and ADA recommendations on diagnosis of diabetes mellitus. Scandinavian Journal of Clinical and Laboratory Investigation, 2001, 61, 169-179.	0.6	65
74	Randomised controlled trial of structured personal care of type 2 diabetes mellitus. BMJ: British Medical Journal, 2001, 323, 970-970.	2.4	252
75	Diabetic retinopathy in newly diagnosed middle-aged and elderly diabetic patients. Prevalence and interrelationship with microalbuminuria and triglycerides. , 2001, 239, 664-672.		26
76	Evaluation of systematic and random factors in measurements of fasting plasma glucose as the basis for analytical quality specifications in the diagnosis of diabetes. 3. Impact of the new WHO and ADA recommendations on diagnosis of diabetes mellitus. Scandinavian Journal of Clinical and Laboratory Investigation, 2001, 61, 191-204.	0.6	44
77	Plasma glucose reference interval in a low-risk population. 2. Impact of the new WHO and ADA recommendations on the diagnosis of diabetes mellitus. Scandinavian Journal of Clinical and Laboratory Investigation, 2001, 61, 181-190.	0.6	32
78	Encouraging GPs to undertake screening and a brief intervention in order to reduce problem drinking: a randomized controlled trial. Family Practice, 1999, 16, 551-557.	0.8	22
79	The UK Prospective Diabetes study. Lancet, The, 1998, 352, 1933.	6.3	2
80	Accuracy of 1-, 5- and 10-year body weight recall given in a standard questionnaire. International Journal of Obesity, 1997, 21, 67-71.	1.6	44
81	Five-year all-cause mortality of 1323 newly diagnosed middle-aged and elderly diabetic patients. Journal of Diabetes and Its Complications, 1997, 11, 83-89.	1.2	17
82	Association between risk factors and overnight urinary albumin/creatinine ratio " Even in its normal range. Journal of Diabetes and Its Complications, 1994, 8, 178-179.	1.2	0
83	Self-referral and self-payment in Danish primary care. Health Policy, 1994, 28, 15-22.	1.4	8
84	Longitudinal Research in General Practice. Scandinavian Journal of Primary Health Care, 1993, 11, 35-35.	0.6	0
85	6.1.1.1 Discrepancy in HbA <sub>1c</sub> Measurements Performed at Different Local Laboratories and at a Selected Central Reference Laboratory. Upsala Journal of Medical Sciences, 1993, 98, 275-282.	0.4	3
86	The Danish Study Diabetes Care in General Practice. Scandinavian Journal of Primary Health Care, 1993, 11, 49-53.	0.6	2
87	The Danish Study Diabetes Care in General Practice. Management of main methodological and practical problems in a long-term multipractice intervention study. Scandinavian Journal of Primary Health Care, Supplement, 1993, 2, 49-53.	0.1	3