

Nicola Martinelli

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/701135/nicola-martinelli-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140
papers

11,318
citations

44
h-index

106
g-index

149
ext. papers

12,898
ext. citations

7.9
avg, IF

5.14
L-index

#	Paper	IF	Citations
140	High Plasma Concentration of Apolipoprotein C-III Confers an Increased Risk of Cerebral Ischemic Events on Cardiovascular Patients Anticoagulated With Warfarin.. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 781383	5.4	
139	Identification of a Novel Serological Marker in Seronegative Rheumatoid Arthritis Using the Peptide Library Approach. <i>Frontiers in Immunology</i> , 2021 , 12, 753400	8.4	1
138	Basophil Blood Cell Count Is Associated With Enhanced Factor II Plasma Coagulant Activity and Increased Risk of Mortality in Patients With Stable Coronary Artery Disease: Not Only Neutrophils as Prognostic Marker in Ischemic Heart Disease. <i>Journal of the American Heart Association</i> , 2021 , 10, e018243	6	5
137	Whole-genome sequencing analysis of semi-supercentenarians. <i>ELife</i> , 2021 , 10,	8.9	11
136	rs629301 CELSR2 polymorphism confers a ten-year equivalent risk of critical stenosis assessed by coronary angiography. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 1542-1547	4.5	0
135	Serum Uric Acid Levels, but Not rs7442295 Polymorphism of SCL2A9 Gene, Predict Mortality in Clinically Stable Coronary Artery Disease. <i>Current Problems in Cardiology</i> , 2021 , 46, 100798	17.1	0
134	A dangerous onychodystrophy. <i>American Journal of Hematology</i> , 2021 , 96, 891-892	7.1	
133	The Asialoglycoprotein Receptor Minor Subunit Gene Contributes to Pharmacokinetics of Factor VIII Concentrates in Hemophilia A. <i>Thrombosis and Haemostasis</i> , 2021 ,	7	2
132	Detection of Urinary Exosomal HSD11B2 mRNA Expression: A Useful Novel Tool for the Diagnostic Approach of Dysfunctional 11βHSD2-Related Hypertension. <i>Frontiers in Endocrinology</i> , 2021 , 12, 681974	5.7	2
131	Portal hypertensive biliopathy and bile duct varices presenting as jaundice. <i>Endoscopy</i> , 2021 , 53, E442-E443	5.4	
130	Rare variants lowering the levels of coagulation factor X are protective against ischemic heart disease. <i>Haematologica</i> , 2020 , 105, e365-e369	6.6	7
129	Aptamer-modified FXa generation assays to investigate hypercoagulability in plasma from patients with ischemic heart disease. <i>Thrombosis Research</i> , 2020 , 189, 140-146	8.2	3
128	Increased Incidence of Ischemic Cerebrovascular Events in Cardiovascular Patients With Elevated Apolipoprotein CIII. <i>Stroke</i> , 2020 , 51, 61-68	6.7	3
127	The Positive Association between Plasma Myristic Acid and ApoCIII Concentrations in Cardiovascular Disease Patients Is Supported by the Effects of Myristic Acid in HepG2 Cells. <i>Journal of Nutrition</i> , 2020 , 150, 2707-2715	4.1	7
126	A relative ADAMTS13 deficiency supports the presence of a secondary microangiopathy in COVID 19. <i>Thrombosis Research</i> , 2020 , 193, 170-172	8.2	36
125	Deep vein thrombosis in SARS-CoV-2 pneumonia-affected patients within standard care units: Exploring a submerged portion of the iceberg. <i>Thrombosis Research</i> , 2020 , 194, 216-219	8.2	10
124	Trace Elements Status and Metallothioneins DNA Methylation Influence Human Hepatocellular Carcinoma Survival Rate. <i>Frontiers in Oncology</i> , 2020 , 10, 596040	5.3	

123	Functional polymorphisms in the LDLR and pharmacokinetics of Factor VIII concentrates. <i>Journal of Thrombosis and Haemostasis</i> , 2019 , 17, 1288-1296	15.4	8
122	Subsequent Event Risk in Individuals With Established Coronary Heart Disease. <i>Circulation Genomic and Precision Medicine</i> , 2019 , 12, e002470	5.2	13
121	Association of Chromosome 9p21 With Subsequent Coronary Heart Disease Events. <i>Circulation Genomic and Precision Medicine</i> , 2019 , 12, e002471	5.2	14
120	Plasma Proteome Profiles of Stable CAD Patients Stratified According to Total Apo C-III Levels. <i>Proteomics - Clinical Applications</i> , 2019 , 13, e1800023	3.1	3
119	Apolipoprotein C-III Strongly Correlates with Activated Factor VII-Anti-Thrombin Complex: An Additional Link between Plasma Lipids and Coagulation. <i>Thrombosis and Haemostasis</i> , 2019 , 119, 192-207	7	11
118	Not Just Arterial Damage: Increased Incidence of Venous Thromboembolic Events in Cardiovascular Patients With Elevated Plasma Levels of Apolipoprotein CIII. <i>Journal of the American Heart Association</i> , 2019 , 8, e010973	6	8
117	Factor H interferes with the adhesion of sickle red cells to vascular endothelium: a novel disease-modulating molecule. <i>Haematologica</i> , 2019 , 104, 919-928	6.6	22
116	Therapeutic oligonucleotides in cardiovascular and metabolic diseases: insights for the internist. <i>Internal and Emergency Medicine</i> , 2018 , 13, 313-318	3.7	3
115	Myristic acid induces proteomic and secretomic changes associated with steatosis, cytoskeleton remodeling, endoplasmic reticulum stress, protein turnover and exosome release in HepG2 cells. <i>Journal of Proteomics</i> , 2018 , 181, 118-130	3.9	17
114	Sialylated isoforms of apolipoprotein C-III and plasma lipids in subjects with coronary artery disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018 , 56, 1542-1550	5.9	6
113	Improvement of maternal and fetal outcomes in women with sickle cell disease treated with early prophylactic erythrocytapheresis. <i>Transfusion</i> , 2018 , 58, 2192-2201	2.9	14
112	Systematic Evaluation of Pleiotropy Identifies 6 Further Loci Associated With 'Coronary Artery' Disease. <i>Journal of the American College of Cardiology</i> , 2017 , 69, 823-836	15.1	146
111	A decade of progress on the genetic basis of coronary artery disease. Practical insights for the internist. <i>European Journal of Internal Medicine</i> , 2017 , 41, 10-17	3.9	12
110	Increased plasma thrombin potential is associated with stable coronary artery disease: An angiographically-controlled study. <i>Thrombosis Research</i> , 2017 , 155, 16-22	8.2	10
109	Relations between lipoprotein(a) concentrations, LPA genetic variants, and the risk of mortality in patients with established coronary heart disease: a molecular and genetic association study. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 534-543	18.1	69
108	Impact of Selection Bias on Estimation of Subsequent Event Risk. <i>Circulation: Cardiovascular Genetics</i> , 2017 , 10,		19
107	One-carbon genetic variants and the role of MTHFD1 1958G>A in liver and colon cancer risk according to global DNA methylation. <i>PLoS ONE</i> , 2017 , 12, e0185792	3.7	11
106	The RFC1 80G>A, among Common One-Carbon Polymorphisms, Relates to Survival Rate According to DNA Global Methylation in Primary Liver Cancers. <i>PLoS ONE</i> , 2016 , 11, e0167534	3.7	4

105	Activated factor VII-antithrombin complex predicts mortality in patients with stable coronary artery disease: a cohort study. <i>Journal of Thrombosis and Haemostasis</i> , 2016 , 14, 655-66	15.4	15
104	Coding Variation in ANGPTL4, LPL, and SVEP1 and the Risk of Coronary Disease. <i>New England Journal of Medicine</i> , 2016 , 374, 1134-44	59.2	325
103	Infections and autoimmunity: role of human cytomegalovirus in autoimmune endothelial cell damage. <i>Lupus</i> , 2015 , 24, 419-32	2.6	18
102	Exome sequencing identifies rare LDLR and APOA5 alleles conferring risk for myocardial infarction. <i>Nature</i> , 2015 , 518, 102-6	50.4	463
101	An integrated genomic-transcriptomic approach supports a role for the proto-oncogene BCL3 in atherosclerosis. <i>Thrombosis and Haemostasis</i> , 2015 , 113, 655-63	7	8
100	The MTHFD1 1958G>A Relates to Survival Rate According to PBMCs DNA Global Methylation in Cancer. <i>FASEB Journal</i> , 2015 , 29, 749.3	0.9	
99	Association of low-frequency and rare coding-sequence variants with blood lipids and coronary heart disease in 56,000 whites and blacks. <i>American Journal of Human Genetics</i> , 2014 , 94, 223-32	11	233
98	A novel molecular diagnostic marker for familial and early-onset coronary artery disease and myocardial infarction in the LRP8 gene. <i>Circulation: Cardiovascular Genetics</i> , 2014 , 7, 514-20		17
97	Delta-5 and delta-6 desaturases: crucial enzymes in polyunsaturated fatty acid-related pathways with pleiotropic influences in health and disease. <i>Advances in Experimental Medicine and Biology</i> , 2014 , 824, 61-81	3.6	92
96	Loss-of-function mutations in APOC3, triglycerides, and coronary disease. <i>New England Journal of Medicine</i> , 2014 , 371, 22-31	59.2	721
95	Iron deficiency in the elderly population, revisited in the hepcidin era. <i>Frontiers in Pharmacology</i> , 2014 , 5, 83	5.6	70
94	Multi-allelic haplotype association identifies novel information different from single-SNP analysis: a new protective haplotype in the LRP8 gene is against familial and early-onset CAD and MI. <i>Gene</i> , 2013 , 521, 78-81	3.8	12
93	Air particulate matter and cardiovascular disease: a narrative review. <i>European Journal of Internal Medicine</i> , 2013 , 24, 295-302	3.9	178
92	Paraoxonases: ancient substrate hunters and their evolving role in ischemic heart disease. <i>Advances in Clinical Chemistry</i> , 2013 , 59, 65-100	5.8	22
91	Paraoxonase-1 status in patients with hereditary hemochromatosis. <i>Journal of Lipid Research</i> , 2013 , 54, 1484-92	6.3	19
90	Global DNA hypomethylation in peripheral blood mononuclear cells as a biomarker of cancer risk. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2013 , 22, 348-55	4	55
89	Factor II activity is similarly increased in patients with elevated apolipoprotein CIII and in carriers of the factor II 20210A allele. <i>Journal of the American Heart Association</i> , 2013 , 2, e000440	6	21
88	Exome sequencing and directed clinical phenotyping diagnose cholesterol ester storage disease presenting as autosomal recessive hypercholesterolemia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 2909-14	9.4	76

87	Global DNA hypomethylation in peripheral blood mononuclear cells as a biomarker of cancer risk. <i>FASEB Journal</i> , 2013 , 27, 248.1	0.9	
86	Iron Status Independently Associates With Bone Mineral Density At Population Level. Insights From The Val Borbera Study. <i>Blood</i> , 2013 , 122, 4672-4672	2.2	
85	Activated Factor VII Antithrombin Complex Plasma Concentration Is An Independent Predictor Of Total and Cardiovascular Mortality In Patients With Coronary Artery Disease and Its Prognostic Significance Is Improved By Using Factor VII Genotype-Specific Threshold Levels. <i>Blood</i> , 2013 , 122, 2339-2339	2.2	
84	Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. <i>Lancet, The</i> , 2012 , 380, 572-80	40	1523
83	Serum levels of the hepcidin-20 isoform in a large general population: the Val Borbera study. <i>Journal of Proteomics</i> , 2012 , 76 Spec No., 28-35	3.9	27
82	Promoter methylation in coagulation F7 gene influences plasma FVII concentrations and relates to coronary artery disease. <i>Journal of Medical Genetics</i> , 2012 , 49, 192-9	5.8	50
81	Increased serum hepcidin levels in subjects with the metabolic syndrome: a population study. <i>PLoS ONE</i> , 2012 , 7, e48250	3.7	53
80	Low levels of serum paraoxonase activities are characteristic of metabolic syndrome and may influence the metabolic-syndrome-related risk of coronary artery disease. <i>Experimental Diabetes Research</i> , 2012 , 2012, 231502		19
79	DISHphagia: an unusual cause of dysphagia. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 2573-4	5.6	1
78	Access rate to the emergency department for venous thromboembolism in relationship with coarse and fine particulate matter air pollution. <i>PLoS ONE</i> , 2012 , 7, e34831	3.7	38
77	Identification of ADAMTS7 as a novel locus for coronary atherosclerosis and association of ABO with myocardial infarction in the presence of coronary atherosclerosis: two genome-wide association studies. <i>Lancet, The</i> , 2011 , 377, 383-92	40	399
76	Polymorphisms of cyclo-oxygenases and 5-lipo-oxygenase-activating protein are associated with chronic spontaneous urticaria and urinary leukotriene E4. <i>European Journal of Dermatology</i> , 2011 , 21, 47-52	0.8	8
75	Hepcidin levels and their determinants in different types of myelodysplastic syndromes. <i>PLoS ONE</i> , 2011 , 6, e23109	3.7	81
74	Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. <i>Nature Genetics</i> , 2011 , 43, 333-8	36.3	1394
73	Large-scale gene-centric analysis identifies novel variants for coronary artery disease. <i>PLoS Genetics</i> , 2011 , 7, e1002260	6	175
72	Abdominal aortic aneurysm is associated with a variant in low-density lipoprotein receptor-related protein 1. <i>American Journal of Human Genetics</i> , 2011 , 89, 619-27	11	145
71	Folic acid effects on s-adenosylmethionine, s-adenosylhomocysteine, and DNA methylation in patients with intermediate hyperhomocysteinemia. <i>Journal of the American College of Nutrition</i> , 2011 , 30, 11-8	3.5	36
70	Differences and similarities between allergic and nonallergic rhinitis in a large sample of adult patients with rhinitis symptoms. <i>International Archives of Allergy and Immunology</i> , 2011 , 155, 263-70	3.7	56

69	An unusual heart failure: cardiac amyloidosis due to light-chain myeloma. <i>Circulation</i> , 2011 , 123, e583-4	16.7	3
68	Serum Hepcidin Levels Correlate with Phenotypes of the Metabolic Syndrome At Population Level. <i>Blood</i> , 2011 , 118, 348-348	2.2	1
67	Genome-wide association study identifies a sequence variant within the DAB2IP gene conferring susceptibility to abdominal aortic aneurysm. <i>Nature Genetics</i> , 2010 , 42, 692-7	36.3	155
66	Evaluation of hepcidin isoforms in hemodialysis patients by a proteomic approach based on SELDI-TOF MS. <i>Journal of Biomedicine and Biotechnology</i> , 2010 , 2010, 329646		30
65	Lack of association between the Trp719Arg polymorphism in kinesin-like protein-6 and coronary artery disease in 19 case-control studies. <i>Journal of the American College of Cardiology</i> , 2010 , 56, 1552-63	15.1	75
64	Detection of a large deletion in the P-selectin (SELP) gene. <i>Molecular and Cellular Probes</i> , 2010 , 24, 161-5	3	6
63	Polymorphisms at LDLR locus may be associated with coronary artery disease through modulation of coagulation factor VIII activity and independently from lipid profile. <i>Blood</i> , 2010 , 116, 5688-97	2.2	64
62	Apolipoprotein C-III predicts cardiovascular mortality in severe coronary artery disease and is associated with an enhanced plasma thrombin generation. <i>Journal of Thrombosis and Haemostasis</i> , 2010 , 8, 463-71	15.4	39
61	Hepcidin Levels and Their Determinants In Different Types of Myelodysplastic Syndromes. <i>Blood</i> , 2010 , 116, 4250-4250	2.2	
60	Serum DNase I, soluble Fas/FasL levels and cell surface Fas expression in patients with SLE: a possible explanation for the lack of efficacy of hrDNase I treatment. <i>International Immunology</i> , 2009 , 21, 237-43	4.9	26
59	Association and functional analyses of MEF2A as a susceptibility gene for premature myocardial infarction and coronary artery disease. <i>Circulation: Cardiovascular Genetics</i> , 2009 , 2, 165-72		25
58	Genetic architecture of coronary artery disease in the genome-wide era: implications for the emerging "golden dozen" loci. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 671-82	5.3	29
57	A score of risk factors associated with ischemic digital ulcers in patients affected by systemic sclerosis treated with iloprost. <i>Clinical Rheumatology</i> , 2009 , 28, 807-13	3.9	30
56	Additive effect of LRP8/APOER2 R952Q variant to APOE epsilon2/epsilon3/epsilon4 genotype in modulating apolipoprotein E concentration and the risk of myocardial infarction: a case-control study. <i>BMC Medical Genetics</i> , 2009 , 10, 41	2.1	20
55	Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. <i>Nature Genetics</i> , 2009 , 41, 342-7	36.3	627
54	Genome-wide association of early-onset myocardial infarction with single nucleotide polymorphisms and copy number variants. <i>Nature Genetics</i> , 2009 , 41, 334-41	36.3	884
53	Reduced serum hepcidin levels in patients with chronic hepatitis C. <i>Journal of Hepatology</i> , 2009 , 51, 845-53	5.4	123
52	Evaluation of serum s-IgE/total IgE ratio in predicting clinical response to allergen-specific immunotherapy. <i>Journal of Allergy and Clinical Immunology</i> , 2009 , 123, 1103-10, 1110.e1-4	11.5	119

51	Novel serum paraoxonase activity assays are associated with coronary artery disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 432-40	5.9	26
50	A Desaturase hypothesis for atherosclerosis: Janus-faced enzymes in omega-6 and omega-3 polyunsaturated fatty acid metabolism. <i>Journal of Nutrigenetics and Nutrigenomics</i> , 2009 , 2, 129-39		19
49	Clinical importance of eosinophil count in nasal fluid in patients with allergic and non-allergic rhinitis. <i>International Journal of Immunopathology and Pharmacology</i> , 2009 , 22, 1077-87	3	9
48	FADS genotypes and desaturase activity estimated by the ratio of arachidonic acid to linoleic acid are associated with inflammation and coronary artery disease. <i>American Journal of Clinical Nutrition</i> , 2008 , 88, 941-9	7	241
47	Combined effect of hemostatic gene polymorphisms and the risk of myocardial infarction in patients with advanced coronary atherosclerosis. <i>PLoS ONE</i> , 2008 , 3, e1523	3.7	28
46	SNPs of the FADS gene cluster are associated with polyunsaturated fatty acids in a cohort of patients with cardiovascular disease. <i>Lipids</i> , 2008 , 43, 289-99	1.6	192
45	Association between four SNPs on chromosome 9p21 and myocardial infarction is replicated in an Italian population. <i>Journal of Human Genetics</i> , 2008 , 53, 144-150	4.3	102
44	Comparative effect of tacrolimus 0.1% ointment and clobetasol 0.05% ointment in patients with oral lichen planus. <i>Journal of Clinical Periodontology</i> , 2008 , 35, 244-9	7.7	67
43	Gene sequence variations of the platelet P2Y12 receptor are associated with coronary artery disease. <i>BMC Medical Genetics</i> , 2007 , 8, 59	2.1	46
42	ALOX5AP gene variants and risk of coronary artery disease: an angiography-based study. <i>European Journal of Human Genetics</i> , 2007 , 15, 959-66	5.3	35
41	Two new highly polymorphic markers in the 3QTR region of the PLA2G7 gene. <i>International Journal of Immunogenetics</i> , 2007 , 34, 465-8	2.3	
40	ApoE epsilon2/epsilon3/epsilon4 polymorphism, ApoC-III/ApoE ratio and metabolic syndrome. <i>Clinical and Experimental Medicine</i> , 2007 , 7, 164-72	4.9	37
39	Altered renal folate handling in hypertensive patients with nephroangiosclerotic damage. <i>Journal of Human Hypertension</i> , 2007 , 21, 327-9	2.6	3
38	Scleroderma patients nailfold videocapillaroscopic patterns are associated with disease subset and disease severity. <i>Rheumatology</i> , 2007 , 46, 1566-9	3.9	91
37	Increased factor VIII coagulant activity levels in male carriers of the factor V R2 polymorphism. <i>Blood Coagulation and Fibrinolysis</i> , 2007 , 18, 125-9	1	7
36	The -1131 T>C and S19W APOA5 gene polymorphisms are associated with high levels of triglycerides and apolipoprotein C-III, but not with coronary artery disease: an angiographic study. <i>Atherosclerosis</i> , 2007 , 191, 409-17	3.1	58
35	Anti-TNFalpha therapy in rheumatoid arthritis and autoimmunity. <i>Rheumatology International</i> , 2006 , 26, 209-14	3.6	60
34	Evaluation of finger skin temperature in scleroderma patients cyclically treated with iloprost. <i>Joint Bone Spine</i> , 2006 , 73, 57-61	2.9	4

33	Relationship between human leucocyte antigen class I and class II and chronic idiopathic urticaria associated with aspirin and/or NSAIDs hypersensitivity. <i>Mediators of Inflammation</i> , 2006 , 2006, 62489	4.3	15
32	Comparison of topical tacrolimus 0.1 % in pectin ointment with clobetasol 0.5% ointment in adults with moderate to severe desquamative gingivitis: A 4-week, randomized, double-blind clinical trial. <i>Clinical Therapeutics</i> , 2006 , 28, 1296-302	3.5	15
31	Hyperhomocysteinemia and mortality after coronary artery bypass grafting. <i>PLoS ONE</i> , 2006 , 1, e83	3.7	15
30	Tacrolimus ointment in nickel sulphate-induced steroid-resistant allergic contact dermatitis. <i>Allergy and Asthma Proceedings</i> , 2006 , 27, 527-31	2.6	17
29	Infective endocarditis with lung and systemic embolization in an injection drug user. <i>European Heart Journal</i> , 2006 , 27, 2938	9.5	
28	Homocysteine, traditional risk factors and impaired renal function in coronary artery disease. <i>European Journal of Clinical Investigation</i> , 2006 , 36, 698-704	4.6	6
27	On the association of the oxidised LDL receptor 1 (OLR1) gene in patients with acute myocardial infarction or coronary artery disease. <i>European Journal of Human Genetics</i> , 2006 , 14, 127-30	5.3	38
26	Reply to Novelli. <i>European Journal of Human Genetics</i> , 2006 , 14, 895-895	5.3	9
25	Évaluation de la température cutanée des patients atteints de sclérodermie traitée périodiquement par iloprost. <i>Revue Du Rhumatisme (Edition Francaise)</i> , 2006 , 73, 53-57	0.1	1
24	ALOX5AP Gene Variants and Risk of Coronary Artery Disease in Italy. An Angiography-Based Study.. <i>Blood</i> , 2006 , 108, 1459-1459	2.2	
23	Long-term evaluation of lung function in patients affected by scleroderma treated with cyclic iloprost infusions. <i>Rheumatology International</i> , 2005 , 25, 250-4	3.6	4
22	Antibodies against cyclic citrullinated peptides in patients affected by rheumatoid arthritis before and after infliximab treatment. <i>Rheumatology International</i> , 2005 , 26, 58-62	3.6	45
21	Interaction between metabolic syndrome and PON1 polymorphisms as a determinant of the risk of coronary artery disease. <i>Clinical and Experimental Medicine</i> , 2005 , 5, 20-30	4.9	19
20	Reply to J Dierkes et al. <i>American Journal of Clinical Nutrition</i> , 2005 , 81, 727-728	7	6
19	Prevalence of body iron excess in the metabolic syndrome. <i>Diabetes Care</i> , 2005 , 28, 2061-3	14.6	146
18	Apolipoprotein C-III, n-3 polyunsaturated fatty acids, and "insulin-resistant" T-455C APOC3 gene polymorphism in heart disease patients: example of gene-diet interaction. <i>Clinical Chemistry</i> , 2005 , 51, 360-7	5.5	44
17	Food-additive-induced urticaria: a survey of 838 patients with recurrent chronic idiopathic urticaria. <i>International Archives of Allergy and Immunology</i> , 2005 , 138, 235-42	3.7	56
16	Hyperhomocysteinemia in Relation to Total and Cardiovascular Death after Coronary Artery Bypass Grafting. A Prospective Study.. <i>Blood</i> , 2005 , 106, 1640-1640	2.2	

15	Influence of polymorphisms in the factor VII gene promoter on activated factor VII levels and on the risk of myocardial infarction in advanced coronary atherosclerosis. <i>Thrombosis and Haemostasis</i> , 2004 , 92, 541-9	7	36
14	Low plasma vitamin B-6 concentrations and modulation of coronary artery disease risk. <i>American Journal of Clinical Nutrition</i> , 2004 , 79, 992-8	7	94
13	Modulation of factor V levels in plasma by polymorphisms in the C2 domain. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2004 , 24, 200-6	9.4	23
12	Randomized placebo-controlled trial comparing fluticasone aqueous nasal spray in mono-therapy, fluticasone plus cetirizine, fluticasone plus montelukast and cetirizine plus montelukast for seasonal allergic rhinitis. <i>Clinical and Experimental Allergy</i> , 2004 , 34, 259-67	4.1	136
11	Comparing tacrolimus ointment and oral cyclosporine in adult patients affected by atopic dermatitis: a randomized study. <i>Clinical and Experimental Allergy</i> , 2004 , 34, 639-45	4.1	52
10	Randomized placebo-controlled trial comparing fluticasone aqueous nasal spray in mono-therapy, fluticasone plus cetirizine, fluticasone plus montelukast and cetirizine plus montelukast for seasonal allergic rhinitis. <i>Clinical and Experimental Allergy</i> , 2004 , 34, 1329-1329	4.1	3
9	Interaction between smoking and PON2 Ser311Cys polymorphism as a determinant of the risk of myocardial infarction. <i>European Journal of Clinical Investigation</i> , 2004 , 34, 14-20	4.6	42
8	Monosodium benzoate hypersensitivity in subjects with persistent rhinitis. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2004 , 59, 192-7	9.3	37
7	Randomized placebo-controlled trial comparing desloratadine and montelukast in monotherapy and desloratadine plus montelukast in combined therapy for chronic idiopathic urticaria. <i>Journal of Allergy and Clinical Immunology</i> , 2004 , 114, 619-25	11.5	103
6	Apolipoprotein C-III, metabolic syndrome, and risk of coronary artery disease. <i>Journal of Lipid Research</i> , 2003 , 44, 2374-81	6.3	98
5	Results of double-blind placebo-controlled challenge with nickel salts in patients affected by recurrent aphthous stomatitis. <i>International Archives of Allergy and Immunology</i> , 2003 , 131, 296-300	3.7	8
4	Serum levels of soluble CD30 in adult patients affected by atopic dermatitis and its relation to age, duration of disease and Scoring Atopic Dermatitis index. <i>Mediators of Inflammation</i> , 2003 , 12, 123-5	4.3	11
3	Nickel sulphate, food additives and seronegative arthritis: is there any relationship?. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2003 , 58, 958-9	9.3	1
2	Interaction of antibodies against cytomegalovirus with heat-shock protein 60 in pathogenesis of atherosclerosis. <i>Lancet, The</i> , 2003 , 362, 1971-7	4.0	80
1	A1298C methylenetetrahydrofolate reductase mutation and coronary artery disease: relationships with C677T polymorphism and homocysteine/folate metabolism. <i>Clinical and Experimental Medicine</i> , 2002 , 2, 7-12	4.9	42