
Shannon C Kenney

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7015778/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | A New Model of Epstein-Barr Virus Infection Reveals an Important Role for Early Lytic Viral Protein Expression in the Development of Lymphomas. Journal of Virology, 2011, 85, 165-177. | 3.4 | 239 |
| 2 | Reactivation of Latent Epstein-Barr Virus by Methotrexate: A Potential Contributor to Methotrexate-Associated Lymphomas. Journal of the National Cancer Institute, 2004, 96, 1691-1702. | 6.3 | 228 |
| 3 | Regulation of the latent-lytic switch in Epstein–Barr virus. Seminars in Cancer Biology, 2014, 26, 60-68. | 9.6 | 219 |
| 4 | Epstein-Barr Virus Immediate-Early Protein BZLF1 Is SUMO-1 Modified and Disrupts Promyelocytic Leukemia Bodies. Journal of Virology, 2001, 75, 2388-2399. | 3.4 | 213 |
| 5 | Lytic Induction Therapy for Epstein-Barr Virus-Positive B-Cell Lymphomas. Journal of Virology, 2004, 78, 1893-1902. | 3.4 | 200 |
| 6 | Epstein-Barr Virus Lytic Infection Contributes to Lymphoproliferative Disease in a SCID Mouse Model. Journal of Virology, 2005, 79, 13993-14003. | 3.4 | 198 |
| 7 | The EBV lytic switch protein, Z, preferentially binds to and activates the methylated viral genome. Nature Genetics, 2004, 36, 1099-1104. | 21.4 | 170 |
| 8 | Epstein-Barr Virus Immediate-Early Proteins BZLF1 and BRLF1 Activate the ATF2 Transcription Factor by Increasing the Levels of Phosphorylated p38 and c-Jun N-Terminal Kinases. Journal of Virology, 2000, 74, 1224-1233. | 3.4 | 161 |
| 9 | Direct BRLF1 binding is required for cooperative BZLF1/BRLF1 activation of the Epstein-Barr virus early promoter, BMRF1. Nucleic Acids Research, 1993, 21, 1999-2007. | 14.5 | 141 |
| 10 | Chemotherapy induces lytic EBV replication and confers ganciclovir susceptibility to EBV-positive epithelial cell tumors. Cancer Research, 2002, 62, 1920-6. | 0.9 | 133 |
| 11 | X-Box-Binding Protein 1 Activates Lytic Epstein-Barr Virus Gene Expression in Combination with Protein Kinase D. Journal of Virology, 2007, 81, 7363-7370. | 3.4 | 105 |
| 12 | An Epstein-Barr Virus (EBV) Mutant with Enhanced BZLF1 Expression Causes Lymphomas with Abortive Lytic EBV Infection in a Humanized Mouse Model. Journal of Virology, 2012, 86, 7976-7987. | 3.4 | 102 |
| 13 | Epstein-Barr Virus Lytic Infection Is Required for Efficient Production of the Angiogenesis Factor Vascular Endothelial Growth Factor in Lymphoblastoid Cell Lines. Journal of Virology, 2005, 79, 13984-13992. | 3.4 | 93 |
| 14 | PD-1/CTLA-4 Blockade Inhibits Epstein-Barr Virus-Induced Lymphoma Growth in a Cord Blood Humanized-Mouse Model. PLoS Pathogens, 2016, 12, e1005642. | 4.7 | 87 |
| 15 | Differentiation-Dependent KLF4 Expression Promotes Lytic Epstein-Barr Virus Infection in Epithelial Cells. PLoS Pathogens, 2015, 11, e1005195. | 4.7 | 79 |
| 16 | BZLF1 Activation of the Methylated Form of the BRLF1 Immediate-Early Promoter Is Regulated by BZLF1 Residue 186. Journal of Virology, 2005, 79, 7338-7348. | 3.4 | 75 |
| 17 | Cellular Differentiation Regulator BLIMP1 Induces Epstein-Barr Virus Lytic Reactivation in Epithelial and B Cells by Activating Transcription from both the R and Z Promoters. Journal of Virology, 2015, 89, 1731-1743. | 3.4 | 75 |
| 18 | The Cellular Ataxia Telangiectasia-Mutated Kinase Promotes Epstein-Barr Virus Lytic Reactivation in Response to Multiple Different Types of Lytic Reactivation-Inducing Stimuli. Journal of Virology, 2012, 86, 13360-13370. | 3.4 | 71 |

SHANNON C KENNEY

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Methylation-Dependent Binding of the Epstein-Barr Virus BZLF1 Protein to Viral Promoters. PLoS Pathogens, 2009, 5, e1000356. | 4.7 | 70 |
| 20 | A cancer-associated Epstein-Barr virus BZLF1 promoter variant enhances lytic infection. PLoS Pathogens, 2018, 14, e1007179. | 4.7 | 68 |
| 21 | Leflunomide/teriflunomide inhibit Epstein-Barr virus (EBV)-induced lymphoproliferative disease and lytic viral replication. Oncotarget, 2017, 8, 44266-44280. | 1.8 | 61 |
| 22 | Rescue of the Epstein–Barr Virus BZLF1 Mutant, Z(S186A), Early Gene Activation Defect by the BRLF1 Gene Product. Virology, 1998, 251, 187-197. | 2.4 | 59 |
| 23 | Adoptively transferred Vγ9Vδ2 T cells show potent antitumor effects in a preclinical B cell lymphomagenesis model. JCI Insight, 2017, 2, . | 5.0 | 56 |
| 24 | LMP1-deficient Epstein-Barr virus mutant requires T cells for lymphomagenesis. Journal of Clinical Investigation, 2015, 125, 304-315. | 8.2 | 56 |
| 25 | Roles of lytic viral infection and IL-6 in early versus late passage lymphoblastoid cell lines and EBV-associated lymphoproliferative disease. International Journal of Cancer, 2007, 121, 1274-1281. | 5.1 | 55 |
| 26 | Hypoxia-inducible factor-1α plays roles in Epstein-Barr virus's natural life cycle and tumorigenesis by inducing lytic infection through direct binding to the immediate-early BZLF1 gene promoter. PLoS Pathogens, 2017, 13, e1006404. | 4.7 | 55 |
| 27 | The BRRF1 Early Gene of Epstein-Barr Virus Encodes a Transcription Factor That Enhances Induction of Lytic Infection by BRLF1. Journal of Virology, 2004, 78, 4983-4992. | 3.4 | 54 |
| 28 | Viral Genome Methylation Differentially Affects the Ability of BZLF1 versus BRLF1 To Activate Epstein-Barr Virus Lytic Gene Expression and Viral Replication. Journal of Virology, 2013, 87, 935-950. | 3.4 | 53 |
| 29 | Human papillomavirus promotes Epstein-Barr virus maintenance and lytic reactivation in immortalized oral keratinocytes. Virology, 2016, 495, 52-62. | 2.4 | 50 |
| 30 | ZEB1 and c-Jun Levels Contribute to the Establishment of Highly Lytic Epstein-Barr Virus Infection in Gastric AGS Cells. Journal of Virology, 2007, 81, 10113-10122. | 3.4 | 49 |
| 31 | Epstein-Barr Virus Immediate-Early Protein BRLF1 Interacts with CBP, Promoting Enhanced BRLF1 Transactivation. Journal of Virology, 2001, 75, 6228-6234. | 3.4 | 44 |
| 32 | Lenalidomide, Thalidomide, and Pomalidomide Reactivate the Epstein–Barr Virus Lytic Cycle through Phosphoinositide 3-Kinase Signaling and Ikaros Expression. Clinical Cancer Research, 2016, 22, 4901-4912. | 7.0 | 41 |
| 33 | Differentiation-Dependent LMP1 Expression Is Required for Efficient Lytic Epstein-Barr Virus Reactivation in Epithelial Cells. Journal of Virology, 2017, 91, . | 3.4 | 40 |
| 34 | Latent Membrane Protein 1 (LMP1) and LMP2A Collaborate To Promote Epstein-Barr Virus-Induced B Cell Lymphomas in a Cord Blood-Humanized Mouse Model but Are Not Essential. Journal of Virology, 2017, 91, . | 3.4 | 33 |
| 35 | Cellular Transcription Factor Oct-1 Interacts with the Epstein-Barr Virus BRLF1 Protein To Promote Disruption of Viral Latency. Journal of Virology, 2011, 85, 8940-8953. | 3.4 | 32 |
| 36 | The B-Cell Specific Transcription Factor, Oct-2, Promotes Epstein-Barr Virus Latency by Inhibiting the Viral Immediate-Early Protein, BZLF1. PLoS Pathogens, 2012, 8, e1002516. | 4.7 | 32 |

SHANNON C KENNEY

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | The Epstein-Barr Virus BRRF1 Protein, Na, Induces Lytic Infection in a TRAF2- and p53-Dependent Manner. Journal of Virology, 2011, 85, 4318-4329. | 3.4 | 30 |
| 38 | 5-hydroxymethylation of the EBV genome regulates the latent to lytic switch. Proceedings of the National Academy of Sciences of the United States of America, 2015, 112, E7257-65. | 7.1 | 28 |
| 39 | Epstein-Barr Virus Infection Promotes Epithelial Cell Growth by Attenuating Differentiation-Dependent Exit from the Cell Cycle. MBio, 2019, 10, . | 4.1 | 25 |
| 40 | B cells infected with Type 2 Epstein-Barr virus (EBV) have increased NFATc1/NFATc2 activity and enhanced lytic gene expression in comparison to Type 1 EBV infection. PLoS Pathogens, 2020, 16, e1008365. | 4.7 | 24 |
| 41 | An EBNA3C-deleted Epstein-Barr virus (EBV) mutant causes B-cell lymphomas with delayed onset in a cord blood-humanized mouse model. PLoS Pathogens, 2018, 14, e1007221. | 4.7 | 22 |
| 42 | CCAAT/enhancer binding proteins α and β regulate the tumor necrosis factor receptor 1 gene promoter. Molecular Immunology, 2009, 46, 2706-2713. | 2.2 | 20 |
| 43 | Complete and Durable Responses in Primary Central Nervous System Posttransplant Lymphoproliferative Disorder with Zidovudine, Ganciclovir, Rituximab, and Dexamethasone. Clinical Cancer Research, 2018, 24, 3273-3281. | 7.0 | 20 |
| 44 | Restricted TET2 Expression in Germinal Center Type B Cells Promotes Stringent Epstein-Barr Virus Latency. Journal of Virology, 2017, 91, . | 3.4 | 18 |
| 45 | EBNA2-deleted Epstein-Barr virus (EBV) isolate, P3HR1, causes Hodgkin-like lymphomas and diffuse large B cell lymphomas with type II and Wp-restricted latency types in humanized mice. PLoS Pathogens, 2020, 16, e1008590. | 4.7 | 16 |
| 46 | Epstein–Barr Virus Gene BARF1 Expression is Regulated by the Epithelial Differentiation Factor ΔNp63α in Undifferentiated Nasopharyngeal Carcinoma. Cancers, 2018, 10, 76. | 3.7 | 14 |
| 47 | Human Cytomegalovirus Productively Replicates <i>In Vitro</i> in Undifferentiated Oral Epithelial Cells. Journal of Virology, 2018, 92, . | 3.4 | 10 |
| 48 | Reduced IRF4 expression promotes lytic phenotype in Type 2 EBV-infected B cells. PLoS Pathogens, 2022, 18, e1010453. | 4.7 | 10 |
| 49 | Hsp90 inhibitors: A potential treatment for latent EBV infection?. Cell Cycle, 2010, 9, 1665-1666. | 2.6 | 9 |
| 50 | Hippo signaling effectors YAP and TAZ induce Epstein-Barr Virus (EBV) lytic reactivation through TEADs in epithelial cells. PLoS Pathogens, 2021, 17, e1009783. | 4.7 | 9 |
| 51 | ΔNp63α promotes Epstein-Barr virus latency in undifferentiated epithelial cells. PLoS Pathogens, 2021, 17, e1010045. | 4.7 | 8 |
| 52 | Development of a novel inducer for EBV lytic therapy. Bioorganic and Medicinal Chemistry Letters, 2019, 29, 2259-2264. | 2.2 | 7 |
| 53 | An EBNA3A-Mutated Epstein-Barr Virus Retains the Capacity for Lymphomagenesis in a Cord Blood-Humanized Mouse Model. Journal of Virology, 2020, 94, . | 3.4 | 5 |
| 54 | Thalidomide, Lenalidomide and Pomalidomide Disrupt Epstein-Barr Virus (EBV) Latency: Clinical Implications. Blood, 2013, 122, 3499-3499. | 1.4 | 2 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | The Epstein-Barr virus EBNA2 protein induces a subset of NOTCH target genes in thyroid cancer cell lines but fails to suppress proliferation. Surgery, 2017, 161, 195-201. | 1.9 | 1 |
| 56 | Validation of Arrayâ€based RNA Expression Profiles in Paraffinâ€embedded Samples of Epsteinâ€Barr Virusâ€related Malignancy. FASEB Journal, 2011, 25, lb313. | 0.5 | 0 |
| 57 | Epstein-Barr Virus Kinase-Targeted Therapy for Primary Central Nervous System Post-Transplant Lymphoproliferative Disorder. Blood, 2014, 124, 1750-1750. | 1.4 | Ο |