

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7017261/seung-u-kim-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| | | | |
|--------------------|-------------------------|----------------|-----------------|
| 334 papers | 7,840 citations | 44 h-index | 72 g-index |
| 373 ext. papers | 9,890 ext. citations | 4.8 avg, IF | 6.13 L-index |

| # | Paper | IF | Citations |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 334 | New trends on obesity and NAFLD in Asia. <i>Journal of Hepatology</i> , 2017 , 67, 862-873 | 13.4 | 448 |
| 333 | Risk assessment of hepatitis B virus-related hepatocellular carcinoma development using liver stiffness measurement (FibroScan). <i>Hepatology</i> , 2011 , 53, 885-94 | 11.2 | 285 |
| 332 | Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017-Part 1: Definition, risk factors and assessment. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018 , 33, 70-85 | 4 | 211 |
| 331 | Performance of transient elastography for the staging of liver fibrosis in patients with chronic hepatitis B: a meta-analysis. <i>PLoS ONE</i> , 2012 , 7, e44930 | 3.7 | 202 |
| 330 | Sarcopaenia is associated with NAFLD independently of obesity and insulin resistance: Nationwide surveys (KNHANES 2008-2011). <i>Journal of Hepatology</i> , 2015 , 63, 486-93 | 13.4 | 186 |
| 329 | Sarcopenia is associated with significant liver fibrosis independently of obesity and insulin resistance in nonalcoholic fatty liver disease: Nationwide surveys (KNHANES 2008-2011). <i>Hepatology</i> , 2016 , 63, 776-86 | 11.2 | 175 |
| 328 | Usefulness of FibroScan for detection of early compensated liver cirrhosis in chronic hepatitis B. <i>Digestive Diseases and Sciences</i> , 2009 , 54, 1758-63 | 4 | 115 |
| 327 | Controlled attenuation parameter (CAP) for detection of hepatic steatosis in patients with chronic liver diseases: a prospective study of a native Korean population. <i>Liver International</i> , 2014 , 34, 102-9 | 7.9 | 113 |
| 326 | Complete response at first chemoembolization is still the most robust predictor for favorable outcome in hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2015 , 62, 1304-10 | 13.4 | 108 |
| 325 | The Laennec staging system for histological sub-classification of cirrhosis is useful for stratification of prognosis in patients with liver cirrhosis. <i>Journal of Hepatology</i> , 2012 , 57, 556-63 | 13.4 | 101 |
| 324 | Increased risk of hepatocellular carcinoma in chronic hepatitis B patients with transient elastography-defined subclinical cirrhosis. <i>Hepatology</i> , 2015 , 61, 1851-9 | 11.2 | 100 |
| 323 | Revision and update on clinical practice guideline for liver cirrhosis. <i>The Korean Journal of Hepatology</i> , 2012 , 18, 1-21 | | 97 |
| 322 | A multicenter study of entecavir vs. tenofovir on prognosis of treatment-naïve chronic hepatitis B in South Korea. <i>Journal of Hepatology</i> , 2019 , 71, 456-464 | 13.4 | 91 |
| 321 | Prediction of development of liver-related events by transient elastography in hepatitis B patients with complete virological response on antiviral therapy. <i>American Journal of Gastroenterology</i> , 2014 , 109, 1241-9 | 0.7 | 89 |
| 320 | How can we enhance the performance of liver stiffness measurement using FibroScan in diagnosing liver cirrhosis in patients with chronic hepatitis B?. <i>Journal of Clinical Gastroenterology</i> , 2010 , 44, 66-71 | 3 | 88 |
| 319 | Applicability of BCLC stage for prognostic stratification in comparison with other staging systems: single centre experience from long-term clinical outcomes of 1717 treatment-naïve patients with hepatocellular carcinoma. <i>Liver International</i> , 2012 , 32, 1120-7 | 7.9 | 83 |
| 318 | Prediction of liver-related events using fibroscan in chronic hepatitis B patients showing advanced liver fibrosis. <i>PLoS ONE</i> , 2012 , 7, e36676 | 3.7 | 76 |

| | | | |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 317 | Clinical applications of transient elastography. <i>Clinical and Molecular Hepatology</i> , 2012 , 18, 163-73 | 6.9 | 74 |
| 316 | Validation of hepatitis B virus-related hepatocellular carcinoma prediction models in the era of antiviral therapy. <i>Hepatology</i> , 2015 , 62, 1757-66 | 11.2 | 73 |
| 315 | Liver stiffness measurement using acoustic radiation force impulse (ARFI) elastography and effect of necroinflammation. <i>Digestive Diseases and Sciences</i> , 2012 , 57, 1682-91 | 4 | 73 |
| 314 | The Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017-Part 2: Management and special groups. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018 , 33, 86-98 ⁴ | | 71 |
| 313 | Radiological response predicts survival following transarterial chemoembolisation in patients with unresectable hepatocellular carcinoma. <i>Alimentary Pharmacology and Therapeutics</i> , 2012 , 35, 1343-50 | 6.1 | 70 |
| 312 | Prediction of postoperative hepatic insufficiency by liver stiffness measurement (FibroScan((R))) before curative resection of hepatocellular carcinoma: a pilot study. <i>Hepatology International</i> , 2008 , 2, 471-7 | 8.8 | 66 |
| 311 | Prospective comparison of prognostic values of modified Response Evaluation Criteria in Solid Tumours with European Association for the Study of the Liver criteria in hepatocellular carcinoma following chemoembolisation. <i>European Journal of Cancer</i> , 2013 , 49, 826-34 | 7.5 | 65 |
| 310 | Liver stiffness measurement in combination with noninvasive markers for the improved diagnosis of B-viral liver cirrhosis. <i>Journal of Clinical Gastroenterology</i> , 2009 , 43, 267-71 | 3 | 63 |
| 309 | What are the true normal liver stiffness values using FibroScan?: a prospective study in healthy living liver and kidney donors in South Korea. <i>Liver International</i> , 2010 , 30, 268-74 | 7.9 | 62 |
| 308 | Clinical outcomes and predictors for relapse after cessation of oral antiviral treatment in chronic hepatitis B patients. <i>Journal of Gastroenterology</i> , 2016 , 51, 830-9 | 6.9 | 61 |
| 307 | Hepatocellular carcinoma presenting with bone metastasis: clinical characteristics and prognostic factors. <i>Journal of Cancer Research and Clinical Oncology</i> , 2008 , 134, 1377-84 | 4.9 | 59 |
| 306 | Prospective validation of ELF test in comparison with Fibroscan and FibroTest to predict liver fibrosis in Asian subjects with chronic hepatitis B. <i>PLoS ONE</i> , 2012 , 7, e41964 | 3.7 | 59 |
| 305 | Metabolic Dysfunction-Associated Fatty Liver Disease and Incident Cardiovascular Disease Risk: A Nationwide Cohort Study. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , 19, 2138-2147.e10 | 6.9 | 59 |
| 304 | Clinical application of liver stiffness measurement using transient elastography in chronic liver disease from longitudinal perspectives. <i>World Journal of Gastroenterology</i> , 2013 , 19, 1890-900 | 5.6 | 58 |
| 303 | Accuracy of transient elastography in assessing liver fibrosis in chronic viral hepatitis: A multicentre, retrospective study. <i>Liver International</i> , 2015 , 35, 2246-55 | 7.9 | 56 |
| 302 | Randomized, prospective, comparative study on the effects and safety of sorafenib vs. hepatic arterial infusion chemotherapy in patients with advanced hepatocellular carcinoma with portal vein tumor thrombosis. <i>Cancer Chemotherapy and Pharmacology</i> , 2018 , 82, 469-478 | 3.5 | 54 |
| 301 | Prediction of recurrence after curative resection of hepatocellular carcinoma using liver stiffness measurement (FibroScan®). <i>Annals of Surgical Oncology</i> , 2012 , 19, 4278-86 | 3.1 | 54 |
| 300 | Combined use of AFP, PIVKA-II, and AFP-L3 as tumor markers enhances diagnostic accuracy for hepatocellular carcinoma in cirrhotic patients. <i>Scandinavian Journal of Gastroenterology</i> , 2016 , 51, 344-53 ⁴ | 3.4 | 52 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 299 | Addition of tumor multiplicity improves the prognostic performance of the hepatoma arterial-embolization prognostic score. <i>Liver International</i> , 2016 , 36, 100-7 | 7.9 | 51 |
| 298 | Prognostic value of the 7th edition of the AJCC staging system as a clinical staging system in patients with hepatocellular carcinoma. <i>European Journal of Cancer</i> , 2011 , 47, 2568-75 | 7.5 | 50 |
| 297 | Clinical application of transient elastography in patients with chronic viral hepatitis receiving antiviral treatment. <i>Liver International</i> , 2015 , 35, 1103-15 | 7.9 | 49 |
| 296 | Clinical features, image findings, and prognosis of inflammatory pseudotumor of the liver: a multicenter experience of 45 cases. <i>Gut and Liver</i> , 2014 , 8, 58-63 | 4.8 | 49 |
| 295 | Prognostic value of E-fetoprotein and des-E-carboxy prothrombin responses in patients with hepatocellular carcinoma treated with transarterial chemoembolization. <i>BMC Cancer</i> , 2013 , 13, 5 | 4.8 | 48 |
| 294 | The accuracy of noninvasive methods in predicting the development of hepatocellular carcinoma and hepatic decompensation in patients with chronic hepatitis B. <i>Journal of Clinical Gastroenterology</i> , 2012 , 46, 518-25 | 3 | 48 |
| 293 | Obesity is more closely related with hepatic steatosis and fibrosis measured by transient elastography than metabolic health status. <i>Metabolism: Clinical and Experimental</i> , 2017 , 66, 23-31 | 12.7 | 46 |
| 292 | Prospective validation of FibroTest in comparison with liver stiffness for predicting liver fibrosis in Asian subjects with chronic hepatitis B. <i>PLoS ONE</i> , 2012 , 7, e35825 | 3.7 | 44 |
| 291 | Factors that affect the diagnostic accuracy of liver fibrosis measurement by Fibroscan in patients with chronic hepatitis B. <i>Alimentary Pharmacology and Therapeutics</i> , 2010 , 32, 498-505 | 6.1 | 44 |
| 290 | Discrimination of Nonalcoholic Steatohepatitis Using Transient Elastography in Patients with Nonalcoholic Fatty Liver Disease. <i>PLoS ONE</i> , 2016 , 11, e0157358 | 3.7 | 44 |
| 289 | Evolution from WHO to EASL and mRECIST for hepatocellular carcinoma: considerations for tumor response assessment. <i>Expert Review of Gastroenterology and Hepatology</i> , 2015 , 9, 335-48 | 4.2 | 43 |
| 288 | Discordance between liver biopsy and Fibroscan in assessing liver fibrosis in chronic hepatitis b: risk factors and influence of necroinflammation. <i>PLoS ONE</i> , 2012 , 7, e32233 | 3.7 | 43 |
| 287 | Normal liver elasticity values using acoustic radiation force impulse imaging: a prospective study in healthy living liver and kidney donors. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012 , 27, 130-6 | 4 | 42 |
| 286 | Risk prediction for patients with hepatocellular carcinoma undergoing chemoembolization: development of a prediction model. <i>Liver International</i> , 2016 , 36, 92-9 | 7.9 | 41 |
| 285 | Surgical resection after down-staging of locally advanced hepatocellular carcinoma by localized concurrent chemoradiotherapy. <i>Annals of Surgical Oncology</i> , 2014 , 21, 3646-53 | 3.1 | 41 |
| 284 | Prognostic significance of the worst grade in hepatocellular carcinoma with heterogeneous histologic grades of differentiation. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2013 , 28, 1384-90 | 4 | 41 |
| 283 | Liver stiffness measurement using FibroScan is influenced by serum total bilirubin in acute hepatitis. <i>Liver International</i> , 2009 , 29, 810-5 | 7.9 | 41 |
| 282 | Characterization of focal liver masses using acoustic radiation force impulse elastography. <i>World Journal of Gastroenterology</i> , 2013 , 19, 219-26 | 5.6 | 41 |

| | | | |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 281 | Congestive hepatopathy after Fontan operation and related factors assessed by transient elastography. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2014 , 148, 1498-505 | 1.5 | 40 |
| 280 | Survival of hepatocellular carcinoma patients may be improved in surveillance interval not more than 6 months compared with more than 6 months: a 15-year prospective study. <i>Journal of Clinical Gastroenterology</i> , 2013 , 47, 538-44 | 3 | 38 |
| 279 | Non-invasive assessment of changes in liver fibrosis via liver stiffness measurement in patients with chronic hepatitis B: impact of antiviral treatment on fibrosis regression. <i>Hepatology International</i> , 2010 , 4, 673-80 | 8.8 | 37 |
| 278 | Advancing the global public health agenda for NAFLD: a consensus statement. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2021 , | 24.2 | 37 |
| 277 | Number of target lesions for EASL and modified RECIST to predict survivals in hepatocellular carcinoma treated with chemoembolization. <i>Clinical Cancer Research</i> , 2013 , 19, 1503-11 | 12.9 | 36 |
| 276 | Histological subclassification of cirrhosis can predict recurrence after curative resection of hepatocellular carcinoma. <i>Liver International</i> , 2014 , 34, 1008-17 | 7.9 | 35 |
| 275 | Factors associated with significant liver fibrosis assessed using transient elastography in general population. <i>World Journal of Gastroenterology</i> , 2015 , 21, 1158-66 | 5.6 | 34 |
| 274 | Leflunomide increases the risk of silent liver fibrosis in patients with rheumatoid arthritis receiving methotrexate. <i>Arthritis Research and Therapy</i> , 2012 , 14, R232 | 5.7 | 34 |
| 273 | Gastroduodenal complications after concurrent chemoradiation therapy in patients with hepatocellular carcinoma: endoscopic findings and risk factors. <i>International Journal of Radiation Oncology Biology Physics</i> , 2011 , 81, 1343-51 | 4 | 34 |
| 272 | Improvement of Liver Fibrosis after Long-Term Antiviral Therapy Assessed by Fibroscan in Chronic Hepatitis B Patients With Advanced Fibrosis. <i>American Journal of Gastroenterology</i> , 2017 , 112, 882-891 | 0.7 | 33 |
| 271 | Prognostic Significance of ¹⁸ F-FDG Uptake in Hepatocellular Carcinoma Treated with Transarterial Chemoembolization or Concurrent Chemoradiotherapy: A Multicenter Retrospective Cohort Study. <i>Journal of Nuclear Medicine</i> , 2016 , 57, 509-16 | 8.9 | 33 |
| 270 | Risk assessment of clinical outcomes in Asian patients with chronic hepatitis B using enhanced liver fibrosis test. <i>Hepatology</i> , 2014 , 60, 1911-9 | 11.2 | 33 |
| 269 | The usefulness of liver stiffness measurement using FibroScan in chronic hepatitis C in South Korea: a multicenter, prospective study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2011 , 26, 171-8 ⁴ | | 33 |
| 268 | Liver fibrosis assessed with transient elastography is an independent risk factor for ischemic stroke. <i>Atherosclerosis</i> , 2017 , 260, 156-162 | 3.1 | 32 |
| 267 | A nomogram based on liver stiffness predicts postoperative complications in patients with hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2020 , 73, 855-862 | 13.4 | 32 |
| 266 | Role of surgical resection for multiple hepatocellular carcinomas. <i>World Journal of Gastroenterology</i> , 2013 , 19, 366-74 | 5.6 | 32 |
| 265 | Regulation of HK2 expression through alterations in CpG methylation of the HK2 promoter during progression of hepatocellular carcinoma. <i>Oncotarget</i> , 2016 , 7, 41798-41810 | 3.3 | 32 |
| 264 | Validation of PAGE-B model in Asian chronic hepatitis B patients receiving entecavir or tenofovir. <i>Liver International</i> , 2017 , 37, 1788-1795 | 7.9 | 31 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 263 | Impact of spontaneous hepatocellular carcinoma rupture on recurrence pattern and long-term surgical outcomes after partial hepatectomy. <i>World Journal of Surgery</i> , 2014 , 38, 2070-8 | 3.3 | 31 |
| 262 | Acoustic Radiation Force Impulse Measurement in Renal Transplantation: A Prospective, Longitudinal Study With Protocol Biopsies. <i>Medicine (United States)</i> , 2015 , 94, e1590 | 1.8 | 31 |
| 261 | Range of Normal Liver Stiffness and Factors Associated With Increased Stiffness Measurements in Apparently Healthy Individuals. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 54-64.e1 | 6.9 | 31 |
| 260 | Ascitic fluid infection in patients with hepatitis B virus-related liver cirrhosis: culture-negative neutrocytic ascites versus spontaneous bacterial peritonitis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2010 , 25, 122-8 | 4 | 30 |
| 259 | Clinical characteristics and prognosis of hepatocellular carcinoma with different sets of serum AFP and PIVKA-II levels. <i>European Journal of Gastroenterology and Hepatology</i> , 2012 , 24, 849-56 | 2.2 | 30 |
| 258 | Factors affecting the accuracy of controlled attenuation parameter (CAP) in assessing hepatic steatosis in patients with chronic liver disease. <i>PLoS ONE</i> , 2014 , 9, e98689 | 3.7 | 30 |
| 257 | Wisteria floribunda agglutinin-positive human Mac-2 binding protein predicts the risk of HBV-related liver cancer development. <i>Liver International</i> , 2017 , 37, 879-887 | 7.9 | 29 |
| 256 | Lower incidence of hepatocellular carcinoma and cirrhosis in hepatitis C patients with sustained virological response by pegylated interferon and ribavirin. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 573-81 | 4.1 | 29 |
| 255 | Association Between Level of Fibrosis, Rather Than Antiviral Regimen, and Outcomes of Patients With Chronic Hepatitis B. <i>Clinical Gastroenterology and Hepatology</i> , 2016 , 14, 1647-1656.e6 | 6.9 | 29 |
| 254 | Early on-treatment predictions of clinical outcomes using alpha-fetoprotein and des-gamma-carboxy prothrombin responses in patients with advanced hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012 , 27, 313-22 | 4 | 29 |
| 253 | Response rates to medical treatments and long-term clinical outcomes of nonsurgical patients with intestinal Behçet disease. <i>Journal of Clinical Gastroenterology</i> , 2010 , 44, e116-22 | 3 | 29 |
| 252 | Natural history of hepatitis B virus-related cirrhotic patients hospitalized to control ascites. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2008 , 23, 1722-7 | 4 | 29 |
| 251 | Liver stiffness-based model for prediction of hepatocellular carcinoma in chronic hepatitis B virus infection: comparison with histological fibrosis. <i>Liver International</i> , 2015 , 35, 1054-62 | 7.9 | 28 |
| 250 | Combination treatment of localized concurrent chemoradiation therapy and transarterial chemoembolization in locally advanced hepatocellular carcinoma with intrahepatic metastasis. <i>Cancer Chemotherapy and Pharmacology</i> , 2013 , 71, 165-73 | 3.5 | 28 |
| 249 | Transient elastography in chronic hepatitis B: an Asian perspective. <i>World Journal of Gastroenterology</i> , 2010 , 16, 5173-80 | 5.6 | 28 |
| 248 | Comparison of treatment outcomes between balloon-occluded retrograde transvenous obliteration and transjugular intrahepatic portosystemic shunt for gastric variceal bleeding hemostasis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 1487-1494 | 4 | 27 |
| 247 | Long-term changes of liver stiffness values assessed using transient elastography in patients with chronic hepatitis B receiving entecavir. <i>Liver International</i> , 2014 , 34, 1216-23 | 7.9 | 27 |
| 246 | "Normal" liver stiffness values differ between men and women: a prospective study for healthy living liver and kidney donors in a native Korean population. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012 , 27, 781-8 | 4 | 27 |

| | | | |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 245 | Combined measurement of preoperative α -fetoprotein and des- γ -carboxy prothrombin predicts recurrence after curative resection in patients with hepatitis-B-related hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2012 , 131, 2332-41 | 7.5 | 26 |
| 244 | Prognostic factors and 10-year survival in patients with hepatocellular carcinoma after curative hepatectomy. <i>Journal of Gastrointestinal Surgery</i> , 2011 , 15, 598-607 | 3.3 | 26 |
| 243 | The Relationship between Type 2 Diabetes Mellitus and Non-Alcoholic Fatty Liver Disease Measured by Controlled Attenuation Parameter. <i>Yonsei Medical Journal</i> , 2016 , 57, 885-92 | 3 | 26 |
| 242 | Sarcopenia is associated with the risk of significant liver fibrosis in metabolically unhealthy subjects with chronic hepatitis B. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 48, 300-312 | 6.1 | 25 |
| 241 | Inhibition of tumour angiogenesis and growth by small hairpin HIF-1 α and IL-8 in hepatocellular carcinoma. <i>Liver International</i> , 2014 , 34, 632-42 | 7.9 | 25 |
| 240 | Gamma-glutamyl transpeptidase-to-platelet ratio is an independent predictor of hepatitis B virus-related liver cancer. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 1221-1229 | 4 | 25 |
| 239 | Serum Dickkopf-1 as a Biomarker for the Diagnosis of Hepatocellular Carcinoma. <i>Yonsei Medical Journal</i> , 2015 , 56, 1296-306 | 3 | 25 |
| 238 | Hepatic arterial damage after transarterial chemoembolization for the treatment of hepatocellular carcinoma: comparison of drug-eluting bead and conventional chemoembolization in a retrospective controlled study. <i>Acta Radiologica</i> , 2017 , 58, 131-139 | 2 | 24 |
| 237 | Development of a New Nomogram Including Neutrophil-to-Lymphocyte Ratio to Predict Survival in Patients with Hepatocellular Carcinoma Undergoing Transarterial Chemoembolization. <i>Cancers</i> , 2019 , 11, | 6.6 | 24 |
| 236 | Cryoablation of Hepatocellular Carcinoma with High-Risk for Percutaneous Ablation: Safety and Efficacy. <i>CardioVascular and Interventional Radiology</i> , 2016 , 39, 1447-54 | 2.7 | 24 |
| 235 | Early on-treatment change in liver stiffness predicts development of liver-related events in chronic hepatitis B patients receiving antiviral therapy. <i>Liver International</i> , 2013 , 33, 180-9 | 7.9 | 24 |
| 234 | Conventional versus drug-eluting beads chemoembolization for hepatocellular carcinoma: Emphasis on the impact of tumor size. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 487-496 | 4 | 24 |
| 233 | A novel model to predict esophageal varices in patients with compensated cirrhosis using acoustic radiation force impulse elastography. <i>PLoS ONE</i> , 2015 , 10, e0121009 | 3.7 | 24 |
| 232 | Use of Wisteria Floribunda Agglutinin-Positive Human Mac-2 Binding Protein in Assessing Risk of Hepatocellular Carcinoma Due to Hepatitis B Virus. <i>Medicine (United States)</i> , 2016 , 95, e3328 | 1.8 | 24 |
| 231 | Nonalcoholic Fatty Liver Disease and Sarcopenia Are Independently Associated With Cardiovascular Risk. <i>American Journal of Gastroenterology</i> , 2020 , 115, 584-595 | 0.7 | 24 |
| 230 | Clinical outcomes and prognostic factors of patients with advanced hepatocellular carcinoma treated with sorafenib as first-line therapy: a Korean multicenter study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 1463-9 | 4 | 23 |
| 229 | Association between IL28B polymorphisms and spontaneous clearance of hepatitis B virus infection. <i>PLoS ONE</i> , 2013 , 8, e69166 | 3.7 | 23 |
| 228 | External validation of the modified PAGE-B score in Asian chronic hepatitis B patients receiving antiviral therapy. <i>Liver International</i> , 2019 , 39, 1624-1630 | 7.9 | 22 |

| | | | |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 227 | Red cell volume distribution width-to-platelet ratio in assessment of liver fibrosis in patients with chronic hepatitis B. <i>Liver International</i> , 2016 , 36, 24-30 | 7.9 | 22 |
| 226 | Tenofovir versus tenofovir plus entecavir for chronic hepatitis B with lamivudine resistance and entecavir resistance. <i>Journal of Viral Hepatitis</i> , 2017 , 24, 141-147 | 3.4 | 22 |
| 225 | Use of the delta neutrophil index as a prognostic factor of mortality in patients with spontaneous bacterial peritonitis: implications of a simple and useful marker. <i>PLoS ONE</i> , 2014 , 9, e86884 | 3.7 | 22 |
| 224 | Risk assessment of liver-related events using transient elastography in patients with chronic hepatitis B receiving entecavir. <i>Journal of Clinical Gastroenterology</i> , 2014 , 48, 272-8 | 3 | 22 |
| 223 | Clinical features and prognosis of hepatocellular carcinoma in Mongolia: a multicentre study. <i>Hepatology International</i> , 2012 , 6, 763-9 | 8.8 | 22 |
| 222 | Prediction of the varices needing treatment with non-invasive tests in patients with compensated advanced chronic liver disease. <i>Liver International</i> , 2019 , 39, 1071-1079 | 7.9 | 22 |
| 221 | KASL clinical practice guidelines: Management of nonalcoholic fatty liver disease. <i>Clinical and Molecular Hepatology</i> , 2021 , 27, 363-401 | 6.9 | 21 |
| 220 | Feasibility of dynamic risk prediction for hepatocellular carcinoma development in patients with chronic hepatitis B. <i>Liver International</i> , 2018 , 38, 676-686 | 7.9 | 20 |
| 219 | Development and validation of a prognostic model for patients with hepatocellular carcinoma undergoing radiofrequency ablation. <i>Cancer Medicine</i> , 2019 , 8, 5023-5032 | 4.8 | 20 |
| 218 | Prognosis of hepatocellular carcinoma patients with extrahepatic metastasis and the controllability of intrahepatic lesions. <i>Clinical and Experimental Metastasis</i> , 2014 , 31, 475-82 | 4.7 | 20 |
| 217 | A case of hepatocellular carcinoma within hepatocellular adenoma in a non-cirrhotic male. <i>Korean Journal of Internal Medicine</i> , 2009 , 24, 147-52 | 2.5 | 20 |
| 216 | Role of splenectomy in patients with hepatocellular carcinoma and hypersplenism. <i>ANZ Journal of Surgery</i> , 2013 , 83, 865-70 | 1 | 19 |
| 215 | Efficacy of sorafenib monotherapy versus sorafenib-based loco-regional treatments in advanced hepatocellular carcinoma. <i>PLoS ONE</i> , 2013 , 8, e77240 | 3.7 | 19 |
| 214 | Partial virological response to entecavir in treatment-naïve patients with chronic hepatitis B. <i>Antiviral Therapy</i> , 2011 , 16, 469-77 | 1.6 | 19 |
| 213 | Subcirrhotic liver stiffness by FibroScan correlates with lower risk of hepatocellular carcinoma in patients with HBV-related cirrhosis. <i>Hepatology International</i> , 2017 , 11, 268-276 | 8.8 | 18 |
| 212 | Use of transient elastography to predict de novo recurrence after radiofrequency ablation for hepatocellular carcinoma. <i>OncoTargets and Therapy</i> , 2015 , 8, 347-56 | 4.4 | 18 |
| 211 | Liver stiffness value-based risk estimation of late recurrence after curative resection of hepatocellular carcinoma: development and validation of a predictive model. <i>PLoS ONE</i> , 2014 , 9, e99167 | 3.7 | 18 |
| 210 | Transient elastography-based risk estimation of hepatitis B virus-related occurrence of hepatocellular carcinoma: development and validation of a predictive model. <i>OncoTargets and Therapy</i> , 2013 , 6, 1463-9 | 4.4 | 18 |

| | | | |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 209 | Clinical characteristics and treatment outcomes of hepatocellular carcinoma with inferior vena cava/heart invasion. <i>Anticancer Research</i> , 2011 , 31, 4641-6 | 2.3 | 18 |
| 208 | Prognostic value of the combined use of transient elastography and fibrotest in patients with chronic hepatitis B. <i>Liver International</i> , 2015 , 35, 455-62 | 7.9 | 17 |
| 207 | Clinical application of liver stiffness measurement using transient elastography: a surgical perspective. <i>Digestion</i> , 2013 , 88, 258-65 | 3.6 | 17 |
| 206 | Hepatic fibrosis assessed using transient elastography independently associated with coronary artery calcification. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2015 , 30, 1536-42 | 4 | 17 |
| 205 | Abnormal liver stiffness assessed using transient elastography (Fibroscan®) in HIV-infected patients without HBV/HCV coinfection receiving combined antiretroviral treatment. <i>PLoS ONE</i> , 2013 , 8, e52720 | 3.7 | 17 |
| 204 | Hepatocellular Carcinoma Risk Steadily Persists over Time Despite Long-Term Antiviral Therapy for Hepatitis B: A Multicenter Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020 , 29, 832-837 | 4 | 17 |
| 203 | A survey on transarterial chemoembolization refractoriness and a real-world treatment pattern for hepatocellular carcinoma in Korea. <i>Clinical and Molecular Hepatology</i> , 2020 , 26, 24-32 | 6.9 | 17 |
| 202 | From nonalcoholic fatty liver disease to metabolic-associated fatty liver disease: Big wave or ripple?. <i>Clinical and Molecular Hepatology</i> , 2021 , 27, 257-269 | 6.9 | 17 |
| 201 | Risk Assessment of Hepatocellular Carcinoma Using Transient Elastography Vs. Liver Biopsy in Chronic Hepatitis B Patients Receiving Antiviral Therapy. <i>Medicine (United States)</i> , 2016 , 95, e2985 | 1.8 | 17 |
| 200 | Subclassification of Barcelona Clinic Liver Cancer B and C hepatocellular carcinoma: A cohort study of the multicenter registry database. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2016 , 31, 842-7 | 4 | 17 |
| 199 | Liver Fibrosis, Not Steatosis, Associates with Long-Term Outcomes in Ischaemic Stroke Patients. <i>Cerebrovascular Diseases</i> , 2019 , 47, 32-39 | 3.2 | 16 |
| 198 | Probability-based interpretation of liver stiffness measurement in untreated chronic hepatitis B patients. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 1448-56 | 4 | 16 |
| 197 | Extremely low risk of hepatocellular carcinoma development in patients with chronic hepatitis B in immune-tolerant phase. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 52, 196-204 | 6.1 | 16 |
| 196 | Effects of Entecavir and Tenofovir on Renal Function in Patients with Hepatitis B Virus-Related Compensated and Decompensated Cirrhosis. <i>Gut and Liver</i> , 2017 , 11, 828-834 | 4.8 | 16 |
| 195 | Reduced risk of hepatocellular carcinoma by achieving a subcirrhotic liver stiffness through antiviral agents in hepatitis B virus-related advanced fibrosis or cirrhosis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018 , 33, 503-510 | 4 | 16 |
| 194 | Validation of risk prediction models for the development of HBV-related HCC: a retrospective multi-center 10-year follow-up cohort study. <i>Oncotarget</i> , 2017 , 8, 113213-113224 | 3.3 | 16 |
| 193 | Early Ectoprotein response predicts survival in patients with advanced hepatocellular carcinoma treated with sorafenib. <i>Journal of Hepatocellular Carcinoma</i> , 2015 , 2, 39-47 | 5.3 | 16 |
| 192 | Clinical usefulness of double biomarkers AFP and PIVKA-II for subdividing prognostic groups in locally advanced hepatocellular carcinoma. <i>Liver International</i> , 2014 , 34, 313-21 | 7.9 | 16 |

| | | | |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 191 | Adjuvant hepatic arterial infusional chemotherapy with 5-fluorouracil and cisplatin after curative resection of hepatocellular carcinoma. <i>Oncology</i> , 2011 , 81, 184-91 | 3.6 | 16 |
| 190 | Influence of hepatic steatosis on the outcomes of patients with chronic hepatitis B treated with entecavir and tenofovir. <i>Clinical and Molecular Hepatology</i> , 2019 , 25, 283-293 | 6.9 | 16 |
| 189 | Validation of the CAMD Score in Patients With Chronic Hepatitis B Virus Infection Receiving Antiviral Therapy. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 693-699.e1 | 6.9 | 16 |
| 188 | An optimized hepatocellular carcinoma prediction model for chronic hepatitis B with well-controlled viremia. <i>Liver International</i> , 2020 , 40, 1736-1743 | 7.9 | 15 |
| 187 | Long-term outcomes of chronic hepatitis B virus infection in the era of antiviral therapy in Korea. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2014 , 29, 1005-11 | 4 | 15 |
| 186 | Variability in liver stiffness values from different intercostal spaces. <i>Liver International</i> , 2009 , 29, 760-6 | 7.9 | 15 |
| 185 | Dynamics of the liver stiffness value using transient elastography during the perioperative period in patients with valvular heart disease. <i>PLoS ONE</i> , 2014 , 9, e92795 | 3.7 | 15 |
| 184 | Using transient elastography to predict hepatocellular carcinoma recurrence after radiofrequency ablation. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 1079-1086 | 4 | 14 |
| 183 | Can preoperative diffusion-weighted MRI predict postoperative hepatic insufficiency after curative resection of HBV-related hepatocellular carcinoma? A pilot study. <i>Magnetic Resonance Imaging</i> , 2010 , 28, 802-11 | 3.3 | 14 |
| 182 | Anti-hepatitis A virus seroprevalence among patients with chronic viral liver disease in Korea. <i>European Journal of Gastroenterology and Hepatology</i> , 2007 , 19, 923-6 | 2.2 | 14 |
| 181 | Risk assessment of hepatocellular carcinoma development for indeterminate hepatic nodules in patients with chronic hepatitis B. <i>Clinical and Molecular Hepatology</i> , 2019 , 25, 390-399 | 6.9 | 14 |
| 180 | Factors Influencing the Diagnostic Accuracy of Acoustic Radiation Force Impulse Elastography in Patients with Chronic Hepatitis B. <i>Gut and Liver</i> , 2016 , 10, 275-82 | 4.8 | 14 |
| 179 | Knockdown of HIF-1 α and IL-8 induced apoptosis of hepatocellular carcinoma triggers apoptosis of vascular endothelial cells. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2016 , 21, 85-95 | 5.4 | 13 |
| 178 | Prevalence and Predictors of Significant Fibrosis Among Subjects with Transient Elastography-Defined Nonalcoholic Fatty Liver Disease. <i>Digestive Diseases and Sciences</i> , 2017 , 62, 2150-2158 | 4.58 | 13 |
| 177 | Prognostic role of magnetic resonance imaging vs. computed tomography for hepatocellular carcinoma undergoing chemoembolization. <i>Liver International</i> , 2015 , 35, 1722-30 | 7.9 | 13 |
| 176 | Reproducibility of European Association for the Study of the Liver criteria and modified Response Evaluation Criteria in Solid Tumors in patients treated with sorafenib. <i>Liver International</i> , 2018 , 38, 1655-1663 | 7.9 | 13 |
| 175 | Incidence and risk factors of delayed postpolypectomy bleeding in patients with chronic liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2016 , 51, 618-24 | 2.4 | 13 |
| 174 | Relationship between Cerebral Microbleeds and Liver Stiffness Determined by Transient Elastography. <i>PLoS ONE</i> , 2015 , 10, e0139227 | 3.7 | 13 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 173 | Antiviral efficacy of lamivudine versus entecavir in patients with hepatitis B virus-related advanced hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012 , 27, 1528-34 | 4 | 13 |
| 172 | Normal enhanced liver fibrosis (ELF) values in apparently healthy subjects undergoing a health check-up and in living liver donors in South Korea. <i>Liver International</i> , 2013 , 33, 706-13 | 7.9 | 13 |
| 171 | Transarterial Radioembolization Versus Concurrent Chemoradiation Therapy for Locally Advanced Hepatocellular Carcinoma: A Propensity Score Matching Analysis. <i>International Journal of Radiation Oncology Biology Physics</i> , 2017 , 99, 396-406 | 4 | 12 |
| 170 | Optimal time for restoring the reliability of liver stiffness measurement in patients with chronic hepatitis B experiencing acute exacerbation. <i>Journal of Clinical Gastroenterology</i> , 2012 , 46, 602-7 | 3 | 12 |
| 169 | Dickkopf-1 induces angiogenesis via VEGF receptor 2 regulation independent of the Wnt signaling pathway. <i>Oncotarget</i> , 2017 , 8, 58974-58984 | 3.3 | 12 |
| 168 | Antiviral efficacies of currently available rescue therapies for multidrug-resistant chronic hepatitis B. <i>Clinical and Molecular Hepatology</i> , 2013 , 19, 29-35 | 6.9 | 12 |
| 167 | Noninvasive monitoring of hepatic steatosis: controlled attenuation parameter and magnetic resonance imaging-proton density fat fraction in patients with nonalcoholic fatty liver disease. <i>Expert Review of Gastroenterology and Hepatology</i> , 2019 , 13, 523-530 | 4.2 | 11 |
| 166 | Normal controlled attenuation parameter values: a prospective study of healthy subjects undergoing health checkups and liver donors in Korea. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 234-42 | 4 | 11 |
| 165 | Predictors of failure to detect early hepatocellular carcinoma in patients with chronic hepatitis B who received regular surveillance. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 47, 1201-1212 | 6.1 | 11 |
| 164 | How many valid measurements are necessary to assess liver fibrosis using FibroScan® in patients with chronic viral hepatitis? An analysis of subjects with at least 10 valid measurements. <i>Yonsei Medical Journal</i> , 2012 , 53, 337-45 | 3 | 11 |
| 163 | Serum Wisteria floribunda agglutinin-positive human Mac-2 binding protein level predicts recurrence of hepatitis B virus-related hepatocellular carcinoma after curative resection. <i>Clinical and Molecular Hepatology</i> , 2020 , 26, 33-44 | 6.9 | 11 |
| 162 | Successful hemostasis of intractable rectal variceal bleeding using variceal embolization. <i>World Journal of Gastroenterology</i> , 2015 , 21, 2558-62 | 5.6 | 11 |
| 161 | Quasispecies and pre-existing drug-resistant mutations of hepatitis B virus in patients with chronic hepatitis B. <i>Gut and Liver</i> , 2013 , 7, 329-34 | 4.8 | 11 |
| 160 | Predicting Liver-Related Events Using Transient Elastography in Chronic Hepatitis C Patients with Sustained Virological Response. <i>Gut and Liver</i> , 2016 , 10, 429-36 | 4.8 | 11 |
| 159 | Genetically Engineered Mouse Models for Liver Cancer. <i>Cancers</i> , 2019 , 12, | 6.6 | 11 |
| 158 | Unhealthy lifestyle habits and physical inactivity among Asian patients with non-alcoholic fatty liver disease. <i>Liver International</i> , 2020 , 40, 2719-2731 | 7.9 | 11 |
| 157 | Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. <i>Modern Rheumatology</i> , 2020 , 30, 70-77 | 3.3 | 11 |
| 156 | Comparison between chronic hepatitis B patients with untreated immune-tolerant phase vs. those with virological response by antivirals. <i>Scientific Reports</i> , 2019 , 9, 2508 | 4.9 | 10 |

| | | | |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----|
| 155 | Discontinuation of nucleos(t)ide analogues is not associated with a higher risk of HBsAg seroreversion after antiviral-induced HBsAg seroclearance: a nationwide multicentre study. <i>Gut</i> , 2020 , 69, 2214-2222 | 19.2 | 10 |
| 154 | Efficacy and Safety of Liver-Directed Concurrent Chemoradiotherapy and Sequential Sorafenib for Advanced Hepatocellular Carcinoma: A Prospective Phase 2 Trial. <i>International Journal of Radiation Oncology Biology Physics</i> , 2020 , 107, 106-115 | 4 | 10 |
| 153 | Alpha-Fetoprotein, Des-Gamma-Carboxy Prothrombin, and Modified RECIST Response as Predictors of Survival after Transarterial Radioembolization for Hepatocellular Carcinoma. <i>Journal of Vascular and Interventional Radiology</i> , 2019 , 30, 1194-1200.e1 | 2.4 | 10 |
| 152 | Maintaining remission in lamivudine-resistant patients with a virological response to adefovir add-on lamivudine after stopping lamivudine therapy. <i>Liver International</i> , 2014 , 34, 1543-9 | 7.9 | 10 |
| 151 | Non-invasive assessment of liver fibrosis: time to move from cross-sectional studies to longitudinal ones. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2010 , 25, 1472-3 | 4 | 10 |
| 150 | Combination of preS deletions and A1762T/G1764A mutations in HBV subgenotype C2 increases the risk of developing HCC. <i>Intervirology</i> , 2012 , 55, 296-302 | 2.5 | 10 |
| 149 | Discordance between liver biopsy and FibroTest in assessing liver fibrosis in chronic hepatitis B. <i>PLoS ONE</i> , 2013 , 8, e55759 | 3.7 | 10 |
| 148 | Tenofovir disoproxil fumarate monotherapy for nucleos(t)ide-naïve chronic hepatitis B patients in Korea: data from the clinical practice setting in a single-center cohort. <i>Clinical and Molecular Hepatology</i> , 2014 , 20, 261-6 | 6.9 | 10 |
| 147 | Sarcopenia Is Independently Associated with the Degree of Liver Fibrosis in Patients with Type 2 Diabetes Mellitus. <i>Gut and Liver</i> , 2020 , 14, 626-635 | 4.8 | 10 |
| 146 | Design and validation of risk prediction model for hepatocellular carcinoma development after sustained virological response in patients with chronic hepatitis C. <i>European Journal of Gastroenterology and Hepatology</i> , 2020 , 32, 378-385 | 2.2 | 10 |
| 145 | Transient Elastography is Superior to FIB-4 in Assessing the Risk of Hepatocellular Carcinoma in Patients With Chronic Hepatitis B. <i>Medicine (United States)</i> , 2016 , 95, e3434 | 1.8 | 10 |
| 144 | Prognosis of Untreated Minimally Active Chronic Hepatitis B Patients in Comparison With Virological Responders by Antivirals. <i>Clinical and Translational Gastroenterology</i> , 2019 , 10, e00036 | 4.2 | 10 |
| 143 | Bariatric Arterial Embolization with Non-spherical Polyvinyl Alcohol Particles for Ghrelin Suppression in a Swine Model. <i>CardioVascular and Interventional Radiology</i> , 2017 , 40, 744-749 | 2.7 | 9 |
| 142 | Staged partial hepatectomy versus transarterial chemoembolization for the treatment of spontaneous hepatocellular carcinoma rupture: a multicenter analysis in Korea. <i>Annals of Surgical Treatment and Research</i> , 2019 , 96, 275-282 | 2 | 9 |
| 141 | Progression of Untreated Minimally Active Chronic HBV Infection Compared to Inactive Infection. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 2808-2810.e2 | 6.9 | 9 |
| 140 | Natural History of Untreated HBeAg-Positive Chronic HBV Infection With Persistently Elevated HBV DNA but Normal Alanine Aminotransferase. <i>Clinical and Translational Gastroenterology</i> , 2020 , 11, e00140 | 4.2 | 9 |
| 139 | Acute variceal hemorrhage in patients with liver cirrhosis: weekend versus weekday admissions. <i>Yonsei Medical Journal</i> , 2012 , 53, 318-27 | 3 | 9 |
| 138 | Silencing of hypoxia-inducible factor-1 induces anti-tumor effects in hepatoma cell lines under tumor hypoxia. <i>PLoS ONE</i> , 2014 , 9, e103304 | 3.7 | 9 |

| | | | |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 137 | Non-invasive prediction of forthcoming cirrhosis-related complications. <i>World Journal of Gastroenterology</i> , 2014 , 20, 2613-23 | 5.6 | 9 |
| 136 | Comparison of the Effects of Telbivudine and Entecavir Treatment on Estimated Glomerular Filtration Rate in Patients with Chronic Hepatitis B. <i>Gut and Liver</i> , 2015 , 9, 776-83 | 4.8 | 9 |
| 135 | Liver Cirrhosis, Not Antiviral Therapy, Predicts Clinical Outcome in Cohorts with Heterogeneous Hepatitis B Viral Status. <i>Gut and Liver</i> , 2019 , 13, 197-205 | 4.8 | 9 |
| 134 | Lenvatinib is independently associated with the reduced risk of progressive disease when compared with sorafenib in patients with advanced hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021 , 36, 1317-1325 | 4 | 9 |
| 133 | Longitudinal assessment of alpha-fetoprotein for early detection of hepatocellular carcinoma in patients with cirrhosis. <i>Scandinavian Journal of Gastroenterology</i> , 2019 , 54, 1283-1290 | 2.4 | 8 |
| 132 | Orthotopic liver transplantation after the combined use of locoregional therapy and sorafenib for advanced hepatocellular carcinoma. <i>OncoTargets and Therapy</i> , 2013 , 6, 755-9 | 4.4 | 8 |
| 131 | The Modified Response Evaluation Criteria in Solid Tumors (RECIST) Yield a More Accurate Prognoses Than the RECIST 1.1 in Hepatocellular Carcinoma Treated with Transarterial Radioembolization. <i>Gut and Liver</i> , 2020 , 14, 765-774 | 4.8 | 8 |
| 130 | Hepatic Decompensation in Cirrhotic Patients Receiving Antiviral Therapy for Chronic Hepatitis B. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , 19, 1950-1958.e7 | 6.9 | 8 |
| 129 | Prediction of virologic response to tenofovir mono-rescue therapy for multidrug resistant chronic hepatitis B. <i>Journal of Medical Virology</i> , 2016 , 88, 1027-34 | 19.7 | 8 |
| 128 | Hepatic Steatosis Index in the Detection of Fatty Liver in Patients with Chronic Hepatitis B Receiving Antiviral Therapy. <i>Gut and Liver</i> , 2021 , 15, 117-127 | 4.8 | 8 |
| 127 | Feasibility of dynamic risk assessment for patients with repeated trans-arterial chemoembolization for hepatocellular carcinoma. <i>BMC Cancer</i> , 2019 , 19, 363 | 4.8 | 7 |
| 126 | Non-invasive prediction of development of hepatocellular carcinoma using transient elastography in patients with chronic liver disease. <i>Expert Review of Gastroenterology and Hepatology</i> , 2014 , 8, 501-11 | 4.2 | 7 |
| 125 | Spontaneous bacterial peritonitis in patients with hepatitis B virus-related liver cirrhosis: community-acquired versus nosocomial. <i>Yonsei Medical Journal</i> , 2012 , 53, 328-36 | 3 | 7 |
| 124 | Clinical course of virologic breakthrough after emergence of YMDD mutations in HBeAg-positive chronic hepatitis B. <i>Intervirolgy</i> , 2008 , 51, 293-8 | 2.5 | 7 |
| 123 | Outcome of adefovir add-on lamivudine rescue therapy of up to 5 years in patients with lamivudine-resistant chronic hepatitis B. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2016 , 31, 241-7 | 4 | 7 |
| 122 | The impact of steatosis on liver stiffness measurement in patients with chronic hepatitis B. <i>Hepato-Gastroenterology</i> , 2010 , 57, 832-8 | | 7 |
| 121 | Transarterial Chemoembolization in Treatment-Naïve and Recurrent Hepatocellular Carcinoma: A Propensity-Matched Outcome Analysis. <i>Digestive Diseases and Sciences</i> , 2019 , 64, 3660-3668 | 4 | 6 |
| 120 | Hepatic Fibrosis Assessed Using Fibrosis-4 Index Is Predictive of All-Cause Mortality in Patients with Chronic Obstructive Pulmonary Disease. <i>International Journal of COPD</i> , 2020 , 15, 831-839 | 3 | 6 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 119 | Predictive score for hepatocellular carcinoma after hepatitis B e antigen loss in patients treated with entecavir or tenofovir. <i>Journal of Viral Hepatitis</i> , 2020 , 27, 1052-1060 | 3.4 | 6 |
| 118 | Efficacy of switching to telbivudine plus adefovir in suboptimal responders to lamivudine plus adefovir. <i>World Journal of Gastroenterology</i> , 2013 , 19, 7671-9 | 5.6 | 6 |
| 117 | Combination of Transient Elastography and an Enhanced Liver Fibrosis Test to Assess the Degree of Liver Fibrosis in Patients with Chronic Hepatitis B. <i>Gut and Liver</i> , 2018 , 12, 190-200 | 4.8 | 6 |
| 116 | Daclatasvir Plus Asunaprevir for the Treatment of Patients with Hepatitis C Virus Genotype 1b Infection: Real-World Efficacy, Changes in Liver Stiffness and Fibrosis Markers, and Safety. <i>Gut and Liver</i> , 2018 , 12, 324-330 | 4.8 | 6 |
| 115 | Prognosis of Early-Stage Hepatocellular Carcinoma: Comparison between Trans-Arterial Chemoembolization and Radiofrequency Ablation. <i>Cancers</i> , 2020 , 12, | 6.6 | 6 |
| 114 | The efficacies of entecavir and tenofovir in terms of enhancing prognosis after curative treatment of hepatitis B virus-related hepatocellular carcinoma. <i>European Journal of Internal Medicine</i> , 2021 , 89, 48-55 | 3.9 | 6 |
| 113 | Entecavir versus tenofovir in patients with chronic hepatitis B: Enemies or partners in the prevention of hepatocellular carcinoma. <i>Clinical and Molecular Hepatology</i> , 2021 , 27, 402-412 | 6.9 | 6 |
| 112 | Risk stratification using sarcopenia status among subjects with metabolic dysfunction-associated fatty liver disease. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2021 , 12, 1168-1178 | 10.3 | 6 |
| 111 | Association between the severity of liver fibrosis and cardiovascular outcomes in patients with type 2 diabetes. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021 , 36, 1703-1713 | 4 | 6 |
| 110 | Prognostic factors of sorafenib therapy in hepatocellular carcinoma patients with failure of transarterial chemoembolization. <i>Hepatology International</i> , 2017 , 11, 292-299 | 8.8 | 5 |
| 109 | Follow-Up Liver Stiffness Measurements after Liver Resection Influence Oncologic Outcomes of Hepatitis-B-Associated Hepatocellular Carcinoma with Liver Cirrhosis. <i>Cancers</i> , 2019 , 11, | 6.6 | 5 |
| 108 | Optimal criteria for hepatocellular carcinoma diagnosis using CT in patients undergoing liver transplantation. <i>European Radiology</i> , 2019 , 29, 1022-1031 | 8 | 5 |
| 107 | Graft function measured by transient elastography in living donor liver transplantation: preliminary. <i>Transplantation Proceedings</i> , 2013 , 45, 3028-31 | 1.1 | 5 |
| 106 | Durability of the virological response after lamivudine discontinuation in lamivudine-resistant patients with a complete virological response after lamivudine and adefovir combination therapy. <i>Journal of Medical Virology</i> , 2017 , 89, 85-90 | 19.7 | 5 |
| 105 | Inter-observer variability of response evaluation criteria for hepatocellular carcinoma treated with chemoembolization. <i>Digestive and Liver Disease</i> , 2015 , 47, 682-8 | 3.3 | 5 |
| 104 | Acoustic radiation force impulse elastography: a better option for patients with extrahepatic cholestasis. <i>European Journal of Gastroenterology and Hepatology</i> , 2012 , 24, 215-6 | 2.2 | 5 |
| 103 | Conditional survival estimate in patients with Barcelona Clinic Liver Cancer stage B/C hepatocellular carcinoma treated with hepatic arterial infusion chemotherapy with/without concurrent radiotherapy. <i>Oncotarget</i> , 2017 , 8, 79914-79926 | 3.3 | 5 |
| 102 | Fatty liver associated with metabolic derangement in patients with chronic kidney disease: A controlled attenuation parameter study. <i>Kidney Research and Clinical Practice</i> , 2017 , 36, 48-57 | 3.6 | 5 |

| | | | |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 101 | An artificial intelligence model to predict hepatocellular carcinoma risk in Korean and Caucasian patients with chronic hepatitis B. <i>Journal of Hepatology</i> , 2021 , | 13.4 | 5 |
| 100 | An observational study on long-term renal outcome in patients with chronic hepatitis B treated with tenofovir disoproxil fumarate. <i>Journal of Viral Hepatitis</i> , 2020 , 27, 316-322 | 3.4 | 5 |
| 99 | Sarcopenia assessed using bioimpedance analysis is associated independently with significant liver fibrosis in patients with chronic liver diseases. <i>European Journal of Gastroenterology and Hepatology</i> , 2020 , 32, 58-65 | 2.2 | 5 |
| 98 | Advanced Liver Fibrosis Predicts Unfavorable Long-Term Prognosis in First-Ever Ischemic Stroke or Transient Ischemic Attack. <i>Cerebrovascular Diseases</i> , 2020 , 49, 474-480 | 3.2 | 5 |
| 97 | Hepatocellular Carcinoma Risk According to Regimens for Eradication of Hepatitis C Virus; Interferon or Direct Acting Antivirals. <i>Cancers</i> , 2020 , 12, | 6.6 | 5 |
| 96 | Clinical Utility of a New Automated Hepatitis C Virus Core Antigen Assay for Prediction of Treatment Response in Patients with Chronic Hepatitis C. <i>Journal of Korean Medical Science</i> , 2016 , 31, 1431-7 | 4.7 | 5 |
| 95 | Switching from tenofovir and nucleoside analogue therapy to tenofovir monotherapy in virologically suppressed chronic hepatitis B patients with antiviral resistance. <i>Journal of Medical Virology</i> , 2018 , 90, 497-502 | 19.7 | 5 |
| 94 | Impact of tenofovir alafenamide vs. entecavir on hepatocellular carcinoma risk in patients with chronic hepatitis B. <i>Hepatology International</i> , 2021 , 15, 1083-1092 | 8.8 | 5 |
| 93 | Predictors of Discordance in the Assessment of Skeletal Muscle Mass between Computed Tomography and Bioimpedance Analysis. <i>Journal of Clinical Medicine</i> , 2019 , 8, | 5.1 | 4 |
| 92 | Refractoriness to transarterial chemoembolization in patients with recurrent hepatocellular carcinoma after curative resection. <i>PLoS ONE</i> , 2019 , 14, e0214613 | 3.7 | 4 |
| 91 | Non-O1, Non-O139Vibrio choleraeSepticemia after Acupuncture. <i>The Ewha Medical Journal</i> , 2013 , 36, S22 | 0.1 | 4 |
| 90 | The use of health functional foods in gastrointestinal cancer patients. <i>Clinical Nutrition Research</i> , 2013 , 2, 19-25 | 1.7 | 4 |
| 89 | Non-invasive assessment of liver fibrosis: the gap between ideal and real. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2011 , 26, 937-9 | 4 | 4 |
| 88 | Appraisal of Long-Term Outcomes of Liver-Directed Concurrent Chemoradiotherapy for Hepatocellular Carcinoma with Major Portal Vein Invasion. <i>Journal of Hepatocellular Carcinoma</i> , 2020 , 7, 403-412 | 5.3 | 4 |
| 87 | Survival Estimates after Stopping Sorafenib in Patients with Hepatocellular Carcinoma: NEXT Score Development and Validation. <i>Gut and Liver</i> , 2017 , 11, 693-701 | 4.8 | 4 |
| 86 | MRI Ancillary Features for LI-RADS Category 3 and 4 Observations: Improved Categorization to Indicate the Risk of Hepatic Malignancy. <i>American Journal of Roentgenology</i> , 2020 , 215, 1354-1362 | 5.4 | 4 |
| 85 | Cardiovascular Risk Is Elevated in Lean Subjects with Nonalcoholic Fatty Liver Disease. <i>Gut and Liver</i> , 2021 , | 4.8 | 4 |
| 84 | Validation of risk prediction scores for hepatocellular carcinoma in patients with chronic hepatitis B treated with entecavir or tenofovir. <i>Journal of Viral Hepatitis</i> , 2021 , 28, 95-104 | 3.4 | 4 |

| | | | |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 83 | Comparison of the current guidelines for diagnosing hepatocellular carcinoma using gadoxetic acid-enhanced magnetic resonance imaging. <i>European Radiology</i> , 2021 , 31, 4492-4503 | 8 | 4 |
| 82 | Disease duration and MedsgerQ severity score are associated with significant liver fibrosis in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, S68-74 | 2.2 | 4 |
| 81 | Compensated heart failure does not always alleviate overestimation of liver stiffness by transient elastography. <i>European Journal of Gastroenterology and Hepatology</i> , 2013 , 25, 385-7 | 2.2 | 3 |
| 80 | Predictors of refractory ascites development in patients with hepatitis B virus-related cirrhosis hospitalized to control ascitic decompensation. <i>Yonsei Medical Journal</i> , 2013 , 54, 145-53 | 3 | 3 |
| 79 | Dynamics of liver stiffness-based risk prediction model during antiviral therapy in patients with chronic hepatitis B. <i>European Journal of Gastroenterology and Hepatology</i> , 2021 , 33, 885-893 | 2.2 | 3 |
| 78 | A multi-centre study of trends in hepatitis B virus-related hepatocellular carcinoma risk over time during long-term entecavir therapy. <i>Journal of Viral Hepatitis</i> , 2020 , 27, 1352-1358 | 3.4 | 3 |
| 77 | Negligible risks of hepatocellular carcinoma during biomarker-defined immune-tolerant phase for patients with chronic hepatitis B. <i>Clinical and Molecular Hepatology</i> , 2021 , 27, 295-304 | 6.9 | 3 |
| 76 | Serum Dickkopf-1 in Combined with CA 19-9 as a Biomarker of Intrahepatic Cholangiocarcinoma. <i>Cancers</i> , 2021 , 13, | 6.6 | 3 |
| 75 | Predictive Performance of CAGE-B and SAGE-B Models in Asian Treatment-Naive Patients Who Started Entecavir for Chronic Hepatitis B. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , | 6.9 | 3 |
| 74 | Diagnostic Performance of Serum Asialo- α -acid Glycoprotein for Advanced Liver Fibrosis or Cirrhosis in Patients with Chronic Hepatitis B or Nonalcoholic Fatty Liver Disease. <i>Korean journal of gastroenterology = Taehan Sohwagi Hakhoe chi, The</i> , 2019 , 74, 341-348 | 0.6 | 3 |
| 73 | The Influence of Histologic Inflammation on the Improvement of Liver Stiffness Values Over 1 and 3 Years. <i>Journal of Clinical Medicine</i> , 2019 , 8, | 5.1 | 3 |
| 72 | Predictive validation of qualitative fibrosis staging in patients with chronic hepatitis B on antiviral therapy. <i>Scientific Reports</i> , 2019 , 9, 15628 | 4.9 | 3 |
| 71 | The LAW index as an accurate indicator of the initiation of antiviral treatment in patients with chronic hepatitis B. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 208-214 | 4 | 2 |
| 70 | The Paradoxical Protective Effect of Liver Steatosis on Severity and Functional Outcome of Ischemic Stroke. <i>Frontiers in Neurology</i> , 2019 , 10, 375 | 4.1 | 2 |
| 69 | Risk of Hepatocellular Carcinoma With Tenofovir vs Entecavir in Patients With Chronic Hepatitis B. <i>Gastroenterology</i> , 2020 , 158, 2310-2311 | 13.3 | 2 |
| 68 | Noninvasive Prediction of Erosive Esophagitis Using a Controlled Attenuation Parameter (CAP)-Based Risk Estimation Model. <i>Digestive Diseases and Sciences</i> , 2016 , 61, 507-16 | 4 | 2 |
| 67 | Partial virological response to adefovir add-on lamivudine rescue therapy in patients with lamivudine-resistant chronic hepatitis B. <i>Digestion</i> , 2013 , 87, 196-203 | 3.6 | 2 |
| 66 | Optimal timing of combining sorafenib with locoregional treatments in patients with advanced hepatocellular carcinoma. <i>European Journal of Gastroenterology and Hepatology</i> , 2013 , 25, 514-5 | 2.2 | 2 |

| | | | |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 65 | Advanced liver fibrosis measured by transient elastography predicts chronic kidney disease development in individuals with non-alcoholic fatty liver disease.. <i>Diabetologia</i> , 2021 , 65, 518 | 10.3 | 2 |
| 64 | Influence of Besifovir Dipivoxil Maleate Combined with L-Carnitine on Hepatic Steatosis in Patients with Chronic Hepatitis B. <i>Journal of Korean Medical Science</i> , 2020 , 35, e104 | 4.7 | 2 |
| 63 | Combined use of rosuvastatin and ezetimibe improves hepatic steatosis in patients with dyslipidemia. <i>European Journal of Gastroenterology and Hepatology</i> , 2020 , 32, 1538-1544 | 2.2 | 2 |
| 62 | Establishment and validation of a risk prediction model in patients with hepatocellular carcinoma treated with transarterial radioembolization. <i>European Journal of Gastroenterology and Hepatology</i> , 2020 , 32, 739-747 | 2.2 | 2 |
| 61 | Association between liver fibrosis and appendicular skeletal muscle mass during antiviral therapy in chronic hepatitis B. <i>Digestive and Liver Disease</i> , 2020 , 52, 1338-1345 | 3.3 | 2 |
| 60 | Reply to: "Response to A nomogram based on liver stiffness predicts postoperative complications in patients with hepatocellular carcinoma". <i>Journal of Hepatology</i> , 2020 , 73, 1270-1271 | 13.4 | 2 |
| 59 | Direct-Acting Antivirals Improve Treatment Outcomes in Patients with Hepatitis C Virus-Related Hepatocellular Carcinoma Treated with Transarterial Chemoembolization: A Nationwide, Multi-center, Retrospective Cohort Study. <i>Digestive Diseases and Sciences</i> , 2021 , 66, 2427-2438 | 4 | 2 |
| 58 | External validation of CAGE-B and SAGE-B scores for Asian chronic hepatitis B patients with well-controlled viremia by antivirals. <i>Journal of Viral Hepatitis</i> , 2021 , 28, 951-958 | 3.4 | 2 |
| 57 | Clinical predictors of silent but substantial liver fibrosis in primary Sjogren's syndrome. <i>Modern Rheumatology</i> , 2016 , 26, 576-82 | 3.3 | 2 |
| 56 | Evolution of liver fibrosis and steatosis markers in patients with type 2 diabetes after metformin treatment for 2 years. <i>Journal of Diabetes and Its Complications</i> , 2021 , 35, 107747 | 3.2 | 2 |
| 55 | Impact of antiviral therapy on risk prediction model for hepatocellular carcinoma development in patients with chronic hepatitis B. <i>Hepatology Research</i> , 2021 , 51, 406-416 | 5.1 | 2 |
| 54 | Noninvasive evaluation of liver fibrosis: comparison of the stretched exponential diffusion-weighted model to other diffusion-weighted MRI models and transient elastography. <i>European Radiology</i> , 2021 , 31, 4813-4823 | 8 | 2 |
| 53 | Effect of tenofovir alafenamide vs. tenofovir disoproxil fumarate on hepatocellular carcinoma risk in chronic hepatitis B. <i>Journal of Viral Hepatitis</i> , 2021 , 28, 1570-1578 | 3.4 | 2 |
| 52 | Episodic Detectable Viremia Does Not Affect Prognosis in Untreated Compensated Cirrhosis With Serum Hepatitis B Virus DNA . <i>American Journal of Gastroenterology</i> , 2021 , | 0.7 | 2 |
| 51 | Addition of liver stiffness enhances the predictive accuracy of the PAGE-B model for hepatitis B-related hepatocellular carcinoma. <i>Alimentary Pharmacology and Therapeutics</i> , 2021 , 53, 919-927 | 6.1 | 2 |
| 50 | Entecavir and tenofovir on renal function in patients with hepatitis B virus-related hepatocellular carcinoma. <i>Journal of Viral Hepatitis</i> , 2020 , 27, 932-940 | 3.4 | 1 |
| 49 | Clinical relevance of dynamic risk assessment for developing hepatocellular carcinoma during prolonged antiviral therapy. <i>Journal of Hepatology</i> , 2020 , 73, 222-223 | 13.4 | 1 |
| 48 | Radiologic Evaluation of Non-Alcoholic Fatty Liver Disease in Diabetic Patient. <i>Journal of Korean Diabetes</i> , 2017 , 18, 88 | 0.1 | 1 |

| | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 47 | Reply to "The problem of the most appropriate curative treatment for hepatocellular carcinoma. When to embolize? When to operate?". <i>Journal of Hepatology</i> , 2015 , 63, 281-2 | 13.4 | 1 |
| 46 | Acoustic radiation force impulse elastography is useful to exclude nonliver-related ascites. <i>European Journal of Gastroenterology and Hepatology</i> , 2011 , 23, 1080-1 | 2.2 | 1 |
| 45 | Correlation of the grade of hepatic steatosis between controlled attenuation parameter and ultrasound in patients with fatty liver: a multi-center retrospective cohort study. <i>Korean Journal of Internal Medicine</i> , 2020 , 35, 1346-1353 | 2.5 | 1 |
| 44 | Efficacy of entecavir, tenofovir disoproxil fumarate, and tenofovir alafenamide in treatment-naïve hepatitis B patients. <i>Hepatology International</i> , 2021 , 15, 1328-1336 | 8.8 | 1 |
| 43 | Proposal of a New Risk Score for Patients Treated with Transarterial Chemoembolization due to Recurrent Hepatocellular Carcinoma after Curative Resection: A Multicenter Study. <i>Gut and Liver</i> , 2020 , 14, 477-485 | 4.8 | 1 |
| 42 | Multicenter Retrospective Risk Assessment of Esophageal Variceal Bleeding in Patients with Cirrhosis: An Acoustic Radiation Force Impulse Elastography-Based Prediction Model. <i>Gut and Liver</i> , 2019 , 13, 206-214 | 4.8 | 1 |
| 41 | Letter: risk of hepatocellular carcinoma in immune-tolerant phase of chronic hepatitis B-authorsQ reply. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 52, 913-914 | 6.1 | 1 |
| 40 | Treatment efficacy by hepatic arterial infusion chemotherapy vs. sorafenib after liver-directed concurrent chemoradiotherapy for advanced hepatocellular carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021 , 147, 3123-3133 | 4.9 | 1 |
| 39 | Change in the Recurrence Pattern and Predictors over Time after Complete Cure of Hepatocellular Carcinoma. <i>Gut and Liver</i> , 2021 , 15, 420-429 | 4.8 | 1 |
| 38 | Subclinical Ascites Does Not Affect the Long-term Prognosis in Hepatitis B Virus-related Cirrhosis Patients Receiving Antivirals. <i>Journal of Clinical Gastroenterology</i> , 2016 , 50, 676-85 | 3 | 1 |
| 37 | Fibrosis-matched outcomes between chronic hepatitis B patients with drug-induced virological response and inactive carriers. <i>Liver International</i> , 2019 , 39, 81-89 | 7.9 | 1 |
| 36 | Prognostic Value of Alpha-Fetoprotein in Patients Who Achieve a Complete Response to Transarterial Chemoembolization for Hepatocellular Carcinoma. <i>Yonsei Medical Journal</i> , 2021 , 62, 12-20 ³ | | 1 |
| 35 | Predictors of Complete Response in Patients with Hepatocellular Carcinoma Treated with Trans-Arterial Radioembolization. <i>Current Oncology</i> , 2021 , 28, 965-977 | 2.8 | 1 |
| 34 | A Case of Lymphocyte-Rich Hepatocellular Carcinoma in a Patient Who Was Treated for Colon Cancer. <i>Journal of Liver Cancer</i> , 2021 , 21, 69-75 | 0.7 | 1 |
| 33 | Pharmacologic Treatment Using Non-Hypoglycemic Agents for Nonalcoholic Fatty Liver Disease. <i>Journal of Korean Diabetes</i> , 2018 , 19, 97 | 0.1 | 1 |
| 32 | A qualitative study of self-management experiences in people with non-alcoholic fatty liver disease. <i>Nursing Open</i> , 2021 , 8, 3135-3142 | 2.1 | 1 |
| 31 | Long-term effects of entecavir and tenofovir treatment on the fibrotic burden in patients with chronic hepatitis B. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021 , | 4 | 1 |
| 30 | Impact of HBeAg on Hepatocellular Carcinoma Risk During Oral Antiviral Treatment in Patients With Chronic Hepatitis B. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , | 6.9 | 1 |

| | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 29 | Metabolic Dysfunction-Associated Fatty Liver Disease and Risk of Incident Chronic Kidney Disease: A Nationwide Cohort Study.. <i>Diabetes and Metabolism</i> , 2022 , 101344 | 5.4 | 1 |
| 28 | Screening, confirmation, and treatment rates of hepatitis C virus infection in a tertiary academic medical center in South Korea. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021 , 36, 2479-2485 | 4.85 | 0 |
| 27 | The Prognostic Role of On-Treatment Liver Stiffness for Hepatocellular Carcinoma Development in Patients with Chronic Hepatitis B. <i>Journal of Hepatocellular Carcinoma</i> , 2021 , 8, 467-476 | 5.3 | 0 |
| 26 | Development and initial validation of the nonalcoholic fatty liver disease self-management questionnaire. <i>Research in Nursing and Health</i> , 2021 , 44, 844-853 | 2 | 0 |
| 25 | Letter to the Editor: Re: Tenofovir Versus Entecavir on Recurrence of Hepatitis B Virus-Related Hepatocellular Carcinoma After Surgical Resection. <i>Hepatology</i> , 2021 , 73, 466 | 11.2 | 0 |
| 24 | Association Between Curative Treatment after Transarterial Radioembolization and Better Survival Outcomes in Patients with Hepatocellular Carcinoma. <i>Cancer Investigation</i> , 2021 , 39, 274-283 | 2.1 | 0 |
| 23 | Controlled attenuation parameter value and the risk of hepatocellular carcinoma in chronic hepatitis B patients under antiviral therapy. <i>Hepatology International</i> , 2021 , 15, 892-900 | 8.8 | 0 |
| 22 | Risk assessment of hepatocellular carcinoma and liver-related events using ultrasonography and transient elastography in patients with chronic hepatitis B. <i>Journal of Viral Hepatitis</i> , 2021 , 28, 1362-1372 | 3.4 | 0 |
| 21 | No influence of hepatic steatosis on the 3-year outcomes of patients with quiescent chronic hepatitis B. <i>Journal of Viral Hepatitis</i> , 2021 , 28, 1545-1553 | 3.4 | 0 |
| 20 | Revised Korean Antiviral Guideline Reduces the Hepatitis B-related Hepatocellular Carcinoma Risk in Cirrhotic Patients. <i>Journal of Korean Medical Science</i> , 2021 , 36, e105 | 4.7 | 0 |
| 19 | Noninvasive surrogates are poor predictors of liver fibrosis in patients with Fontan circulation.. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2021 , | 1.5 | 0 |
| 18 | Comparison of FibroScan-Aspartate Aminotransferase (FAST) Score and Other Non-invasive Surrogates in Predicting High-Risk Non-alcoholic Steatohepatitis Criteria.. <i>Frontiers in Medicine</i> , 2022 , 9, 869190 | 4.9 | 0 |
| 17 | Liver-Directed Concurrent Chemoradiotherapy versus Sorafenib in Hepatocellular Carcinoma with Portal Vein Tumor Thrombosis. <i>Cancers</i> , 2022 , 14, 2396 | 6.6 | 0 |
| 16 | Fibrotic burden during antiviral therapy for chronic hepatitis B, not ALT level, independently predicts liver cancer risk.. <i>Liver International</i> , 2022 , | 7.9 | 0 |
| 15 | The effect of pharmacological treatment and lifestyle modification in patients with nonalcoholic fatty liver disease: An umbrella review of meta-analyses of randomized controlled trials.. <i>Obesity Reviews</i> , 2022 , e13464 | 10.6 | 0 |
| 14 | Exercise Reduces the Risk of Chronic Kidney Disease in Individuals with Nonalcoholic Fatty Liver Disease: A Nationwide Cohort Study. <i>Diabetes and Metabolism</i> , 2022 , 101362 | 5.4 | 0 |
| 13 | Reply to "Response assessment for HCC patients treated with repeated TACE: The optimal time-point is still an open issue". <i>Journal of Hepatology</i> , 2015 , 63, 1532 | 13.4 | |
| 12 | Can Non-invasive Fibrosis Assessment Predict Sustained Virological Response to Telaprevir with Pegylated Interferon and Ribavirin in Chronic Hepatitis C Patients?. <i>Korean journal of gastroenterology = Taehan Sohwagi Hakhoe chi, The</i> , 2014 , 64, 250 | 0.6 | |

- 11 Clinical Implications of Shear Wave Elastography in Patients with Chronic Hepatitis B. *Journal of Clinical Ultrasound*, **2018**, 3, 45-49 0
- 10 AuthorsQresponse: Additional more factors should be included in predicting risk of hepatocellular carcinoma in patients with chronic hepatitis B virus infection. *Journal of Viral Hepatitis*, **2021**, 28, 574 3.4
- 9 Efficacy and safety of localized concurrent chemoradiation therapy and sorafenib sequential therapy in advanced hepatocellular carcinoma: A prospective phase II trial.. *Journal of Clinical Oncology*, **2017**, 35, e15678-e15678 2.2
- 8 Reply to: "Liver stiffness: A novel predictor of postoperative complications in patients with hepatocellular carcinoma". *Journal of Hepatology*, **2020**, 73, 988-989 13.4
- 7 Predictors of Sarcopenia in an Obese Asian Population. *Nutrition and Cancer*, **2021**, 1-10 2.8
- 6 Reply to Qrognostic model for patients receiving arterial-embolization for hepatocellular carcinoma: issues to considerQLiver International, **2016**, 36, 312 7.9
- 5 Reply. *Hepatology*, **2016**, 63, 672-3 11.2
- 4 Advanced Stage Hepatocellular Carcinoma Successfully Treated with Liver-directed Concurrent Chemoradiotherapy and Sequential Transarterial Radio-embolization. *Journal of Liver Cancer*, **2021**, 21, 97-103 0.7
- 3 Prevalence and Risk Factors of Cardiovascular Disease in Patients with Chronic Hepatitis B. *Digestive Diseases and Sciences*, **2021**, 1 4
- 2 Dickkopf-1 promotes angiogenesis by upregulating VEGF receptor 2-mediated mTOR/p70S6K signaling in hepatocellular carcinoma. *American Journal of Cancer Research*, **2021**, 11, 4788-4806 4.4
- 1 Letter: fatty liver disease could have been a confounding factor for phase change in patients with chronic hepatitis B in the immune-tolerant phase-authorsQreply. *Alimentary Pharmacology and Therapeutics*, **2020**, 52, 1094-1095 6.1