

Sarah Tosato

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7020666/sarah-tosato-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75
papers

7,353
citations

28
h-index

85
g-index

86
ext. papers

9,064
ext. citations

7.7
avg, IF

4.26
L-index

#	Paper	IF	Citations
75	Use of multiple polygenic risk scores for distinguishing schizophrenia-spectrum disorder and affective psychosis categories in a first-episode sample; the EU-GEI study.. <i>Psychological Medicine</i> , 2022 , 1-10	6.9	2
74	Facial Emotion Recognition in Psychosis and Associations With Polygenic Risk for Schizophrenia: Findings From the Multi-Center EU-GEI Case-Control Study.. <i>Schizophrenia Bulletin</i> , 2022 ,	1.3	1
73	Three-Month Follow-up Study of Mental Health Outcomes After a National COVID-19 Lockdown: Comparing Patients With Mood or Anxiety Disorders Living in an Area With a Higher Versus Lower Infection Incidence.. <i>Journal of Clinical Psychiatry</i> , 2022 , 83,	4.6	3
72	Mapping genomic loci implicates genes and synaptic biology in schizophrenia.. <i>Nature</i> , 2022 ,	50.4	35
71	Targeting Metabolic Abnormalities in Mental Health Prevention Strategies 2022 , 225-249		
70	Pre-training inter-rater reliability of clinical instruments in an international psychosis research project. <i>Schizophrenia Research</i> , 2021 , 230, 104-107	3.6	5
69	Perceived major experiences of discrimination, ethnic group, and risk of psychosis in a six-country case-control study. <i>Psychological Medicine</i> , 2021 , 1-9	6.9	3
68	A Comparison of Ten Polygenic Score Methods for Psychiatric Disorders Applied Across Multiple Cohorts. <i>Biological Psychiatry</i> , 2021 , 90, 611-620	7.9	17
67	Lack of Support for the Genes by Early Environment Interaction Hypothesis in the Pathogenesis of Schizophrenia. <i>Schizophrenia Bulletin</i> , 2021 ,	1.3	6
66	The Independent Effects of Psychosocial Stressors on Subclinical Psychosis: Findings From the Multinational EU-GEI Study. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1674-1684	1.3	3
65	Childhood and Adulthood Severe Stressful Experiences and Biomarkers Related to Glucose Metabolism: A Possible Association?. <i>Frontiers in Psychiatry</i> , 2021 , 12, 629137	5	1
64	Duration of Untreated Psychosis in First-Episode Psychosis is not Associated With Common Genetic Variants for Major Psychiatric Conditions: Results From the Multi-Center EU-GEI Study. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1653-1662	1.3	0
63	Jumping to conclusions, general intelligence, and psychosis liability: findings from the multi-centre EU-GEI case-control study. <i>Psychological Medicine</i> , 2021 , 51, 623-633	6.9	15
62	Migration history and risk of psychosis: results from the multinational EU-GEI study. <i>Psychological Medicine</i> , 2021 , 1-13	6.9	6
61	The continuity of effect of schizophrenia polygenic risk score and patterns of cannabis use on transdiagnostic symptom dimensions at first-episode psychosis: findings from the EU-GEI study. <i>Translational Psychiatry</i> , 2021 , 11, 423	8.6	1
60	Depression is associated with increased disease activity and higher disability in a large Italian cohort of patients with rheumatoid arthritis. <i>Advances in Rheumatology</i> , 2021 , 61, 57	3	2
59	The relationship of symptom dimensions with premorbid adjustment and cognitive characteristics at first episode psychosis: Findings from the EU-GEI study. <i>Schizophrenia Research</i> , 2021 , 236, 69-79	3.6	0

58	Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. <i>Molecular Psychiatry</i> , 2021 , 26, 4529-4543	15.1	7
57	The incidence of psychotic disorders among migrants and minority ethnic groups in Europe: findings from the multinational EU-GEI study. <i>Psychological Medicine</i> , 2020 , 1-10	6.9	7
56	Daily use of high-potency cannabis is associated with more positive symptoms in first-episode psychosis patients: the EU-GEI case-control study. <i>Psychological Medicine</i> , 2020 , 1-9	6.9	13
55	Childhood trauma and glucose metabolism in patients with first-episode psychosis. <i>Psychoneuroendocrinology</i> , 2020 , 113, 104536	5	10
54	Dietary habits and physical activity in first-episode psychosis patients treated in community services. Effect on early anthropometric and cardio-metabolic alterations. <i>Schizophrenia Research</i> , 2020 , 216, 374-381	3.6	2
53	Correlations between immune and metabolic serum markers and schizophrenia/bipolar disorder polygenic risk score in first-episode psychosis. <i>Microbial Biotechnology</i> , 2020 , 14, 507-511	3.3	9
52	Premorbid Adjustment and IQ in Patients With First-Episode Psychosis: A Multisite Case-Control Study of Their Relationship With Cannabis Use. <i>Schizophrenia Bulletin</i> , 2020 , 46, 517-529	1.3	10
51	Pathways to care, DUP, and types of interventions over 5 years following psychosis onset: findings from a naturalistic study conducted in routine generalist mental health services. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2020 , 55, 175-186	4.5	1
50	The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): a multicentre case-control study. <i>Lancet Psychiatry</i> , 2019 , 6, 427-436	23.3	322
49	Psychiatric Disorders 2019 , 215-235		
48	Dimensional structure of first episode psychosis. <i>Microbial Biotechnology</i> , 2019 , 13, 1431-1438	3.3	13
47	The impact of gender and childhood abuse on age of psychosis onset, psychopathology and needs for care in psychosis patients. <i>Schizophrenia Research</i> , 2019 , 210, 164-171	3.6	9
46	Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. <i>Nature Genetics</i> , 2018 , 50, 381-389	36.3	787
45	Immune and metabolic alterations in first episode psychosis (FEP) patients. <i>Brain, Behavior, and Immunity</i> , 2018 , 70, 315-324	16.6	22
44	Prevalence of non-psychotic disorders in ultra-high risk individuals and transition to psychosis: A systematic review. <i>Psychiatry Research</i> , 2018 , 270, 1-12	9.9	23
43	Classification of first-episode psychosis in a large cohort of patients using support vector machine and multiple kernel learning techniques. <i>NeuroImage</i> , 2017 , 145, 238-245	7.9	36
42	Predictors of 9-month hospitalization in patients with first-episode affective and non-affective psychosis. Results from the GET UP pragmatic cluster randomized controlled trial. <i>Schizophrenia Research</i> , 2017 , 190, 187-188	3.6	4
41	Predictors and moderators of treatment outcome in patients receiving multi-element psychosocial intervention for early psychosis: results from the GET UP pragmatic cluster randomised controlled trial. <i>British Journal of Psychiatry</i> , 2017 , 210, 342-349	5.4	19

40	Epigenetics and gene expression profile in first-episode psychosis: The role of childhood trauma. <i>Neuroscience and Biobehavioral Reviews</i> , 2017 , 83, 226-237	9	32
39	Family Burden, Emotional Distress and Service Satisfaction in First Episode Psychosis. Data from the GET UP Trial. <i>Frontiers in Psychology</i> , 2017 , 8, 721	3.4	9
38	A Systematized Review of Atypical Antipsychotics in Pregnant Women: Balancing Between Risks of Untreated Illness and Risks of Drug-Related Adverse Effects. <i>Journal of Clinical Psychiatry</i> , 2017 , 78, e477-489	4.6	38
37	Association of AADAC Deletion and Gilles de la Tourette Syndrome in a Large European Cohort. <i>Biological Psychiatry</i> , 2016 , 79, 383-391	7.9	26
36	The use of dynamic susceptibility contrast (DSC) MRI to automatically classify patients with first episode psychosis. <i>Schizophrenia Research</i> , 2015 , 165, 38-44	3.6	18
35	Classification of first-episode psychosis: a multi-modal multi-feature approach integrating structural and diffusion imaging. <i>Journal of Neural Transmission</i> , 2015 , 122, 897-905	4.3	20
34	Association between the COMT gene and neurological abnormalities and poorer executive function in psychosis. <i>Psychiatry Research</i> , 2015 , 230, 742-3	9.9	2
33	Feasibility and Effectiveness of a Multi-Element Psychosocial Intervention for First-Episode Psychosis: Results From the Cluster-Randomized Controlled GET UP PIANO Trial in a Catchment Area of 10 Million Inhabitants. <i>Schizophrenia Bulletin</i> , 2015 , 41, 1192-203	1.3	78
32	Bridging the gap between education and appropriate use of benzodiazepines in psychiatric clinical practice. <i>Neuropsychiatric Disease and Treatment</i> , 2015 , 11, 1885-909	3.1	37
31	Neurological soft signs and prepulse inhibition of the startle reflex in psychosis: A pilot study. <i>International Journal of Psychiatry in Clinical Practice</i> , 2015 , 19, 276-84	2.4	
30	First-contact incidence of psychosis in north-eastern Italy: influence of age, gender, immigration and socioeconomic deprivation. <i>British Journal of Psychiatry</i> , 2014 , 205, 127-34	5.4	40
29	A genome-wide association analysis of a broad psychosis phenotype identifies three loci for further investigation. <i>Biological Psychiatry</i> , 2014 , 75, 386-97	7.9	36
28	Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. <i>Schizophrenia Bulletin</i> , 2014 , 40, 729-36	1.3	186
27	Positive symptoms in first-episode psychosis patients experiencing low maternal care and stressful life events: a pilot study to explore the role of the COMT gene. <i>Stress</i> , 2014 , 17, 410-5	3	15
26	No association between NRG1 and ErbB4 genes and psychopathological symptoms of schizophrenia. <i>NeuroMolecular Medicine</i> , 2014 , 16, 742-51	4.6	4
25	The role of experienced and anticipated discrimination in the lives of people with first-episode psychosis. <i>Psychiatric Services</i> , 2014 , 65, 1034-40	3.3	27
24	Genome-wide association analysis identifies 13 new risk loci for schizophrenia. <i>Nature Genetics</i> , 2013 , 45, 1150-9	36.3	1153
23	The impact of cannabis use on age of onset and clinical characteristics in first-episode psychotic patients. Data from the Psychosis Incident Cohort Outcome Study (PICOS). <i>Journal of Psychiatric Research</i> , 2013 , 47, 438-44	5.2	54

22	COMT, neuropsychological function and brain structure in schizophrenia: a systematic review and neurobiological interpretation. <i>Journal of Psychiatry and Neuroscience</i> , 2013 , 38, 366-80	4.5	59
21	Association study of nonsynonymous single nucleotide polymorphisms in schizophrenia. <i>Biological Psychiatry</i> , 2012 , 71, 169-77	7.9	63
20	A multi-element psychosocial intervention for early psychosis (GET UP PIANO TRIAL) conducted in a catchment area of 10 million inhabitants: study protocol for a pragmatic cluster randomized controlled trial. <i>Trials</i> , 2012 , 13, 73	2.8	38
19	Replication study and meta-analysis in European samples supports association of the 3p21.1 locus with bipolar disorder. <i>Biological Psychiatry</i> , 2012 , 72, 645-50	7.9	15
18	Is neuregulin 1 involved in determining cerebral volumes in schizophrenia? Preliminary results showing a decrease in superior temporal gyrus volume. <i>Neuropsychobiology</i> , 2012 , 65, 119-25	4	22
17	At-risk variant in TCF7L2 for type II diabetes increases risk of schizophrenia. <i>Biological Psychiatry</i> , 2011 , 70, 59-63	7.9	101
16	Common variants on 8p12 and 1q24.2 confer risk of schizophrenia. <i>Nature Genetics</i> , 2011 , 43, 1224-7	36.3	201
15	Effect of COMT genotype on aggressive behaviour in a community cohort of schizophrenic patients. <i>Neuroscience Letters</i> , 2011 , 495, 17-21	3.3	29
14	Common variants at VRK2 and TCF4 conferring risk of schizophrenia. <i>Human Molecular Genetics</i> , 2011 , 20, 4076-81	5.6	162
13	Maternally derived microduplications at 15q11-q13: implication of imprinted genes in psychotic illness. <i>American Journal of Psychiatry</i> , 2011 , 168, 408-17	11.9	84
12	A large replication study and meta-analysis in European samples provides further support for association of AH11 markers with schizophrenia. <i>Human Molecular Genetics</i> , 2010 , 19, 1379-86	5.6	42
11	Disruption of the neurexin 1 gene is associated with schizophrenia. <i>Human Molecular Genetics</i> , 2009 , 18, 988-96	5.6	376
10	The contribution of epidemiology to defining the most appropriate approach to genetic research on schizophrenia. <i>Epidemiologia E Psichiatria Sociale</i> , 2009 , 18, 81-90		15
9	Common variants conferring risk of schizophrenia. <i>Nature</i> , 2009 , 460, 744-7	50.4	1350
8	Can the role of genetic factors in schizophrenia be enlightened by studies of candidate gene mutant mice behaviour?. <i>World Journal of Biological Psychiatry</i> , 2009 , 10, 778-97	3.8	8
7	Large recurrent microdeletions associated with schizophrenia. <i>Nature</i> , 2008 , 455, 232-6	50.4	1427
6	Association study of dysbindin gene with clinical and outcome measures in a representative cohort of Italian schizophrenic patients. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2007 , 144B, 647-59	3.5	43
5	The psychopathology of schizophrenia and the presence of neurological soft signs: a review. <i>Current Opinion in Psychiatry</i> , 2005 , 18, 285-8	4.9	41

4	Association between the neuregulin 1 gene and schizophrenia: a systematic review. <i>Schizophrenia Bulletin</i> , 2005, 31, 613-7	1.3	97
3	The independent and combined influence of schizophrenia polygenic risk score and heavy cannabis use on risk for psychotic disorder: A case-control analysis from the EUGEI study.		2
2	Implementing evidence-based treatments in routine mental health services57-77		
1	Synergistic effects of childhood adversity and polygenic risk in first-episode psychosis: the EU-GEI study. <i>Psychological Medicine</i> ,1-9	6.9	1