

Tatiane C Mariguela

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7027812/publications.pdf>

Version: 2024-02-01

12
papers

545
citations

933264

10
h-index

1199470

12
g-index

12
all docs

12
docs citations

12
times ranked

544
citing authors

#	ARTICLE	IF	CITATIONS
1	Phylogeny and biogeography of Triportheidae (Teleostei: Characiformes) based on molecular data. <i>Molecular Phylogenetics and Evolution</i> , 2016, 96, 130-139.	1.2	21
2	Systematic and historical biogeography of the Bryconidae (Ostariophysi: Characiformes) suggesting a new rearrangement of its genera and an old origin of Mesoamerican ichthyofauna. <i>BMC Evolutionary Biology</i> , 2014, 14, 152.	3.2	32
3	Multilocus molecular phylogeny of Gasteropelecidae (Ostariophysi: Characiformes) reveals the existence of an unsuspected diversity. <i>Molecular Phylogenetics and Evolution</i> , 2013, 69, 1209-1214.	1.2	10
4	Composition and interrelationships of a large Neotropical freshwater fish group, the subfamily Cheirodontinae (Characiformes: Characidae): A case study based on mitochondrial and nuclear DNA sequences. <i>Molecular Phylogenetics and Evolution</i> , 2013, 68, 23-34.	1.2	18
5	Molecular phylogeny of <i>Moenkhausia</i> (Characidae) inferred from mitochondrial and nuclear DNA evidence. <i>Journal of Zoological Systematics and Evolutionary Research</i> , 2013, 51, 327-332.	0.6	21
6	Historical biogeography and cryptic diversity in the Callichthyinae (Siluriformes, Callichthyidae). <i>Journal of Zoological Systematics and Evolutionary Research</i> , 2013, 51, 308-315.	0.6	18
7	Distributions and phylogeographic data of rheophilic freshwater fishes provide evidences on the geographic extension of a central-brazilian amazonian palaeoplateau in the area of the present day Pantanal Wetland. <i>Neotropical Ichthyology</i> , 2013, 11, 319-326.	0.5	24
8	A new species of <i>Tetragonopterus</i> Cuvier, 1816 (Characiformes: Characidae: Tetragonopterinae) from the rio Jari, Amapá, northern Brazil. <i>Neotropical Ichthyology</i> , 2011, 9, 49-56.	0.5	61
9	5S rDNA chromosomal mapping and COI sequence analysis reveal differentiation among distinct populations of a characid fish <i>Serrapinnus notomelas</i> . <i>Reviews in Fish Biology and Fisheries</i> , 2011, 21, 779-788.	2.4	3
10	Phylogenetic relationships within the speciose family Characidae (Teleostei: Ostariophysi): <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 387 Td Biology</i> , 2011, 11, 275.	3.2	270
11	Identification of guitarfish species <i>Rhinobatos</i> <i>percellens</i> , <i>R. horkelli</i> , and <i>Zapteryx</i> <i>abrevirostris</i> (Chondrichthyes) using mitochondrial genes and RFLP technique. <i>Conservation Genetics Resources</i> , 2009, 1, 393-396.	0.4	7
12	New species of <i>Moenkhausia</i> Eigenmann, 1903 (Characiformes: Characidae) with comments on the <i>Moenkhausia oligolepis</i> species complex. <i>Neotropical Ichthyology</i> , 2009, 7, 161-168.	0.5	60