

# Wioleta Chajeka-Wierzchowska

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7034787/publications.pdf>

Version: 2024-02-01

34  
papers

660  
citations

567281

15  
h-index

580821

25  
g-index

36  
all docs

36  
docs citations

36  
times ranked

791  
citing authors

#	ARTICLE	IF	CITATIONS
1	Occurrence of antibiotic resistance among <i>Enterobacteriales</i> isolated from raw and ready-to-eat food – phenotypic and genotypic characteristics. <i>International Journal of Environmental Health Research</i> , 2022, 32, 1733-1744.	2.7	3
2	Microorganisms from starter and protective cultures - Occurrence of antibiotic resistance and conjugal transfer of tet genes in vitro and during food fermentation. <i>LWT - Food Science and Technology</i> , 2022, 153, 112490.	5.2	15
3	Effects of osmotic and high pressure stress on expression of virulence factors among <i>Enterococcus</i> spp. isolated from food of animal origin. <i>Food Microbiology</i> , 2022, 102, 103900.	4.2	16
4	Antimicrobial and Antibiotic Resistance from the Perspective of Polish Veterinary Students: An Inter-University Study. <i>Antibiotics</i> , 2022, 11, 115.	3.7	9
5	A Comparison of Methods for Identifying <i>Enterobacteriales</i> Isolates from Fish and Prawns. <i>Pathogens</i> , 2022, 11, 410.	2.8	15
6	Ceviche-Natural Preservative: Possibility of Microbiota Survival and Effect on <i>L. monocytogenes</i> . <i>Foods</i> , 2022, 11, 860.	4.3	4
7	Linezolid-Resistant <i>Enterococcus</i> spp. Isolates from Foods of Animal Origin – The Genetic Basis of Acquired Resistance. <i>Foods</i> , 2022, 11, 975.	4.3	5
8	Antibiotic Resistance Carriage Causes a Lower Survivability Due to Stress Associated with High-Pressure Treatment among Strains from Starter Cultures. <i>Animals</i> , 2022, 12, 1460.	2.3	3
9	High pressure processing, acidic and osmotic stress increased resistance to aminoglycosides and tetracyclines and the frequency of gene transfer among strains from commercial starter and protective cultures. <i>Food Microbiology</i> , 2022, 107, 104090.	4.2	8
10	<i>Enterococci</i> isolated from plant-derived food - Analysis of antibiotic resistance and the occurrence of resistance genes. <i>LWT - Food Science and Technology</i> , 2021, 139, 110549.	5.2	7
11	In-milk inactivation of <i>Escherichia coli</i> O157:H7 by the environmental lytic bacteriophage ECPS-6. <i>Journal of Food Safety</i> , 2020, 40, e12747.	2.3	18
12	Enterotoxigenic Potential of Coagulase-Negative Staphylococci from Ready-to-Eat Food. <i>Pathogens</i> , 2020, 9, 734.	2.8	29
13	Biofilm Formation Ability and Presence of Adhesion Genes among Coagulase-Negative and Coagulase-Positive Staphylococci Isolates from Raw Cow's Milk. <i>Pathogens</i> , 2020, 9, 654.	2.8	33
14	Virulence Characterization of <i>Listeria monocytogenes</i> , <i>Listeria innocua</i> , and <i>Listeria welshimeri</i> Isolated from Fish and Shrimp Using In Vivo Early Zebrafish Larvae Models and Molecular Study. <i>Pathogens</i> , 2020, 9, 1028.	2.8	6
15	Ready-to-eat dairy products as a source of multidrug-resistant <i>Enterococcus</i> strains: Phenotypic and genotypic characteristics. <i>Journal of Dairy Science</i> , 2020, 103, 4068-4077.	3.4	23
16	Starter cultures as a reservoir of antibiotic resistant microorganisms. <i>LWT - Food Science and Technology</i> , 2020, 127, 109424.	5.2	31
17	Mleko i produkty mleczne jako potencjalne Źródło enterotoksyn gronkowcowych. <i>Przemysł Spożywczy</i> , 2020, 1, 26-29.	0.1	0
18	<i>Enterococci</i> from ready-to-eat food – horizontal gene transfer of antibiotic resistance genes and genotypic characterization by PCR melting profile. <i>Journal of the Science of Food and Agriculture</i> , 2019, 99, 1172-1179.	3.5	22

#	ARTICLE	IF	CITATIONS
19	<i>S. epidermidis</i> strains from artisanal cheese made from unpasteurized milk in Poland - Genetic characterization of antimicrobial resistance and virulence determinants. <i>International Journal of Food Microbiology</i> , 2019, 294, 55-59.	4.7	17
20	Bakterie fermentacji mlekowej w tym szczepy probiotyczne jako rezerwuar genów w oporności na antybiotyki. <i>Zywność</i> , 2019, 120, 22-35.	0.1	0
21	Virulence factors of <i>Enterococcus</i> spp. presented in food. <i>LWT - Food Science and Technology</i> , 2017, 75, 670-676.	5.2	68
22	Prevalence, biofilm formation and virulence markers of <i>Salmonella</i> sp. and <i>Yersinia enterocolitica</i> in food of animal origin in Poland. <i>LWT - Food Science and Technology</i> , 2017, 75, 552-556.	5.2	28
23	Diversity of Antibiotic Resistance Genes in <i>Enterococcus</i> Strains Isolated from Ready-to-Eat Meat Products. <i>Journal of Food Science</i> , 2016, 81, M2799-M2807.	3.1	38
24	Virulence factors, antimicrobial resistance and biofilm formation in <i>Enterococcus</i> spp. isolated from retail shrimps. <i>LWT - Food Science and Technology</i> , 2016, 69, 117-122.	5.2	49
25	Growth potential of <i>Yersinia enterocolitica</i> in blue cheese and in blue cheese with probiotic <i>Lactobacillus acidophilus</i> LA-5®. <i>Journal of Food Science and Technology</i> , 2015, 52, 7540-7544.	2.8	7
26	Coagulase-negative staphylococci (CoNS) isolated from ready-to-eat food of animal origin – Phenotypic and genotypic antibiotic resistance. <i>Food Microbiology</i> , 2015, 46, 222-226.	4.2	93
27	Microbiological contamination of dried and lyophilized garlic as a potential source of food spoilage. <i>Journal of Food Science and Technology</i> , 2015, 52, 1802-1807.	2.8	9
28	Retail Ready-to-Eat Food as a Potential Vehicle for <i>Staphylococcus</i> spp. Harboring Antibiotic Resistance Genes. <i>Journal of Food Protection</i> , 2014, 77, 993-998.	1.7	34
29	<i>Yersinia enterocolitica</i> : A Dangerous, But Often Ignored, Foodborne Pathogen. <i>Food Reviews International</i> , 2014, 30, 53-70.	8.4	27
30	Vidas UP – Enzyme-Linked Fluorescent Immunoassay Based on Recombinant Phage Protein and Fluorescence <i>In Situ</i> Hybridization as Alternative Methods for Detection of <i>Salmonella enterica</i> Serovars in Meat. <i>Foodborne Pathogens and Disease</i> , 2014, 11, 747-752.	1.8	8
31	FLUORESCENCE IN SITU HYBRIDIZATION AS ALTERNATIVE SCREENING METHOD FOR DETERMINING PRESENCE OF <i>SALMONELLA</i> SP. IN CHICKEN MEAT. <i>Zywnosc Nauka Technologia Jakosc/Food Science Technology Quality</i> , 2014, , .	0.1	0
32	<i>Salmonella</i> Detection in Poultry Meat – Validation of Vidas XPRESS Automatic Enzyme-Linked Fluorescent Immunoassay-Based Method. <i>Journal of Food Safety</i> , 2012, 32, 407-414.	2.3	11
33	Occurrence and antibiotic resistance of enterococci in ready-to-eat food of animal origin. <i>African Journal of Microbiology Research</i> , 2012, 6, 6773-6780.	0.4	17
34	<i>STAPHYLOCOCCUS AUREUS</i> FROM READY-TO-EAT FOOD AS A SOURCE OF MULTIPLE ANTIBIOTIC RESISTANCE GENES. <i>CBU International Conference Proceedings</i> , 0, 5, 1108-1112.	0.0	4