

# Thierry Touvier

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7037546/publications.pdf>

Version: 2024-02-01

17  
papers

909  
citations

623188

14  
h-index

887659

17  
g-index

17  
all docs

17  
docs citations

17  
times ranked

2448  
citing authors

#	ARTICLE	IF	CITATIONS
1	Regulation of Bile Acid Synthesis by the Nuclear Receptor Rev-erb $\beta$ . <i>Gastroenterology</i> , 2008, 135, 689-698.e5.	0.6	175
2	Requirement of Inducible Nitric Oxide Synthase for Skeletal Muscle Regeneration after Acute Damage. <i>Journal of Immunology</i> , 2013, 190, 1767-1777.	0.4	114
3	Muscle-specific Drp1 overexpression impairs skeletal muscle growth via translational attenuation. <i>Cell Death and Disease</i> , 2015, 6, e1663-e1663.	2.7	88
4	The amyloidogenic light chain is a stressor that sensitizes plasma cells to proteasome inhibitor toxicity. <i>Blood</i> , 2017, 129, 2132-2142.	0.6	70
5	Deficient nitric oxide signalling impairs skeletal muscle growth and performance: involvement of mitochondrial dysregulation. <i>Skeletal Muscle</i> , 2014, 4, 22.	1.9	58
6	Endoplasmic Reticulum Protein Quality Control Failure in Myelin Disorders. <i>Frontiers in Molecular Neuroscience</i> , 2016, 9, 162.	1.4	54
7	Partial dysferlin reconstitution by adult murine mesoangioblasts is sufficient for full functional recovery in a murine model of dysferlinopathy. <i>Cell Death and Disease</i> , 2010, 1, e61-e61.	2.7	53
8	Cripto regulates skeletal muscle regeneration and modulates satellite cell determination by antagonizing myostatin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, E3231-40.	3.3	48
9	LEPROT and LEPROTL1 cooperatively decrease hepatic growth hormone action in mice. <i>Journal of Clinical Investigation</i> , 2009, 119, 3830-3838.	3.9	47
10	Transplanted Mesoangioblasts Require Macrophage IL-10 for Survival in a Mouse Model of Muscle Injury. <i>Journal of Immunology</i> , 2012, 188, 6267-6277.	0.4	44
11	Autophagy controls neonatal myogenesis by regulating the GH-IGF1 system through a NFE2L2- and DDIT3-mediated mechanism. <i>Autophagy</i> , 2019, 15, 58-77.	4.3	41
12	Necdin is expressed in cachectic skeletal muscle to protect fibers from tumor-induced wasting. <i>Journal of Cell Science</i> , 2009, 122, 1119-1125.	1.2	35
13	Unexpected expression of orexin-B in basal conditions and increased levels in the adult rat hippocampus during pilocarpine-induced epileptogenesis. <i>Brain Research</i> , 2006, 1109, 164-175.	1.1	28
14	Necdin Enhances Myoblasts Survival by Facilitating the Degradation of the Mediator of Apoptosis CCAR1/CARP1. <i>PLoS ONE</i> , 2012, 7, e43335.	1.1	17
15	Treatment with IFB-088 Improves Neuropathy in CMT1A and CMT1B Mice. <i>Molecular Neurobiology</i> , 2022, 59, 4159-4178.	1.9	14
16	Necdin enhances muscle reconstitution of dystrophic muscle by vessel-associated progenitors, by promoting cell survival and myogenic differentiation. <i>Cell Death and Differentiation</i> , 2012, 19, 827-838.	5.0	13
17	Clearance of Cell Remnants and Regeneration of Injured Muscle Depend on Soluble Pattern Recognition Receptor PTX3. <i>Molecular Medicine</i> , 2016, 22, 809-820.	1.9	10