

# Tormod K Guren

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/703875/publications.pdf>

Version: 2024-02-01

53  
papers

1,281  
citations

471371

17  
h-index

395590

33  
g-index

53  
all docs

53  
docs citations

53  
times ranked

1756  
citing authors

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 1  | Ipilimumab in a real-world population: A prospective Phase IV trial with long-term follow-up. <i>International Journal of Cancer</i> , 2022, 150, 100-111.  | 2.3  | 11        |
| 2  | Survival Trends of Right- and Left-Sided Colon Cancer across Four Decades: A Norwegian Population-Based Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2022, 31, 342-351.  | 1.1  | 7         |
| 3  | Multiplex immunohistochemistry of metastatic colorectal cancer and ex vivo tumor avatars. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 2022, 1877, 188682.   | 3.3  | 1         |
| 4  | Pembrolizumab for previously treated advanced anal squamous cell carcinoma: results from the non-randomised, multicohort, multicentre, phase 2 KEYNOTE-158 study. <i>The Lancet Gastroenterology and Hepatology</i> , 2022, 7, 446-454.   | 3.7  | 36        |
| 5  | First-in-human phase 1 dose-escalation study of CANO4, a first-in-class interleukin-1 receptor accessory protein (IL1RAP) antibody in patients with solid tumours. <i>British Journal of Cancer</i> , 2022, 126, 1010-1017.   | 2.9  | 8         |
| 6  | Circulating tumour DNA and its clinical utility in predicting treatment response or survival in patients with metastatic colorectal cancer: a systematic review and meta-analysis. <i>British Journal of Cancer</i> , 2022, 127, 500-513.   | 2.9  | 15        |
| 7  | A national precision cancer medicine implementation initiative for Norway. <i>Nature Medicine</i> , 2022, 28, 885-887.  | 15.2 | 7         |
| 8  | Prototype precision oncology learning ecosystem: Norwegian precision cancer medicine implementation initiative.. <i>Journal of Clinical Oncology</i> , 2022, 40, e13634-e13634.   | 0.8  | 2         |
| 9  | Encorafenib Plus Cetuximab as a New Standard of Care for Previously Treated BRAF V600E-Mutant Metastatic Colorectal Cancer: Updated Survival Results and Subgroup Analyses from the BEACON Study. <i>Journal of Clinical Oncology</i> , 2021, 39, 273-284.  | 0.8  | 254       |
| 10 | Clinicopathological factors associated with tumour-specific mutation detection in plasma of patients with RAS-mutated or BRAF-mutated metastatic colorectal cancer. <i>International Journal of Cancer</i> , 2021, 149, 1385-1397.  | 2.3  | 10        |
| 11 | Combining a Universal Telomerase Based Cancer Vaccine With Ipilimumab in Patients With Metastatic Melanoma - Five-Year Follow Up of a Phase I/IIa Trial. <i>Frontiers in Immunology</i> , 2021, 12, 663865.   | 2.2  | 17        |
| 12 | Efficacy and safety of pembrolizumab in patients with advanced mesothelioma in the open-label, single-arm, phase 2 KEYNOTE-158 study. <i>Lancet Respiratory Medicine</i> , 2021, 9, 613-621.  | 5.2  | 44        |
| 13 | Long-Term Outcomes of a Phase I Study With UV1, a Second Generation Telomerase Based Vaccine, in Patients With Advanced Non-Small Cell Lung Cancer. <i>Frontiers in Immunology</i> , 2020, 11, 572172.  | 2.2  | 21        |
| 14 | Patient-Derived Organoids from Multiple Colorectal Cancer Liver Metastases Reveal Moderate Intra-patient Pharmacotranscriptomic Heterogeneity. <i>Clinical Cancer Research</i> , 2020, 26, 4107-4119.   | 3.2  | 68        |
| 15 | Encorafenib plus cetuximab with or without binimetinib for BRAF V600E metastatic colorectal cancer: Updated survival results from a randomized, three-arm, phase III study versus choice of either irinotecan or FOLFIRI plus cetuximab (BEACON CRC).. <i>Journal of Clinical Oncology</i> , 2020, 38, 4001-4001.           | 0.8  | 35        |
| 16 | Encorafenib plus cetuximab with or without binimetinib for BRAF V600E-mutant metastatic colorectal cancer: Quality-of-life results from a randomized, three-arm, phase III study versus the choice of either irinotecan or FOLFIRI plus cetuximab (BEACON CRC).. <i>Journal of Clinical Oncology</i> , 2020, 38, 4039-4039. | 0.8  | 5         |
| 17 | Pembrolizumab for advanced anal squamous cell carcinoma (ASCC): Results from the multicohort, phase II KEYNOTE-158 study.. <i>Journal of Clinical Oncology</i> , 2020, 38, 1-1.   | 0.8  | 19        |
| 18 | Encorafenib plus cetuximab with or without binimetinib for BRAF V600E-mutant metastatic colorectal cancer: Quality-of-life results from a randomized, three-arm, phase III study versus the choice of either irinotecan or FOLFIRI plus cetuximab (BEACON CRC).. <i>Journal of Clinical Oncology</i> , 2020, 38, 8-8.       | 0.8  | 25        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | A phase I/IIa clinical trial investigating the therapeutic cancer vaccine UV1 in combination with ipilimumab in patients with malignant melanoma: Four-year survival update.. Journal of Clinical Oncology, 2020, 38, 62-62.  | 0.8 | 2         |
| 20 | Safety and efficacy of nivolumab in challenging subgroups with advanced melanoma who progressed on or after ipilimumab treatment: A single-arm, open-label, phase II study (CheckMate 172). European Journal of Cancer, 2019, 121, 144-153.                                   | 1.3 | 27        |
| 21 | Safety and efficacy of nivolumab in patients with rare melanoma subtypes who progressed on or after ipilimumab treatment: a single-arm, open-label, phase II study (CheckMate 172). European Journal of Cancer, 2019, 119, 168-178.   | 1.3 | 61        |
| 22 | Combination therapies with HSP90 inhibitors against colorectal cancer. Biochimica Et Biophysica Acta: Reviews on Cancer, 2019, 1871, 240-247.   | 3.3 | 81        |
| 23 | Transplant oncology: assessment of response and tolerance to systemic chemotherapy for metastatic colorectal cancer after liver transplantation â€” a retrospective study. Transplant International, 2019, 32, 1144-1150.   | 0.8 | 23        |
| 24 | Results from a first-in-man, open label, safety and tolerability trial of CAN04 (nidanimab), a fully humanized monoclonal antibody against the novel antitumor target, IL1RAP, in patients with solid tumor malignancies.. Journal of Clinical Oncology, 2019, 37, 2504-2504. | 0.8 | 5         |
| 25 | Treatment-related survival associations of claudin-2 expression in fibroblasts of colorectal cancer. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2018, 472, 395-405.  | 1.4 | 10        |
| 26 | Graft rejection after immune checkpoint inhibitor therapy in solid organ transplant recipients. Acta Oncologica, 2018, 57, 1414-1418.   | 0.8 | 19        |
| 27 | Prognostic role of carcinoembryonic antigen and carbohydrate antigen 19-9 in metastatic colorectal cancer: a BRAF-mutant subset with high CA 19-9 level and poor outcome. British Journal of Cancer, 2018, 118, 1609-1616.  | 2.9 | 47        |
| 28 | ISO-CC-005: A phase I/II study of Modufolin (MTHF) in combination with 5-FU, irinotecan, and oxaliplatin ± bevacizumab in patients with metastasizing colorectal cancer.. Journal of Clinical Oncology, 2018, 36, 838-838.  | 0.8 | 0         |
| 29 | Cetuximab in treatment of metastatic colorectal cancer: final survival analyses and extended RAS data from the NORDIC-VII study. British Journal of Cancer, 2017, 116, 1271-1278.   | 2.9 | 55        |
| 30 | Health-related quality of life in patients with metastatic colorectal cancer, association with systemic inflammatory response and RAS and BRAF mutation status. European Journal of Cancer, 2017, 81, 26-35.  | 1.3 | 13        |
| 31 | Dalotuzumab in chemorefractory KRAS exon 2 mutant colorectal cancer: Results from a randomised phase II/III trial. International Journal of Cancer, 2017, 140, 431-439.   | 2.3 | 4         |
| 32 | Efficacy and safety of nivolumab (NIVO) in patients with advanced melanoma (MEL) and poor prognostic factors who progressed on or after ipilimumab (IPI): Results from a phase II study (CheckMate 172).. Journal of Clinical Oncology, 2017, 35, 9524-9524.                  | 0.8 | 17        |
| 33 | Phase 2 results: Encorafenib (ENCO) and cetuximab (CETUX) with or without alpelisib (ALP) in patients with advanced BRAF-mutant colorectal cancer (BRAFM CRC).. Journal of Clinical Oncology, 2016, 34, 3544-3544.  | 0.8 | 79        |
| 34 | Nivolumab (NIVO) safety in patients with advanced melanoma (MEL) who have progressed on or after ipilimumab (IPI): A single-arm, open-label, multicenter, phase II study (CheckMate 172).. Journal of Clinical Oncology, 2016, 34, 9526-9526.                                 | 0.8 | 2         |
| 35 | TIMP-1 is under regulation of the EGF signaling axis and promotes an aggressive phenotype in KRAS-mutated colorectal cancer cells: A potential novel approach to the treatment of metastatic colorectal cancer. Oncotarget, 2016, 7, 59441-59457.                             | 0.8 | 7         |
| 36 | Interleukin-6 and C-reactive protein as prognostic biomarkers in metastatic colorectal cancer. Oncotarget, 2016, 7, 75013-75022.  | 0.8 | 61        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | Intact and cleaved plasma soluble urokinase receptor in patients with metastatic colorectal cancer treated with oxaliplatin with or without cetuximab. <i>International Journal of Cancer</i> , 2015, 137, 2470-2477.   | 2.3 | 8         |
| 38 | C-reactive protein and interleukin-6 as markers of systemic inflammatory response and as prognostic factors for metastatic colorectal cancer. Data from the randomized phase III NORDIC-VII study.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3548-3548.  | 0.8 | 0         |
| 39 | Plasma YKL-40 in Patients with Metastatic Colorectal Cancer Treated with First Line Oxaliplatin-Based Regimen with or without Cetuximab: RESULTS from the NORDIC VII Study. <i>PLoS ONE</i> , 2014, 9, e87746.  | 1.1 | 18        |
| 40 | Outcome after liver transplantation compared with chemotherapy in colorectal cancer patients with nonresectable liver-only disease.. <i>Journal of Clinical Oncology</i> , 2014, 32, 531-531.   | 0.8 | 0         |
| 41 | Digitalized multiparametric analyses of tumor stroma for identification of low perivascular PDGFBR expression and low vessel density as independent prognosis markers for stage IV CRC.. <i>Journal of Clinical Oncology</i> , 2014, 32, e14525-e14525.   | 0.8 | 0         |
| 42 | Benefit of EGFR-inhibition therapy for metastatic colorectal cancer patients with KRAS-mutated tumors and high plasma TIMP-1 level: Results from the NORDIC VII study.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3590-3590.  | 0.8 | 0         |
| 43 | Plasma levels of TIMP-1 in chemo-naïve patients with metastatic colorectal cancer treated with first-line FLOX with or without cetuximab: Results from the Nordic VII Study.. <i>Journal of Clinical Oncology</i> , 2013, 31, 392-392.  | 0.8 | 0         |
| 44 | Plasma TIMP-1 in patients with metastatic colorectal cancer treated with first-line oxaliplatin-based therapy with or without cetuximab: Results from the Nordic VII study.. <i>Journal of Clinical Oncology</i> , 2013, 31, e14710-e14710.   | 0.8 | 0         |
| 45 | On-treatment progression-free survival analysis of ziv-aflibercept/FOLFIRI treatment within 28 days of end of treatment in metastatic colorectal cancer: Updated efficacy results from the VELOUR study.. <i>Journal of Clinical Oncology</i> , 2013, 31, 3573-3573.  | 0.8 | 0         |
| 46 | Prognostic significance of tumor stromal and epithelial claudin 2 in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2013, 31, 3597-3597.  | 0.8 | 0         |
| 47 | Tumor perivascular PDGFBR as an independent prognostic factor in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2013, 31, 3571-3571.  | 0.8 | 0         |
| 48 | Molecular analysis of the randomized phase II/III study of the anti-IGF-1R antibody dalotuzumab (MK-0646) in combination with cetuximab (Cx) and irinotecan (Ir) in the treatment of chemorefractory KRAS wild-type metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2012, 30, 3531-3531.                      | 0.8 | 8         |
| 49 | Plasma concentrations of YKL-40 in chemo-naïve patients with metastatic colorectal cancer treated with FLOX with or without cetuximab: Results from the NORDIC VII study.. <i>Journal of Clinical Oncology</i> , 2012, 30, 3548-3548.   | 0.8 | 0         |
| 50 | EGF receptor-mediated, c-Src-dependent, activation of Stat5b is downregulated in mitogenically responsive hepatocytes. <i>Journal of Cellular Physiology</i> , 2003, 196, 113-123.  | 2.0 | 17        |
| 51 | Response to transforming growth factor $\hat{\pm}$ (TGF $\hat{\pm}$ ) and epidermal growth factor (EGF) in hepatocytes: Lower EGF receptor affinity of TGF $\hat{\pm}$ is associated with more sustained activation of p42/p44 mitogen-activated protein kinase and greater efficacy in stimulation of DNA synthesis. , 1998, 175, 10-18. |     | 55        |
| 52 | Dual effects of glucagon and cyclic amp on dna synthesis in cultured rat hepatocytes: Stimulatory regulation in early G1 and inhibition shortly before the s phase entry. <i>Journal of Cellular Physiology</i> , 1990, 144, 523-530.   | 2.0 | 47        |
| 53 | Survival-associated heterogeneity of marker-defined perivascular cells in colorectal cancer. <i>Oncotarget</i> , 0, 7, 41948-41958.   | 0.8 | 30        |