

Marek Chmelik

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7050373/publications.pdf>

Version: 2024-02-01

51
papers

2,211
citations

218381

26
h-index

223531

46
g-index

52
all docs

52
docs citations

52
times ranked

2760
citing authors

#	ARTICLE	IF	CITATIONS
1	A systematic review on the use of quantitative imaging to detect cancer therapy adverse effects in normal-appearing brain tissue. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2022, 35, 163-186.	1.1	7
2	In Vivo ¹ H MRSpectroscopy of Biliary Components of Human Gallbladder at 7T. <i>Journal of Magnetic Resonance Imaging</i> , 2021, 53, 98-107.	1.9	3
3	Concentration of Gallbladder Phosphatidylcholine in Cholangiopathies: A Phosphorus- ³¹ Magnetic Resonance Spectroscopy Pilot Study. <i>Journal of Magnetic Resonance Imaging</i> , 2021, , .	1.9	2
4	Absolute Quantification of Phosphorus-Containing Metabolites in the Liver Using ³¹ P MRSI and Hepatic Lipid Volume Correction at 7T Suggests No Dependence on Body Mass Index or Age. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 49, 597-607.	1.9	16
5	Effect of Rehabilitation on Fatigue Level in Patients with Multiple Sclerosis. <i>Medical Science Monitor</i> , 2018, 24, 5761-5770.	0.5	8
6	In-vivo ³¹ P-MRS of skeletal muscle and liver: A way for non-invasive assessment of their metabolism. <i>Analytical Biochemistry</i> , 2017, 529, 193-215.	1.1	78
7	Gliptin therapy reduces hepatic and myocardial fat in type 2 diabetic patients. <i>European Journal of Clinical Investigation</i> , 2017, 47, 829-838.	1.7	11
8	(² H- and ¹ H)-CAIPIRINHA accelerated MR spectroscopic imaging of the brain at 7T. <i>Magnetic Resonance in Medicine</i> , 2017, 78, 429-440.	1.9	46
9	Skeletal muscle alkaline Pi pool is decreased in overweight-to-obese sedentary subjects and relates to mitochondrial capacity and phosphodiester content. <i>Scientific Reports</i> , 2016, 6, 20087.	1.6	26
10	Feasibility and repeatability of localized ³¹ P MRS four-angle saturation transfer (FAST) of the human gastrocnemius muscle using a surface coil at 7T. <i>NMR in Biomedicine</i> , 2016, 29, 57-65.	1.6	14
11	Improved spectral resolution and high reliability of in vivo ¹ H MRS at 7 T allow the characterization of the effect of acute exercise on carnosine in skeletal muscle. <i>NMR in Biomedicine</i> , 2016, 29, 24-32.	1.6	22
12	Dynamic ³¹ P MRSI using spiral spectroscopic imaging can map mitochondrial capacity in muscles of the human calf during plantar flexion exercise at 7T. <i>NMR in Biomedicine</i> , 2016, 29, 1825-1834.	1.6	38
13	Diffusion-weighted imaging of breast tumours at 3T and 7T: a comparison. <i>European Radiology</i> , 2016, 26, 1466-1473.	2.3	18
14	Lipid suppression via double inversion recovery with symmetric frequency sweep for robust 2D CRAPPA accelerated MRSI of the brain at 7T. <i>NMR in Biomedicine</i> , 2015, 28, 1413-1425.	1.6	48
15	Dynamic ³¹ P MR spectroscopy of plantar flexion: Influence of ergometer design, magnetic field strength (3 and 7 T), and RF coil design. <i>Medical Physics</i> , 2015, 42, 1678-1689.	1.6	26
16	Ultrashort-TE stimulated echo acquisition mode (STEAM) improves the quantification of lipids and fatty acid chain unsaturation in the human liver at 7T. <i>NMR in Biomedicine</i> , 2015, 28, 1283-1293.	1.6	27
17	Phosphatidylcholine contributes to in vivo ³¹ P MRS signal from the human liver. <i>European Radiology</i> , 2015, 25, 2059-2066.	2.3	19
18	Dixon imaging-based partial volume correction improves quantification of choline detected by breast 3D-MRSI. <i>European Radiology</i> , 2015, 25, 830-836.	2.3	2

#	ARTICLE	IF	CITATIONS
19	Mapping of brain macromolecules and their use for spectral processing of ^1H -MRSI data with an ultra-short acquisition delay at 7 T. <i>NeuroImage</i> , 2015, 121, 126-135.	2.1	62
20	Depth-resolved surface coil MRS (DRESS)-localized dynamic ^{31}P -MRS of the exercising human gastrocnemius muscle at 7 T. <i>NMR in Biomedicine</i> , 2014, 27, 1346-1352.	1.6	35
21	<i>In vivo</i> ^{31}P magnetic resonance spectroscopy of the human liver at 7 T: an initial experience. <i>NMR in Biomedicine</i> , 2014, 27, 478-485.	1.6	38
22	Dynamic Contrast-Enhanced Magnetic Resonance Imaging of Breast Tumors at 3 and 7 T. <i>Investigative Radiology</i> , 2014, 49, 354-362.	3.5	27
23	Lower Fasting Muscle Mitochondrial Activity Relates to Hepatic Steatosis in Humans. <i>Diabetes Care</i> , 2014, 37, 468-474.	4.3	26
24	Flip-angle mapping of ^{31}P coils by steady-state MR spectroscopic imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 40, 391-397.	1.9	14
25	Two-dimensional spectroscopic imaging with combined free induction decay and long-TE acquisition (FID echo spectroscopic imaging, FIDESI) for the detection of intramyocellular lipids in calf muscle at 7 T. <i>NMR in Biomedicine</i> , 2014, 27, 980-987.	1.6	10
26	Application of localized ^{31}P MRS saturation transfer at 7 T for measurement of ATP metabolism in the liver: reproducibility and initial clinical application in patients with non-alcoholic fatty liver disease. <i>European Radiology</i> , 2014, 24, 1602-1609.	2.3	27
27	One-dimensional image-selected <i>in vivo</i> spectroscopy localized phosphorus saturation transfer at 7T. <i>Magnetic Resonance in Medicine</i> , 2014, 72, 1509-1515.	1.9	17
28	<i>In vivo</i> relaxation behavior of liver compounds at 7 tesla, measured by single-voxel proton MR spectroscopy. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 40, 1365-1374.	1.9	19
29	Fully adiabatic ^{31}P 2D-CSI with reduced chemical shift displacement error at 7 T "GOIA-1D-ISIS/2D-CSI". <i>Magnetic Resonance in Medicine</i> , 2013, 69, 1233-1244.	1.9	28
30	Time-resolved phosphorous magnetization transfer of the human calf muscle at 3T and 7T: A feasibility study. <i>European Journal of Radiology</i> , 2013, 82, 745-751.	1.2	28
31	Interrelation of ^{31}P -MRS metabolism measurements in resting and exercised quadriceps muscle of overweight-to-obese sedentary individuals. <i>NMR in Biomedicine</i> , 2013, 26, 1714-1722.	1.6	29
32	Coil combination of multichannel MRSI data at 7 T: MUSICAL. <i>NMR in Biomedicine</i> , 2013, 26, 1796-1805.	1.6	45
33	Readout-segmented Echo-planar Imaging Improves the Diagnostic Performance of Diffusion-weighted MR Breast Examinations at 3.0 T. <i>Radiology</i> , 2012, 263, 64-76.	3.6	180
34	A Single Nucleotide Polymorphism Associates With the Response of Muscle ATP Synthesis to Long-Term Exercise Training in Relatives of Type 2 Diabetic Humans. <i>Diabetes Care</i> , 2012, 35, 350-357.	4.3	25
35	High-resolution mapping of human brain metabolites by free induction decay ^1H MRSI at 7 T. <i>NMR in Biomedicine</i> , 2012, 25, 873-882.	1.6	91
36	Fatty Liver Index Predicts Further Metabolic Deteriorations in Women with Previous Gestational Diabetes. <i>PLoS ONE</i> , 2012, 7, e32710.	1.1	49

#	ARTICLE	IF	CITATIONS
37	Skeletal Muscle Phosphodiester Content Relates to Body Mass and Glycemic Control. PLoS ONE, 2011, 6, e21846.	1.1	22
38	Impaired insulin stimulation of muscular ATP production in patients with type 1 diabetes. Journal of Internal Medicine, 2011, 269, 189-199.	2.7	42
39	Vascular function in obese children with non-alcoholic fatty liver disease. Pediatric Obesity, 2011, 6, 120-127.	3.2	23
40	In vivo ³¹ P spectroscopy by fully adiabatic extended image selected in vivo spectroscopy: A comparison between 3 T and 7 T. Magnetic Resonance in Medicine, 2011, 66, 923-930.	1.9	40
41	Three-dimensional Proton MR Spectroscopic Imaging at 3 T for the Differentiation of Benign and Malignant Breast Lesions. Radiology, 2011, 261, 752-761.	3.6	61
42	Body and Liver Fat Mass Rather Than Muscle Mitochondrial Function Determine Glucose Metabolism in Women With a History of Gestational Diabetes Mellitus. Diabetes Care, 2011, 34, 430-436.	4.3	42
43	Postprandial and Fasting Hepatic Glucose Fluxes in Long-Standing Type 1 Diabetes. Diabetes, 2011, 60, 1752-1758.	0.3	33
44	Liver ATP Synthesis Is Lower and Relates to Insulin Sensitivity in Patients With Type 2 Diabetes. Diabetes Care, 2011, 34, 448-453.	4.3	177
45	Short-Term Exercise Training Does Not Stimulate Skeletal Muscle ATP Synthesis in Relatives of Humans With Type 2 Diabetes. Diabetes, 2009, 58, 1333-1341.	0.3	62
46	Abnormal hepatic energy homeostasis in type 2 diabetes. Hepatology, 2009, 50, 1079-1086.	3.6	166
47	Quantitative ATP synthesis in human liver measured by localized ³¹ P spectroscopy using the magnetization transfer experiment. NMR in Biomedicine, 2008, 21, 437-443.	1.6	61
48	Metabolic changes in the normal ageing brain: Consistent findings from short and long echo time proton spectroscopy. European Journal of Radiology, 2008, 68, 320-327.	1.2	76
49	Reduced Basal ATP Synthetic Flux of Skeletal Muscle in Patients with Previous Acromegaly. PLoS ONE, 2008, 3, e3958.	1.1	29
50	Muscle Mitochondrial ATP Synthesis and Glucose Transport/Phosphorylation in Type 2 Diabetes. PLoS Medicine, 2007, 4, e154.	3.9	216
51	Ankylosing spondylitis on unidentified individual from early modern times found in reformed church (Silická Brezová, Slovakia): a case-based review. Rheumatology International, 0, , .	1.5	0