

Svein Friis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7055561/publications.pdf>

Version: 2024-02-01

60
papers

2,632
citations

304743

22
h-index

189892

50
g-index

64
all docs

64
docs citations

64
times ranked

2475
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Predictive validity of attention-deficit/hyperactivity disorder from ages 3 to 5 Years. <i>European Child and Adolescent Psychiatry</i> , 2022, 31, 1-10. | 4.7 | 7 |
| 2 | Cognitive, emotional, and social functioning of preschoolers with attention deficit hyperactivity problems. <i>BMC Psychiatry</i> , 2022, 22, 78. | 2.6 | 8 |
| 3 | The effect of special educational assistance in early childhood education and care on psycho-social difficulties in elementary school children. <i>Child and Adolescent Psychiatry and Mental Health</i> , 2022, 16, 14. | 2.5 | 0 |
| 4 | Continuity of care and therapeutic relationships as critical elements in acute psychiatric care. <i>World Psychiatry</i> , 2022, 21, 241-242. | 10.4 | 6 |
| 5 | Youth with hearing loss: Emotional and behavioral problems and quality of life. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2021, 145, 110718. | 1.0 | 11 |
| 6 | The factor structure of social cognition in schizophrenia: Weak evidence for separable domains. <i>Schizophrenia Research: Cognition</i> , 2021, 26, 100208. | 1.3 | 1 |
| 7 | Lipid alterations in adolescents with early-onset psychosis may be independent of antipsychotic medication. <i>Schizophrenia Research</i> , 2020, 216, 295-301. | 2.0 | 10 |
| 8 | Symptom remission at 12-weeks is a strong predictor for long term outcome. <i>Psychological Medicine</i> , 2020, 50, 2464-2464. | 4.5 | 0 |
| 9 | Increased interleukin 18 activity in adolescents with early-onset psychosis is associated with cortisol and depressive symptoms. <i>Psychoneuroendocrinology</i> , 2020, 112, 104513. | 2.7 | 21 |
| 10 | Reduced levels of circulating adhesion molecules in adolescents with early-onset psychosis. <i>NPJ Schizophrenia</i> , 2020, 6, 20. | 3.6 | 10 |
| 11 | Natural History, Not Lead Time. <i>American Journal of Psychiatry</i> , 2020, 177, 1185-1185. | 7.2 | 6 |
| 12 | The Associations Between Pre- and Postnatal Maternal Symptoms of Distress and Preschooler's Symptoms of ADHD, Oppositional Defiant Disorder, Conduct Disorder, and Anxiety. <i>Journal of Attention Disorders</i> , 2020, 24, 1057-1069. | 2.6 | 23 |
| 13 | Parsing the impact of early detection on duration of untreated psychosis (DUP): Applying quantile regression to data from the Scandinavian TIPS study. <i>Schizophrenia Research</i> , 2019, 210, 128-134. | 2.0 | 15 |
| 14 | Early Substance Use Cessation Improves Cognition's 10 Years Outcome in First-Episode Psychosis Patients. <i>Frontiers in Psychiatry</i> , 2019, 10, 495. | 2.6 | 6 |
| 15 | The predictive validity of the Strengths and Difficulties Questionnaire for child attention-deficit/hyperactivity disorder. <i>European Child and Adolescent Psychiatry</i> , 2019, 28, 625-633. | 4.7 | 13 |
| 16 | Attention-Deficit/Hyperactivity Disorder in Preschoolers: The Accuracy of a Short Screener. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2018, 57, 428-435. | 0.5 | 32 |
| 17 | Relation between premorbid adjustment, duration of untreated psychosis and close interpersonal trauma in first-episode psychosis. <i>Microbial Biotechnology</i> , 2018, 12, 316-323. | 1.7 | 26 |
| 18 | Is going into stable symptomatic remission associated with a more positive development of life satisfaction? A 10-year follow-up study of first episode psychosis. <i>Schizophrenia Research</i> , 2018, 193, 364-369. | 2.0 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 19 | A 10-year follow-up study of violent victimization in first episode psychosis: Risk and protective factors. <i>Psychiatry Research</i> , 2018, 259, 545-549. | 3.3 | 9 |
| 20 | F136. PARSING DUP TO REFINE EARLY DETECTION: QUANTILE REGRESSION OF RESULTS FROM THE SCANDINAVIAN TIPS STUDY. <i>Schizophrenia Bulletin</i> , 2018, 44, S272-S273. | 4.3 | 0 |
| 21 | Dr. Åvergaard et al. Reply. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2018, 57, 701-702. | 0.5 | 0 |
| 22 | Reported use of evidence in clinical practice: a survey of rehabilitation practices in Norway. <i>BMC Health Services Research</i> , 2018, 18, 379. | 2.2 | 8 |
| 23 | Depression and functioning are important to subjective quality of life after a first episode psychosis. <i>Comprehensive Psychiatry</i> , 2018, 86, 107-114. | 3.1 | 6 |
| 24 | Co-Occurrence of ODD and CD in Preschool Children With Symptoms of ADHD. <i>Journal of Attention Disorders</i> , 2017, 21, 741-752. | 2.6 | 30 |
| 25 | Causes and predictors of premature death in first-episode schizophrenia spectrum disorders. <i>World Psychiatry</i> , 2017, 16, 217-218. | 10.4 | 21 |
| 26 | The Effect of Substance Use on 10-Year Outcome in First-Episode Psychosis. <i>Schizophrenia Bulletin</i> , 2017, 43, 843-851. | 4.3 | 55 |
| 27 | Vocational rehabilitation for adults with psychotic disorders in a Scandinavian welfare society. <i>BMC Psychiatry</i> , 2017, 17, 24. | 2.6 | 23 |
| 28 | Characteristics of adolescents frequently restrained in acute psychiatric units in Norway: a nationwide study. <i>Child and Adolescent Psychiatry and Mental Health</i> , 2017, 11, 3. | 2.5 | 14 |
| 29 | Neurocognition and Duration of Psychosis: A 10-year Follow-up of First-Episode Patients. <i>Schizophrenia Bulletin</i> , 2016, 42, sbv083. | 4.3 | 77 |
| 30 | Participants' perceived benefits of family intervention following a first episode of psychosis: a qualitative study. <i>Microbial Biotechnology</i> , 2016, 10, 152-159. | 1.7 | 24 |
| 31 | Analyzing the Duration of Untreated Psychosis. <i>JAMA Psychiatry</i> , 2016, 73, 1094. | 11.0 | 10 |
| 32 | Depressive symptoms in first-episode psychosis: a 10-year follow-up study. <i>Microbial Biotechnology</i> , 2016, 10, 227-233. | 1.7 | 35 |
| 33 | The Global Body Examination (GBE): A useful instrument for examination of patients with long-lasting musculoskeletal and/or psychological disorders. <i>European Journal of Physiotherapy</i> , 2016, 18, 137-143. | 1.3 | 2 |
| 34 | A nationwide study of why and how acute adolescent psychiatric units use restraint. <i>Psychiatry Research</i> , 2016, 237, 60-66. | 3.3 | 18 |
| 35 | Subjective quality of life in first-episode psychosis. A ten year follow-up study. <i>Schizophrenia Research</i> , 2016, 172, 23-28. | 2.0 | 56 |
| 36 | Early Predictors of Ten-Year Course in First-Episode Psychosis. <i>Psychiatric Services</i> , 2016, 67, 438-443. | 2.0 | 47 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | Challenges for group leaders working with families dealing with early psychosis: a qualitative study. <i>BMC Psychiatry</i> , 2015, 15, 141. | 2.6 | 9 |
| 38 | Patients'™ and Family Members'™ Experiences of a Psychoeducational Family Intervention after a First Episode Psychosis: A Qualitative Study. <i>Issues in Mental Health Nursing</i> , 2014, 35, 58-68. | 1.2 | 23 |
| 39 | Treatment and violent behavior in persons with first episode psychosis during a 10-year prospective follow-up study. <i>Schizophrenia Research</i> , 2014, 156, 272-276. | 2.0 | 42 |
| 40 | Characteristics of patients frequently subjected to pharmacological and mechanical restraint"™A register study in three Norwegian acute psychiatric wards. <i>Psychiatry Research</i> , 2014, 215, 127-133. | 3.3 | 41 |
| 41 | Early detection, early symptom progression and symptomatic remission after ten years in a first episode of psychosis study. <i>Schizophrenia Research</i> , 2013, 143, 337-343. | 2.0 | 41 |
| 42 | Impairment in Young Preschool Children with Symptoms of Attention-Deficit/Hyperactivity Disorder and Co-occurring Oppositional Defiant Disorder and Conduct Disorder. <i>Scandinavian Journal of Child and Adolescent Psychiatry and Psychology</i> , 2013, 2, 95-105. | 0.6 | 5 |
| 43 | The TIPS study and bias: Replies to Dr Amos. <i>Australian and New Zealand Journal of Psychiatry</i> , 2012, 46, 905-907. | 2.3 | 2 |
| 44 | Long-Term Follow-Up of the TIPS Early Detection in Psychosis Study: Effects on 10-Year Outcome. <i>American Journal of Psychiatry</i> , 2012, 169, 374-380. | 7.2 | 262 |
| 45 | The Global Body Examination (GBE). A useful instrument for evaluation of respiration. <i>Advances in Physiotherapy</i> , 2012, 14, 146-154. | 0.2 | 2 |
| 46 | Neurocognition and occupational functioning in patients with first-episode psychosis: A 2-year follow-up study. <i>Psychiatry Research</i> , 2011, 188, 334-342. | 3.3 | 41 |
| 47 | Development of the Posture domain in the Global Body Examination (GBE). <i>Advances in Physiotherapy</i> , 2010, 12, 157-165. | 0.2 | 4 |
| 48 | Reflections on Psychotherapeutic approaches to schizophrenic psychoses: past, present and future, by Alanen, de Ch'vez, Silver & Martindale. <i>Scandinavian Psychoanalytic Review</i> , 2010, 33, 47-51. | 0.4 | 3 |
| 49 | Personality disorders in first-episode psychosis. <i>Personality and Mental Health</i> , 2008, 2, 230-239. | 1.2 | 18 |
| 50 | Prevention of Negative Symptom Psychopathologies in First-Episode Schizophrenia. <i>Archives of General Psychiatry</i> , 2008, 65, 634. | 12.3 | 163 |
| 51 | The Key to Reducing Duration of Untreated First Psychosis: Information Campaigns. <i>Schizophrenia Bulletin</i> , 2007, 34, 466-472. | 4.3 | 169 |
| 52 | Effects on referral patterns of reducing intensive informational campaigns about first-episode psychosis. <i>Microbial Biotechnology</i> , 2007, 1, 340-345. | 1.7 | 11 |
| 53 | Substance abuse in first-episode non-affective psychosis. <i>Schizophrenia Research</i> , 2006, 88, 55-62. | 2.0 | 84 |
| 54 | Early Detection of the First Episode of Schizophrenia and Suicidal Behavior. <i>American Journal of Psychiatry</i> , 2006, 163, 800-804. | 7.2 | 186 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 55 | Premorbid adjustment in first-episode non-affective psychosis: Distinct patterns of pre-onset course. <i>British Journal of Psychiatry</i> , 2004, 185, 108-115. | 2.8 | 140 |
| 56 | Reducing the Duration of Untreated First-Episode Psychosis. <i>Archives of General Psychiatry</i> , 2004, 61, 143. | 12.3 | 384 |
| 57 | Early detection strategies for untreated first-episode psychosis. <i>Schizophrenia Research</i> , 2001, 51, 39-46. | 2.0 | 208 |
| 58 | Shortened Duration of Untreated First Episode of Psychosis: Changes in Patient Characteristics at Treatment. <i>American Journal of Psychiatry</i> , 2001, 158, 1917-1919. | 7.2 | 151 |
| 59 | The Ullevål acute-ward follow-up study: Reflections on the treatment of psychotic patients who committed suicide. <i>Nordic Journal of Psychiatry</i> , 1995, 49, 5-10. | 1.3 | 3 |
| 60 | Problems in Treating Late-Onset Schizophrenia: An 8-year follow-up study. <i>Nordic Journal of Psychiatry</i> , 1994, 48, 235-240. | 1.3 | 2 |