

# Annie Sittler

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/7066240/annie-sittler-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

10  
papers

684  
citations

9  
h-index

10  
g-index

10  
ext. papers

739  
ext. citations

9  
avg, IF

2.8  
L-index

#	Paper	IF	Citations
10	SUMOylation by SUMO2 is implicated in the degradation of misfolded ataxin-7 via RNF4 in SCA7 models. <i>DMM Disease Models and Mechanisms</i> , <b>2019</b> , 12,	4.1	10
9	Deregulation of autophagy in postmortem brains of Machado-Joseph disease patients. <i>Neuropathology</i> , <b>2018</b> , 38, 113-124	2	32
8	Control of Huntington's Disease-Associated Phenotypes by the Striatum-Enriched Transcription Factor Foxp2. <i>Cell Reports</i> , <b>2017</b> , 21, 2688-2695	10.6	17
7	Lentiviral vector-mediated overexpression of mutant ataxin-7 recapitulates SCA7 pathology and promotes accumulation of the FUS/TLS and MBNL1 RNA-binding proteins. <i>Molecular Neurodegeneration</i> , <b>2016</b> , 11, 58	19	7
6	The autophagy/lysosome pathway is impaired in SCA7 patients and SCA7 knock-in mice. <i>Acta Neuropathologica</i> , <b>2014</b> , 128, 705-22	14.3	42
5	Interferon $\gamma$ induces clearance of mutant ataxin 7 and improves locomotion in SCA7 knock-in mice. <i>Brain</i> , <b>2013</b> , 136, 1732-45	11.2	50
4	SUMOylation attenuates the aggregation propensity and cellular toxicity of the polyglutamine expanded ataxin-7. <i>Human Molecular Genetics</i> , <b>2010</b> , 19, 181-95	5.6	69
3	PML clastosomes prevent nuclear accumulation of mutant ataxin-7 and other polyglutamine proteins. <i>Journal of Cell Biology</i> , <b>2006</b> , 174, 65-76	7.3	75
2	Membrane filter assay for detection of amyloid-like polyglutamine-containing protein aggregates. <i>Methods in Enzymology</i> , <b>1999</b> , 309, 375-86	1.7	193
1	SH3GL3 associates with the Huntingtin exon 1 protein and promotes the formation of polyglu-containing protein aggregates. <i>Molecular Cell</i> , <b>1998</b> , 2, 427-36	17.6	189