
Susanna Lallukka

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7073583/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Saturated Fat Is More Metabolically Harmful for the Human Liver Than Unsaturated Fat or Simple Sugars. Diabetes Care, 2018, 41, 1732-1739.	8.6	266
2	A population-based study on the prevalence of NASH using scores validated against liver histology. Journal of Hepatology, 2014, 60, 839-846.	3.7	107
3	Human PNPLA3-I148M variant increases hepatic retention of polyunsaturated fatty acids. JCI Insight, 2019, 4, .	5.0	93
4	Predictors of Liver Fat and Stiffness in Non-Alcoholic Fatty Liver Disease (NAFLD) – an 11-Year Prospective Study. Scientific Reports, 2017, 7, 14561.	3.3	18
5	Obesity/insulin resistance rather than liver fat increases coagulation factor activities and expression in humans. Thrombosis and Haemostasis, 2017, 117, 286-294.	3.4	18
6	The PNPLA3â€I148M variant increases polyunsaturated triglycerides in human adipose tissue. Liver International, 2020, 40, 2128-2138.	3.9	17
7	Assessment of Lifestyle Factors Helps to Identify Liver Fibrosis Due to Non-Alcoholic Fatty Liver Disease in Obesity. Nutrients, 2021, 13, 169.	4.1	9
8	Effects of Weighted Hula-Hooping Compared to Walking on Abdominal Fat, Trunk Muscularity, and Metabolic Parameters in Overweight Subjects: A Randomized Controlled Study. Obesity Facts, 2019, 12, 385-396.	3.4	7
9	Natural Course of Nonalcoholic Fatty Liver Disease and Type 2 Diabetes in Patients With Human Immunodeficiency Virus With and Without Combination Antiretroviral Therapy–associated Lipodystrophy: A 16-Year Follow-up Study. Clinical Infectious Diseases, 2020, 70, 1708-1716.	5.8	6
10	Reply to Krahn and Sebastiani. Clinical Infectious Diseases, 2020, 71, 245-245.	5.8	0

10