

Youichirou Ootsuka

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/707418/youichirou-ootsuka-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

55
papers

1,385
citations

25
h-index

35
g-index

56
ext. papers

1,499
ext. citations

3.6
avg, IF

4.69
L-index

#	Paper	IF	Citations
55	Multifaceted roles of orexin neurons in mediating methamphetamine-induced changes in body temperature and heart rate.. <i>IBRO Neuroscience Reports</i> , 2022 , 12, 108-120		0
54	Inhibition of the dorsomedial hypothalamus substantially decreases brown adipose tissue sympathetic discharge induced by activation of the lateral habenula. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2021 , 230, 102745	2.4	2
53	Alpha-adrenergic receptor agonists prevent emotional hyperthermia. <i>Brain Research</i> , 2020 , 1732, 146678-7	3.7	1
52	Blockade of 5-HT _{2A} receptors inhibits emotional hyperthermia in mice. <i>Journal of Physiological Sciences</i> , 2019 , 69, 1097-1102	2.3	2
51	Activating dopamine D ₂ receptors reduces brown adipose tissue thermogenesis induced by psychological stress and by activation of the lateral habenula. <i>Scientific Reports</i> , 2019 , 9, 19512	4.9	5
50	Neurons in ventral tegmental area tonically inhibit sympathetic outflow to brown adipose tissue: possible mediation of thermogenic signals from lateral habenula. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2019 , 316, R6-R12	3.2	3
49	Inactivation of Serotonergic Neurons in the Rostral Medullary Raph \square Attenuates Stress-Induced Tachypnea and Tachycardia in Mice. <i>Frontiers in Physiology</i> , 2018 , 9, 832	4.6	9
48	Clozapine, chlorpromazine and risperidone dose-dependently reduce emotional hyperthermia, a biological marker of salience. <i>Psychopharmacology</i> , 2017 , 234, 3259-3269	4.7	7
47	Lateral habenula regulation of emotional hyperthermia: mediation via the medullary raph \square <i>Scientific Reports</i> , 2017 , 7, 4102	4.9	14
46	Body Temperature Measurements for Metabolic Phenotyping in Mice. <i>Frontiers in Physiology</i> , 2017 , 8, 520	4.6	54
45	The integrated ultradian organization of behavior and physiology in mice and the contribution of orexin to the ultradian patterning. <i>Neuroscience</i> , 2016 , 334, 119-133	3.9	12
44	Attenuated cold defense responses in orexin neuron-ablated rats. <i>Temperature</i> , 2016 , 3, 465-475	5.2	11
43	Locus coeruleus noradrenergic innervation of the amygdala facilitates alerting-induced constriction of the rat tail artery. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2016 , 310, R1109-19	3.2	6
42	Timing of activities of daily life is jaggy: How episodic ultradian changes in body and brain temperature are integrated into this process. <i>Temperature</i> , 2016 , 3, 371-383	5.2	28
41	Control of cutaneous blood flow by central nervous system. <i>Temperature</i> , 2015 , 2, 392-405	5.2	43
40	Activation of the habenula complex evokes autonomic physiological responses similar to those associated with emotional stress. <i>Physiological Reports</i> , 2015 , 3, e12297	2.6	19
39	Brown adipose tissue thermogenesis contributes to emotional hyperthermia in a resident rat suddenly confronted with an intruder rat. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2014 , 306, R394-400	3.2	33

38	Reduced brown adipose tissue thermogenesis during environmental interactions in transgenic rats with ataxin-3-mediated ablation of hypothalamic orexin neurons. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2014 , 307, R978-89	3.2	24
37	Brown adipose tissue thermogenesis, the basic rest-activity cycle, meal initiation, and bodily homeostasis in rats. <i>Physiology and Behavior</i> , 2013 , 121, 61-9	3.5	24
36	Brown adipose tissue thermogenesis precedes food intake in genetically obese Zucker (fa/fa) rats. <i>Physiology and Behavior</i> , 2013 , 118, 129-37	3.5	5
35	Inactivation of neuronal function in the amygdaloid region reduces tail artery blood flow alerting responses in conscious rats. <i>Neuroscience</i> , 2013 , 228, 13-22	3.9	16
34	TRPA1 detects environmental chemicals and induces avoidance behavior and arousal from sleep. <i>Scientific Reports</i> , 2013 , 3, 3100	4.9	18
33	The impact of hypothermia on emergence from isoflurane anesthesia in orexin neuron-ablated mice. <i>Anesthesia and Analgesia</i> , 2013 , 116, 1001-1005	3.9	9
32	Heating and eating: brown adipose tissue thermogenesis precedes food ingestion as part of the ultradian basic rest-activity cycle in rats. <i>Physiology and Behavior</i> , 2012 , 105, 966-74	3.5	44
31	SR59230A, a beta-3 adrenoceptor antagonist, inhibits ultradian brown adipose tissue thermogenesis and interrupts associated episodic brain and body heating. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2011 , 301, R987-94	3.2	23
30	Atypical antipsychotics cause an acute increase in cutaneous hand blood flow in patients with schizophrenia and schizoaffective disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2011 , 45, 646-53	2.6	6
29	Sympathetic cutaneous vasomotor alerting responses (SCVARs) are associated with hippocampal theta rhythm in non-moving conscious rats. <i>Brain Research</i> , 2009 , 1298, 123-30	3.7	24
28	Brown adipose tissue thermogenesis heats brain and body as part of the brain-coordinated ultradian basic rest-activity cycle. <i>Neuroscience</i> , 2009 , 164, 849-61	3.9	74
27	When administered to rats in a cold environment, 3,4-methylenedioxymethamphetamine reduces brown adipose tissue thermogenesis and increases tail blood flow: effects of pretreatment with 5-HT1A and dopamine D2 antagonists. <i>Neuroscience</i> , 2008 , 154, 1619-26	3.9	26
26	Selective blockade of 5-HT2A receptors attenuates the increased temperature response in brown adipose tissue to restraint stress in rats. <i>Stress</i> , 2008 , 11, 125-33	3	41
25	Fever response to intravenous prostaglandin E2 is mediated by the brain but does not require afferent vagal signaling. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2008 , 294, R1294-303	3.2	45
24	Bursts of brown adipose tissue (BAT) metabolism underlie periodic fluctuations in body temperature (Tb) in rats: a newly discovered ultradian rhythm. <i>FASEB Journal</i> , 2008 , 22, 956.4	0.9	
23	Independent vasomotor control of rat tail and proximal hairy skin. <i>Journal of Physiology</i> , 2007 , 582, 421-33	3.3	31
22	Activation of dopamine D2 receptors in the CNS inhibits sympathetic cutaneous vasomotor alerting responses (SCVARs), contributing to clozapine's SCVAR-inhibiting action. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2007 , 31, 328-36	5.5	13
21	Dopamine D2 receptor stimulation inhibits cold-initiated thermogenesis in brown adipose tissue in conscious rats. <i>Neuroscience</i> , 2007 , 147, 127-35	3.9	32

20	Activation of 5-HT _{1A} receptors in rostral medullary raphe inhibits cutaneous vasoconstriction elicited by cold exposure in rabbits. <i>Brain Research</i> , 2006 , 1073-1074, 252-61	3.7	32
19	Comparison between two rat sympathetic pathways activated in cold defense. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2006 , 291, R589-95	3.2	45
18	Effect of peripheral administration of leptin on the renal sympathetic nerve activity in high-fat diet-related hypertensive rats. <i>Life Sciences</i> , 2006 , 78, 1149-54	6.8	9
17	Thermogenesis in brown adipose tissue: increase by 5-HT _{2A} receptor activation and decrease by 5-HT _{1A} receptor activation in conscious rats. <i>Neuroscience Letters</i> , 2006 , 395, 170-4	3.3	46
16	Clozapine reverses increased brown adipose tissue thermogenesis induced by 3,4-methylenedioxymethamphetamine and by cold exposure in conscious rats. <i>Neuroscience</i> , 2006 , 141, 2067-73	3.9	42
15	Inhibition of medullary raphe parapyramidal neurons prevents cutaneous vasoconstriction elicited by alerting stimuli and by cold exposure in conscious rabbits. <i>Brain Research</i> , 2005 , 1051, 189-93	3.7	33
14	Activation of slowly conducting medullary raphe-spinal neurons, including serotonergic neurons, increases cutaneous sympathetic vasomotor discharge in rabbit. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2005 , 288, R909-18	3.2	34
13	Activation of 5-HT _{1A} receptors in the medullary raphe reduces cardiovascular changes elicited by acute psychological and inflammatory stresses in rabbits. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2005 , 289, R596-R604	3.2	59
12	Spinal 5-HT _{2A} receptors regulate cutaneous sympathetic vasomotor outflow in rabbits and rats; relevance for cutaneous vasoconstriction elicited by MDMA (3,4-methylenedioxymethamphetamine, "Ecstasy") and its reversal by clozapine. <i>Brain Research</i> , 2004 , 1014, 34-44	3.7	40
11	Inhibition of rostral medullary raphe neurons prevents cold-induced activity in sympathetic nerves to rat tail and rabbit ear arteries. <i>Neuroscience Letters</i> , 2004 , 357, 58-62	3.3	71
10	5-Hydroxytryptamine 1A receptors inhibit cold-induced sympathetically mediated cutaneous vasoconstriction in rabbits. <i>Journal of Physiology</i> , 2003 , 552, 303-14	3.9	44
9	Direct measurement of renal sympathetic nervous activity in high-fat diet-related hypertensive rats. <i>Life Sciences</i> , 2002 , 71, 537-46	6.8	39
8	Rhythmic activities of the sympatho-excitatory neurons in the medulla of rabbits: neurons controlling cutaneous vasomotion. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2002 , 101, 48-59	2.4	15
7	Different cardiovascular neuron groups in the ventral reticular formation of the rostral medulla in rabbits: single neurone studies. <i>Journal of the Autonomic Nervous System</i> , 2000 , 79, 74-83		10
6	Leptin injection into white adipose tissue elevates renal sympathetic nerve activity dose-dependently through the afferent nerves pathway in rats. <i>Neuroscience Letters</i> , 2000 , 293, 107-10	3.3	52
5	Functional significance of the 10 Hz rhythmic discharges in sympathetic nerves. <i>Clinical and Experimental Pharmacology and Physiology</i> , 1998 , 25, 464-7	3	3
4	Functionally different neurons are organized topographically in the rostral ventrolateral medulla of rabbits. <i>Journal of the Autonomic Nervous System</i> , 1997 , 67, 67-78		34
3	The spinally mediated 10-Hz rhythm in the sympathetic nerve activity of cats. <i>Journal of the Autonomic Nervous System</i> , 1995 , 54, 89-103		18

- 2 The 10-Hz rhythm in the sympathetic nerve activity of cats, rats and rabbits. *Neuroscience Letters*, **1995**, 196, 173-6 33 11
- 1 Neurons in the caudal ventrolateral medulla mediate the somato-sympathetic inhibitory reflex response via GABA receptors in the rostral ventrolateral medulla. *Journal of the Autonomic Nervous System*, **1992**, 40, 91-8 44