

Ewan K A Millar

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7074844/publications.pdf>

Version: 2024-02-01

63
papers

3,702
citations

109321

35
h-index

133252

59
g-index

65
all docs

65
docs citations

65
times ranked

7073
citing authors

#	ARTICLE	IF	CITATIONS
1	Prediction of Local Recurrence, Distant Metastases, and Death After Breast-Conserving Therapy in Early-Stage Invasive Breast Cancer Using a Five-Biomarker Panel. <i>Journal of Clinical Oncology</i> , 2009, 27, 4701-4708.	1.6	281
2	PI3K pathway activation in breast cancer is associated with the basal-like phenotype and cancer-specific mortality. <i>International Journal of Cancer</i> , 2010, 126, 1121-1131.	5.1	254
3	Inositol polyphosphate 4-phosphatase II regulates PI3K/Akt signaling and is lost in human basal-like breast cancers. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 22231-22236.	7.1	249
4	The key hypoxia regulated gene CAIX is upregulated in basal-like breast tumours and is associated with resistance to chemotherapy. <i>British Journal of Cancer</i> , 2009, 100, 405-411.	6.4	180
5	Hedgehog Overexpression Is Associated with Stromal Interactions and Predicts for Poor Outcome in Breast Cancer. <i>Cancer Research</i> , 2011, 71, 4002-4014.	0.9	149
6	Myopericytoma: a unifying term for a spectrum of tumours that show overlapping features with myofibroma. A review of 14 cases. <i>Journal of Clinical Pathology</i> , 2006, 59, 67-73.	2.0	146
7	Recruitment of regulatory T cells is correlated with hypoxia-induced CXCR4 expression, and is associated with poor prognosis in basal-like breast cancers. <i>Breast Cancer Research</i> , 2011, 13, R47.	5.0	146
8	Cytoplasmic Localization of β -Catenin is a Marker of Poor Outcome in Breast Cancer Patients. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2010, 19, 301-309.	2.5	139
9	Therapeutic targets in triple negative breast cancer. <i>Journal of Clinical Pathology</i> , 2013, 66, 530-542.	2.0	117
10	Transcription factor ATF3 links host adaptive response to breast cancer metastasis. <i>Journal of Clinical Investigation</i> , 2013, 123, 2893-2906.	8.2	109
11	DNA methylation of oestrogen-regulated enhancers defines endocrine sensitivity in breast cancer. <i>Nature Communications</i> , 2015, 6, 7758.	12.8	105
12	miR-139-5p Modulates Radiotherapy Resistance in Breast Cancer by Repressing Multiple Gene Networks of DNA Repair and ROS Defense. <i>Cancer Research</i> , 2018, 78, 501-515.	0.9	105
13	Proteomic Analysis of Urine to Identify Breast Cancer Biomarker Candidates Using a Label-Free LC-MS/MS Approach. <i>PLoS ONE</i> , 2015, 10, e0141876.	2.5	87
14	Enhanced RAD21 cohesin expression confers poor prognosis and resistance to chemotherapy in high grade luminal, basal and HER2 breast cancers. <i>Breast Cancer Research</i> , 2011, 13, R9.	5.0	83
15	Cyclin D1b protein expression in breast cancer is independent of cyclin D1a and associated with poor disease outcome. <i>Oncogene</i> , 2009, 28, 1812-1820.	5.9	81
16	Malignant phyllodes tumours of the breast display increased stromal p53 protein expression. <i>Histopathology</i> , 1999, 34, 491-496.	2.9	72
17	c-Myc overexpression and endocrine resistance in breast cancer. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2006, 102, 147-155.	2.5	71
18	Cyclin D1b Is Aberrantly Regulated in Response to Therapeutic Challenge and Promotes Resistance to Estrogen Antagonists. <i>Cancer Research</i> , 2008, 68, 5628-5638.	0.9	65

#	ARTICLE	IF	CITATIONS
19	Î²-Catenin Signaling Is a Critical Event in ErbB2-Mediated Mammary Tumor Progression. <i>Cancer Research</i> , 2013, 73, 4474-4487.	0.9	65
20	The Magnitude of Androgen Receptor Positivity in Breast Cancer Is Critical for Reliable Prediction of Disease Outcome. <i>Clinical Cancer Research</i> , 2018, 24, 2328-2341.	7.0	63
21	MCL-1 inhibition provides a new way to suppress breast cancer metastasis and increase sensitivity to dasatinib. <i>Breast Cancer Research</i> , 2016, 18, 125.	5.0	60
22	The impact of breast cosmetic and functional outcomes on quality of life: long-term results from the St. George and Wollongong randomized breast boost trial. <i>Breast Cancer Research and Treatment</i> , 2013, 139, 115-123.	2.5	59
23	ELF5 Drives Lung Metastasis in Luminal Breast Cancer through Recruitment of Gr1+ CD11b+ Myeloid-Derived Suppressor Cells. <i>PLoS Biology</i> , 2015, 13, e1002330.	5.6	59
24	High Notch1 protein expression is an early event in breast cancer development and is associated with the HER2 molecular subtype. <i>Histopathology</i> , 2010, 56, 286-296.	2.9	51
25	Prediction of outcome of early ER+ breast cancer is improved using a biomarker panel, which includes Ki-67 and p53. <i>British Journal of Cancer</i> , 2011, 105, 272-280.	6.4	50
26	Prognostic interaction between expression of p53 and estrogen receptor in patients with node-negative breast cancer: results from IBCSG Trials VIII and IX. <i>Breast Cancer Research</i> , 2012, 14, R143.	5.0	50
27	Breast ductal carcinoma in situ carry mutational driver events representative of invasive breast cancer. <i>Modern Pathology</i> , 2017, 30, 952-963.	5.5	50
28	ID4 controls mammary stem cells and marks breast cancers with a stem cell-like phenotype. <i>Nature Communications</i> , 2015, 6, 6548.	12.8	49
29	The expression of the ubiquitin ligase SIAH2 (seven in absentia homolog 2) is mediated through gene copy number in breast cancer and is associated with a basal-like phenotype and p53 expression. <i>Breast Cancer Research</i> , 2011, 13, R19.	5.0	45
30	Andy's Algorithms: new automated digital image analysis pipelines for Fiji. <i>Scientific Reports</i> , 2017, 7, 15717.	3.3	45
31	MASTL overexpression promotes chromosome instability and metastasis in breast cancer. <i>Oncogene</i> , 2018, 37, 4518-4533.	5.9	45
32	Gene expression profiling in breast cancer: towards individualising patient management. <i>Pathology</i> , 2005, 37, 271-277.	0.6	41
33	Tumour Stroma Ratio Assessment Using Digital Image Analysis Predicts Survival in Triple Negative and Luminal Breast Cancer. <i>Cancers</i> , 2020, 12, 3749.	3.7	39
34	BAG-1 predicts patient outcome and tamoxifen responsiveness in ER-positive invasive ductal carcinoma of the breast. <i>British Journal of Cancer</i> , 2009, 100, 123-133.	6.4	37
35	Global characterization of signalling networks associated with tamoxifen resistance in breast cancer. <i>FEBS Journal</i> , 2013, 280, 5237-5257.	4.7	36
36	Supraclavicular radiotherapy must be limited laterally by the coracoid to avoid significant adjuvant breast nodal radiotherapy lymphoedema risk. <i>Journal of Medical Imaging and Radiation Oncology</i> , 2006, 50, 578-582.	0.6	32

#	ARTICLE	IF	CITATIONS
37	Overexpression of the oncogenic signal transducer Gab2 occurs early in breast cancer development. <i>International Journal of Cancer</i> , 2010, 127, 1486-1492.	5.1	31
38	Proteomics for Breast Cancer Urine Biomarkers. <i>Advances in Clinical Chemistry</i> , 2014, 63, 123-167.	3.7	30
39	An ErbB2/c-Src axis links bioenergetics with PRC2 translation to drive epigenetic reprogramming and mammary tumorigenesis. <i>Nature Communications</i> , 2019, 10, 2901.	12.8	24
40	Extramedullary haematopoiesis in axillary lymph nodes following neoadjuvant chemotherapy for locally advanced breast cancer—a potential diagnostic pitfall. <i>Histopathology</i> , 2009, 54, 622-623.	2.9	22
41	Identification of PUMA as an estrogen target gene that mediates the apoptotic response to tamoxifen in human breast cancer cells and predicts patient outcome and tamoxifen responsiveness in breast cancer. <i>Oncogene</i> , 2011, 30, 3186-3197.	5.9	21
42	Î±VÎ²6 integrin expression in diseased and transplanted kidneys. <i>Kidney International</i> , 2004, 66, 1423-1433.	5.2	18
43	Methylation profiling of ductal carcinoma in situ and its relationship to histopathological features. <i>Breast Cancer Research</i> , 2014, 16, 423.	5.0	18
44	MicroRNA-Related DNA Repair/Cell-Cycle Genes Independently Associated With Relapse After Radiation Therapy for Early Breast Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2015, 93, 1104-1114.	0.8	18
45	Elevated levels of tumour apolipoprotein D independently predict poor outcome in breast cancer patients. <i>Histopathology</i> , 2020, 76, 976-987.	2.9	18
46	Id Proteins Promote a Cancer Stem Cell Phenotype in Mouse Models of Triple Negative Breast Cancer via Negative Regulation of Robo1. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 552.	3.7	18
47	TILs Immunophenotype in Breast Cancer Predicts Local Failure and Overall Survival: Analysis in a Large Radiotherapy Trial with Long-Term Follow-Up. <i>Cancers</i> , 2020, 12, 2365.	3.7	18
48	Multiplexed immunofluorescence identifies high stromal CD68+PD-L1+ macrophages as a predictor of improved survival in triple negative breast cancer. <i>Scientific Reports</i> , 2021, 11, 21608.	3.3	16
49	Molecular assays in breast cancer pathology. <i>Pathology</i> , 2011, 43, 116-127.	0.6	15
50	p27KIP-1/cyclin A and cyclin D1 protein expression in ductal carcinoma in situ of the breast: p27KIP-1 correlates with hormone receptor status but not with local recurrence. <i>Pathology International</i> , 2007, 57, 183-189.	1.3	14
51	Deletion of the Antiphospholipid Syndrome Autoantigen Î² ₂ â€Glycoprotein I Potentiates the Lupus Autoimmune Phenotype in a Tollâ€like Receptor 7â€Mediated Murine Model. <i>Arthritis and Rheumatology</i> , 2014, 66, 2270-2280.	5.6	14
52	Assessment of DNA methylation profiling and copy number variation as indications of clonal relationship in ipsilateral and contralateral breast cancers to distinguish recurrent breast cancer from a second primary tumour. <i>BMC Cancer</i> , 2015, 15, 669.	2.6	14
53	Prostatic adenocarcinoma metastatic to the palatine tonsil: a case report. <i>Journal of Laryngology and Otolaryngology</i> , 1994, 108, 178-180.	0.8	13
54	LRH-1 expression patterns in breast cancer tissues are associated with tumour aggressiveness. <i>Oncotarget</i> , 2017, 8, 83626-83636.	1.8	13

#	ARTICLE	IF	CITATIONS
55	Loss of STARD10 expression identifies a group of poor prognosis breast cancers independent of HER2/Neu and triple negative status. <i>International Journal of Cancer</i> , 2010, 126, 1445-1453.	5.1	11
56	Mucinous differentiation in colonic adenocarcinoma is associated with a reduction in tumour-infiltrating lymphocytes. <i>European Journal of Surgical Oncology</i> , 2001, 27, 273-277.	1.0	9
57	Primary Renal Neuroblastoma in Adults. <i>Urology</i> , 2013, 82, 11-13.	1.0	8
58	Significance and Assessment of Margin Status in Ductal Carcinoma In Situ of the Breast. <i>Advances in Anatomic Pathology</i> , 2001, 8, 338-344.	4.3	7
59	Fecal DNA Virome Is Associated with the Development of Colorectal Neoplasia in a Murine Model of Colorectal Cancer. <i>Pathogens</i> , 2022, 11, 457.	2.8	7
60	ALTEN: A High-Fidelity Primary Tissue-Engineering Platform to Assess Cellular Responses Ex Vivo. <i>Advanced Science</i> , 0, , 2103332.	11.2	3
61	Metachronous bilateral primary low-grade mucosa-associated lymphoid tissue non-Hodgkins lymphoma of the breast. <i>Asia-Pacific Journal of Clinical Oncology</i> , 2009, 5, 154-158.	1.1	0
62	A case of amoebic colitis following remote historical exposure. <i>ANZ Journal of Surgery</i> , 2019, 89, E222-E223.	0.7	0
63	Abstract 2962: LRH-1 expression in breast cancer tissue and its association with phenotype and DNA methylation. , 2015, , .		0