

# Stine Regin Wiegell

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7078157/publications.pdf>

Version: 2024-02-01

23  
papers

1,993  
citations

430874

18  
h-index

642732

23  
g-index

23  
all docs

23  
docs citations

23  
times ranked

791  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Treatment of Familial Benign Chronic Pemphigus With Superficial Radiotherapy. JAMA Dermatology, 2022, , .   | 4.1 | 1         |
| 2  | Dyskeratosis follicularis cured by superficial radiotherapy: long-term follow-up of 10 patients. Journal of the European Academy of Dermatology and Venereology, 2021, 35, e230-e232.   | 2.4 | 1         |
| 3  | Electrochemotherapy with bleomycin for basal cell carcinomas: a systematic review. Journal of the European Academy of Dermatology and Venereology, 2021, 35, 2208-2215.   | 2.4 | 11        |
| 4  | Increased protoporphyrin IX accumulation does not improve the effect of photodynamic therapy for actinic keratosis: a randomized controlled trial. British Journal of Dermatology, 2017, 176, 1241-1246.  | 1.5 | 19        |
| 5  | Pulse photodynamic therapy reduces inflammation without compromising efficacy in the treatment of multiple mild actinic keratoses of the face and scalp: a randomized clinical trial. British Journal of Dermatology, 2016, 174, 979-984.   | 1.5 | 29        |
| 6  | Short-term chemical pretreatment cannot replace curettage in photodynamic therapy. Photodermatology Photoimmunology and Photomedicine, 2016, 32, 146-152.   | 1.5 | 17        |
| 7  | Update on Photodynamic Treatment for Actinic Keratosis. Current Problems in Dermatology, 2014, 46, 122-128.   | 0.7 | 39        |
| 8  | Topical corticosteroid reduces inflammation without compromising the efficacy of photodynamic therapy for actinic keratoses: a randomized clinical trial. British Journal of Dermatology, 2014, 171, 1487-1492.   | 1.5 | 28        |
| 9  | Light protection of the skin after photodynamic therapy reduces inflammation: an unblinded randomized controlled study. British Journal of Dermatology, 2014, 171, 175-178.   | 1.5 | 33        |
| 10 | Daylight-mediated photodynamic therapy of basal cell carcinomas – an explorative study. Journal of the European Academy of Dermatology and Venereology, 2014, 28, 169-175.  | 2.4 | 64        |
| 11 | Weather conditions and daylight-mediated photodynamic therapy: protoporphyrin IX-weighted daylight doses measured in six geographical locations. British Journal of Dermatology, 2013, 168, 186-191.  | 1.5 | 95        |
| 12 | Daylight photodynamic therapy for actinic keratosis: an international consensus. Journal of the European Academy of Dermatology and Venereology, 2012, 26, 673-679.   | 2.4 | 189       |
| 13 | Daylight-mediated photodynamic therapy of moderate to thick actinic keratoses of the face and scalp: a randomized multicentre study. British Journal of Dermatology, 2012, 166, 1327-1332.  | 1.5 | 131       |
| 14 | A randomized, multicentre study of directed daylight exposure times of 1½ vs. 2½ h in daylight-mediated photodynamic therapy with methyl aminolaevulinate in patients with multiple thin actinic keratoses of the face and scalp. British Journal of Dermatology, 2011, 164, 1083-1090. | 1.5 | 157       |
| 15 | Continuous ultra-low-intensity artificial daylight is not as effective as red LED light in photodynamic therapy of multiple actinic keratoses. Photodermatology Photoimmunology and Photomedicine, 2011, 27, 280-285.   | 1.5 | 44        |
| 16 | Photodynamic therapy of actinic keratoses with 8% and 16% methyl aminolaevulinate and home-based daylight exposure: a double-blinded randomized clinical trial. British Journal of Dermatology, 2009, 160, 1308-1314.   | 1.5 | 158       |
| 17 | Short and limited effect of long-pulsed dye laser alone and in combination with photodynamic therapy for inflammatory rosacea. Journal of the European Academy of Dermatology and Venereology, 2009, 23, 200-201.   | 2.4 | 12        |
| 18 | Continuous activation of PpIX by daylight is as effective as and less painful than conventional photodynamic therapy for actinic keratoses; a randomized, controlled, single-blinded study. British Journal of Dermatology, 2008, 158, 740-746.   | 1.5 | 313       |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Pain during photodynamic therapy is associated with protoporphyrin IX fluorescence and fluence rate. <i>British Journal of Dermatology</i> , 2008, 158, 727-733.                           | 1.5 | 120       |
| 20 | Cold Water and Pauses in Illumination Reduces Pain During Photodynamic Therapy: A Randomized Clinical Study. <i>Acta Dermato-Venereologica</i> , 2008, 89, 145-149.                        | 1.3 | 46        |
| 21 | Photodynamic therapy of acne vulgaris using 5-aminolevulinic acid versus methyl aminolevulinate. <i>Journal of the American Academy of Dermatology</i> , 2006, 54, 647-651.                | 1.2 | 189       |
| 22 | Photodynamic therapy of acne vulgaris using methyl aminolaevulinate: a blinded, randomized, controlled trial. <i>British Journal of Dermatology</i> , 2006, 154, 969-976.                  | 1.5 | 169       |
| 23 | Pain Associated With Photodynamic Therapy Using 5-Aminolevulinic Acid or 5-Aminolevulinic Acid Methylester on Tape-Stripped Normal Skin. <i>Archives of Dermatology</i> , 2003, 139, 1173. | 1.4 | 128       |