

Jonathan P Little

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7083316/jonathan-p-little-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

153
papers

7,912
citations

43
h-index

86
g-index

179
ext. papers

9,336
ext. citations

3.9
avg, IF

6.31
L-index

#	Paper	IF	Citations
153	Physiological adaptations to low-volume, high-intensity interval training in health and disease. <i>Journal of Physiology</i> , 2012 , 590, 1077-84	3.9	863
152	Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. <i>Journal of Physiology</i> , 2006 , 575, 901-11	3.9	639
151	Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. <i>Journal of Applied Physiology</i> , 2011 , 111, 1554-60	3.7	471
150	A practical model of low-volume high-intensity interval training induces mitochondrial biogenesis in human skeletal muscle: potential mechanisms. <i>Journal of Physiology</i> , 2010 , 588, 1011-22	3.9	388
149	Endurance exercise rescues progeroid aging and induces systemic mitochondrial rejuvenation in mtDNA mutator mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 4135-40	11.5	264
148	Exercise increases mitochondrial PGC-1alpha content and promotes nuclear-mitochondrial cross-talk to coordinate mitochondrial biogenesis. <i>Journal of Biological Chemistry</i> , 2011 , 286, 10605-17	5.4	238
147	Low-volume interval training improves muscle oxidative capacity in sedentary adults. <i>Medicine and Science in Sports and Exercise</i> , 2011 , 43, 1849-56	1.2	227
146	An acute bout of high-intensity interval training increases the nuclear abundance of PGC-1 α and activates mitochondrial biogenesis in human skeletal muscle. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2011 , 300, R1303-10	3.2	206
145	Muscle time under tension during resistance exercise stimulates differential muscle protein sub-fractional synthetic responses in men. <i>Journal of Physiology</i> , 2012 , 590, 351-62	3.9	197
144	Cardiorespiratory fitness as a predictor of intestinal microbial diversity and distinct metagenomic functions. <i>Microbiome</i> , 2016 , 4, 42	16.6	189
143	Where does HIT fit? An examination of the affective response to high-intensity intervals in comparison to continuous moderate- and continuous vigorous-intensity exercise in the exercise intensity-affect continuum. <i>PLoS ONE</i> , 2014 , 9, e114541	3.7	186
142	Acute high-intensity interval exercise reduces the postprandial glucose response and prevalence of hyperglycaemia in patients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2012 , 14, 575-7	6.7	141
141	Affective and Enjoyment Responses to High-Intensity Interval Training in Overweight-to-Obese and Insufficiently Active Adults. <i>Journal of Sport and Exercise Psychology</i> , 2015 , 37, 138-49	1.5	129
140	Inflammation and insulin/IGF-1 resistance as the possible link between obesity and neurodegeneration. <i>Journal of Neuroimmunology</i> , 2014 , 273, 8-21	3.5	121
139	Training intensity modulates changes in PGC-1 α and p53 protein content and mitochondrial respiration, but not markers of mitochondrial content in human skeletal muscle. <i>FASEB Journal</i> , 2016 , 30, 959-70	0.9	117
138	Effects of high-intensity interval exercise versus continuous moderate-intensity exercise on postprandial glycemic control assessed by continuous glucose monitoring in obese adults. <i>Applied Physiology, Nutrition and Metabolism</i> , 2014 , 39, 835-41	3	110
137	Acute endurance exercise increases the nuclear abundance of PGC-1alpha in trained human skeletal muscle. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2010 , 298, R912-7	3.2	110

136	IL-6 induced STAT3 signalling is associated with the proliferation of human muscle satellite cells following acute muscle damage. <i>PLoS ONE</i> , 2011 , 6, e17392	3.7	101
135	High-intensity interval training as an efficacious alternative to moderate-intensity continuous training for adults with prediabetes. <i>Journal of Diabetes Research</i> , 2015 , 2015, 191595	3.9	96
134	Low-dose creatine combined with protein during resistance training in older men. <i>Medicine and Science in Sports and Exercise</i> , 2008 , 40, 1645-52	1.2	95
133	Exercise and nutritional interventions for improving aging muscle health. <i>Endocrine</i> , 2012 , 42, 29-38	4	88
132	Extremely low volume, whole-body aerobic-resistance training improves aerobic fitness and muscular endurance in females. <i>Applied Physiology, Nutrition and Metabolism</i> , 2012 , 37, 1124-31	3	79
131	Effect of Red Bull energy drink on repeated Wingate cycle performance and bench-press muscle endurance. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2007 , 17, 433-44	4.4	77
130	Short-term high-intensity interval and moderate-intensity continuous training reduce leukocyte TLR4 in inactive adults at elevated risk of type 2 diabetes. <i>Journal of Applied Physiology</i> , 2015 , 119, 508-16	3.7	74
129	Intermittent and continuous high-intensity exercise training induce similar acute but different chronic muscle adaptations. <i>Experimental Physiology</i> , 2014 , 99, 782-91	2.4	74
128	Dissociation of increases in PGC-1 α and its regulators from exercise intensity and muscle activation following acute exercise. <i>PLoS ONE</i> , 2013 , 8, e71623	3.7	72
127	Carbohydrate feeding during recovery alters the skeletal muscle metabolic response to repeated sessions of high-intensity interval exercise in humans. <i>Journal of Applied Physiology</i> , 2010 , 108, 628-36	3.7	72
126	Mitochondrial adaptations to high-volume exercise training are rapidly reversed after a reduction in training volume in human skeletal muscle. <i>FASEB Journal</i> , 2016 , 30, 3413-3423	0.9	69
125	Genes to predict VO ₂ trainability: a systematic review. <i>BMC Genomics</i> , 2017 , 18, 831	4.5	65
124	A randomized controlled trial of the effects of flaxseed lignan complex on metabolic syndrome composite score and bone mineral in older adults. <i>Applied Physiology, Nutrition and Metabolism</i> , 2009 , 34, 89-98	3	65
123	Physical activity and exercise attenuate neuroinflammation in neurological diseases. <i>Brain Research Bulletin</i> , 2016 , 125, 19-29	3.9	64
122	Effectiveness and safety of high-intensity interval training in patients with type 2 diabetes. <i>Diabetes Spectrum</i> , 2015 , 28, 39-44	1.9	62
121	Nutritional ketone salts increase fat oxidation but impair high-intensity exercise performance in healthy adult males. <i>Applied Physiology, Nutrition and Metabolism</i> , 2017 , 42, 1031-1035	3	60
120	Effect of nutritional interventions and resistance exercise on aging muscle mass and strength. <i>Biogerontology</i> , 2012 , 13, 345-58	4.5	60
119	Resistance exercise and nutrition to counteract muscle wasting. <i>Applied Physiology, Nutrition and Metabolism</i> , 2009 , 34, 817-28	3	58

118	Hyporesponsiveness to the anti-inflammatory action of interleukin-10 in type 2 diabetes. <i>Scientific Reports</i> , 2016 , 6, 21244	4.9	57
117	Changes in mechanisms proposed to mediate fat loss following an acute bout of high-intensity interval and endurance exercise. <i>Applied Physiology, Nutrition and Metabolism</i> , 2013 , 38, 1236-44	3	55
116	Modification of insulin sensitivity and glycemic control by activity and exercise. <i>Medicine and Science in Sports and Exercise</i> , 2013 , 45, 1868-77	1.2	55
115	Exercise-induced mitochondrial p53 repairs mtDNA mutations in mutator mice. <i>Skeletal Muscle</i> , 2016 , 6, 7	5.1	53
114	Insulinotropic and muscle protein synthetic effects of branched-chain amino acids: potential therapy for type 2 diabetes and sarcopenia. <i>Nutrients</i> , 2012 , 4, 1664-78	6.7	49
113	Skeletal muscle and beyond: the role of exercise as a mediator of systemic mitochondrial biogenesis. <i>Applied Physiology, Nutrition and Metabolism</i> , 2011 , 36, 598-607	3	48
112	Mitochondrial transcription factor A (Tfam) is a pro-inflammatory extracellular signaling molecule recognized by brain microglia. <i>Molecular and Cellular Neurosciences</i> , 2014 , 60, 88-96	4.8	46
111	Prior ingestion of exogenous ketone monoester attenuates the glycaemic response to an oral glucose tolerance test in healthy young individuals. <i>Journal of Physiology</i> , 2018 , 596, 1385-1395	3.9	43
110	Impact of a single bout of high-intensity interval exercise and short-term interval training on interleukin-6, FNDC5, and METRNL mRNA expression in human skeletal muscle. <i>Journal of Sport and Health Science</i> , 2018 , 7, 191-196	8.2	43
109	Sprint-interval but not continuous exercise increases PGC-1 β protein content and p53 phosphorylation in nuclear fractions of human skeletal muscle. <i>Scientific Reports</i> , 2017 , 7, 44227	4.9	42
108	Conjugated linoleic acid combined with creatine monohydrate and whey protein supplementation during strength training. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2009 , 19, 79-96	4.4	42
107	A Multi-Center Comparison of O Trainability Between Interval Training and Moderate Intensity Continuous Training. <i>Frontiers in Physiology</i> , 2019 , 10, 19	4.6	40
106	Globular adiponectin induces a pro-inflammatory response in human astrocytic cells. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 446, 37-42	3.4	35
105	Reduced cortical BACE1 content with one bout of exercise is accompanied by declines in AMPK, Akt, and MAPK signaling in obese, glucose-intolerant mice. <i>Journal of Applied Physiology</i> , 2015 , 119, 1097-1104	7.7	35
104	High Glucose Enhances Neurotoxicity and Inflammatory Cytokine Secretion by Stimulated Human Astrocytes. <i>Current Alzheimer Research</i> , 2017 , 14, 731-741	3	35
103	Acute high-intensity interval exercise reduces human monocyte Toll-like receptor 2 expression in type 2 diabetes. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2017 , 312, R529-R538	3.2	34
102	Moderate-Intensity Exercise and High-Intensity Interval Training Affect Insulin Sensitivity Similarly in Obese Adults. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	34
101	Just HIT it! A time-efficient exercise strategy to improve muscle insulin sensitivity. <i>Journal of Physiology</i> , 2010 , 588, 3341-2	3.9	34

100	Effect of low- and high-glycemic-index meals on metabolism and performance during high-intensity, intermittent exercise. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2010 , 20, 447-56	4.4	34
99	Markers of skeletal muscle mitochondrial function and lipid accumulation are moderately associated with the homeostasis model assessment index of insulin resistance in obese men. <i>PLoS ONE</i> , 2013 , 8, e66322	3.7	34
98	The saturated fatty acid palmitate induces human monocytic cell toxicity toward neuronal cells: exploring a possible link between obesity-related metabolic impairments and neuroinflammation. <i>Journal of Alzheimer's Disease</i> , 2012 , 30 Suppl 2, S179-83	4.3	33
97	Insulin Modulates In Vitro Secretion of Cytokines and Cytotoxins by Human Glial Cells. <i>Current Alzheimer Research</i> , 2015 , 12, 684-93	3	33
96	Short-Term Exercise Training Alters Leukocyte Chemokine Receptors in Obese Adults. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 1631-1640	1.2	31
95	Microparticles: a new perspective in central nervous system disorders. <i>BioMed Research International</i> , 2014 , 2014, 756327	3	31
94	Impact of high-intensity interval duration on perceived exertion. <i>Medicine and Science in Sports and Exercise</i> , 2015 , 47, 1038-45	1.2	30
93	A ketone monoester drink reduces the glycemic response to an oral glucose challenge in individuals with obesity: a randomized trial. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1491-1501	7	30
92	HIGH-INTENSITY INTERVAL TRAINING. <i>ACSM's Health and Fitness Journal</i> , 2014 , 18, 11-16	0.9	29
91	The effect of a short-term low-carbohydrate, high-fat diet with or without postmeal walks on glycemic control and inflammation in type 2 diabetes: a randomized trial. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2018 , 315, R1210-R1219	3.2	28
90	Do stair climbing exercise "snacks" improve cardiorespiratory fitness?. <i>Applied Physiology, Nutrition and Metabolism</i> , 2019 , 44, 681-684	3	27
89	Physiological basis of brief vigorous exercise to improve health. <i>Journal of Physiology</i> , 2020 , 598, 61-69	3.9	26
88	Short-term exercise training reduces anti-inflammatory action of interleukin-10 in adults with obesity. <i>Cytokine</i> , 2018 , 111, 460-469	4	25
87	Differential impact of acute high-intensity exercise on circulating endothelial microparticles and insulin resistance between overweight/obese males and females. <i>PLoS ONE</i> , 2015 , 10, e0115860	3.7	25
86	The effects of low- and high-glycemic index foods on high-intensity intermittent exercise. <i>International Journal of Sports Physiology and Performance</i> , 2009 , 4, 367-80	3.5	25
85	High-Intensity Interval Training Improves Markers of Oxidative Metabolism in Skeletal Muscle of Individuals With Obesity and Insulin Resistance. <i>Frontiers in Physiology</i> , 2018 , 9, 1451	4.6	25
84	Resistance-based interval exercise acutely improves endothelial function in type 2 diabetes. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2016 , 311, H1258-H1267	5.2	24
83	Creatine, arginine alpha-ketoglutarate, amino acids, and medium-chain triglycerides and endurance and performance. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2008 , 18, 493-508	4.4	24

82	Extremely low-volume, high-intensity interval training improves exercise capacity and increases mitochondrial protein content in human skeletal muscle. <i>Open Journal of Molecular and Integrative Physiology</i> , 2013 , 03, 202-210	1	23
81	Commentary: Why sprint interval training is inappropriate for a largely sedentary population. <i>Frontiers in Psychology</i> , 2015 , 6, 1999	3.4	23
80	High- or moderate-intensity training promotes change in cardiorespiratory fitness, but not visceral fat, in obese men: A randomised trial of equal energy expenditure exercise. <i>Respiratory Physiology and Neurobiology</i> , 2019 , 266, 150-155	2.8	21
79	Combined Interval Training and Post-exercise Nutrition in Type 2 Diabetes: A Randomized Control Trial. <i>Frontiers in Physiology</i> , 2017 , 8, 528	4.6	21
78	Self-Monitoring Using Continuous Glucose Monitors with Real-Time Feedback Improves Exercise Adherence in Individuals with Impaired Blood Glucose: A Pilot Study. <i>Diabetes Technology and Therapeutics</i> , 2016 , 18, 185-93	8.1	20
77	High-intensity interval training for improving postprandial hyperglycemia. <i>Research Quarterly for Exercise and Sport</i> , 2014 , 85, 451-6	1.9	20
76	DXA-derived abdominal fat mass, waist circumference, and blood lipids in postmenopausal women. <i>Obesity</i> , 2009 , 17, 1635-40	8	20
75	Elevated Interleukin-10 Levels in COVID-19: Potentiation of Pro-Inflammatory Responses or Impaired Anti-Inflammatory Action?. <i>Frontiers in Immunology</i> , 2021 , 12, 677008	8.4	20
74	Restricting carbohydrates at breakfast is sufficient to reduce 24-hour exposure to postprandial hyperglycemia and improve glycemic variability. <i>American Journal of Clinical Nutrition</i> , 2019 , 109, 1302-1309	7.09	18
73	Supplements and Nutritional Interventions to Augment High-Intensity Interval Training Physiological and Performance Adaptations-A Narrative Review. <i>Nutrients</i> , 2020 , 12,	6.7	18
72	Reductions in RIP140 are not required for exercise- and AICAR-mediated increases in skeletal muscle mitochondrial content. <i>Journal of Applied Physiology</i> , 2011 , 111, 688-95	3.7	18
71	Effects of postmeal exercise on postprandial glucose excursions in people with type 2 diabetes treated with add-on hypoglycemic agents. <i>Diabetes Research and Clinical Practice</i> , 2017 , 126, 240-247	7.4	17
70	Oral Ketone Supplementation Acutely Increases Markers of NLRP3 Inflammasome Activation in Human Monocytes. <i>Molecular Nutrition and Food Research</i> , 2019 , 63, e1801171	5.9	17
69	Purinergic responses of calcium-dependent signaling pathways in cultured adult human astrocytes. <i>BMC Neuroscience</i> , 2014 , 15, 18	3.2	17
68	Cardiovascular benefits of combined interval training and post-exercise nutrition in type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2018 , 32, 226-233	3.2	17
67	Postmeal exercise blunts postprandial glucose excursions in people on metformin monotherapy. <i>Journal of Applied Physiology</i> , 2017 , 123, 444-450	3.7	16
66	Secreted phospholipase A(2) group IIA is a neurotoxin released by stimulated human glial cells. <i>Molecular and Cellular Neurosciences</i> , 2012 , 49, 430-8	4.8	16
65	High-Intensity Interval or Continuous Moderate Exercise: A 24-Week Pilot Trial. <i>Medicine and Science in Sports and Exercise</i> , 2018 , 50, 2067-2075	1.2	15

64	Sprint exercise snacks: a novel approach to increase aerobic fitness. <i>European Journal of Applied Physiology</i> , 2019 , 119, 1203-1212	3.4	14
63	Low-volume high-intensity interval training for cardiometabolic health. <i>Journal of Physiology</i> , 2021 ,	3.9	14
62	Impact of high-intensity interval training and moderate-intensity continuous training on resting and postexercise cardiac troponin T concentration. <i>Experimental Physiology</i> , 2018 , 103, 370-380	2.4	14
61	The effect of brief intermittent stair climbing on glycemic control in people with type 2 diabetes: a pilot study. <i>Applied Physiology, Nutrition and Metabolism</i> , 2018 , 43, 969-972	3	13
60	One-week high-fat diet leads to reduced toll-like receptor 2 expression and function in young healthy men. <i>Nutrition Research</i> , 2014 , 34, 1045-51	4	13
59	The effect of low-volume high-intensity interval training on cardiometabolic health and psychological responses in overweight/obese middle-aged men. <i>Journal of Sports Sciences</i> , 2020 , 38, 1997-2004	3.6	12
58	Carbohydrate restriction with postmeal walking effectively mitigates postprandial hyperglycemia and improves endothelial function in type 2 diabetes. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2018 , 314, H105-H113	5.2	12
57	Reduction of AMPK activity and altered MAPKs signalling in peripheral blood mononuclear cells in response to acute glucose ingestion following a short-term high fat diet in young healthy men. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 1209-16	12.7	12
56	Creatine Monohydrate Supplementation Does Not Augment Fitness, Performance, or Body Composition Adaptations in Response to Four Weeks of High-Intensity Interval Training in Young Females. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2017 , 27, 285-292	4.4	12
55	Human adipose tissue conditioned media from lean subjects is protective against H2O2 induced neurotoxicity in human SH-SY5Y neuronal cells. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 1221-31	6.3	11
54	Forty high-intensity interval training sessions blunt exercise-induced changes in the nuclear protein content of PGC-1 α and p53 in human skeletal muscle. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2020 , 318, E224-E236	6	11
53	Does SIRT1 determine exercise-induced skeletal muscle mitochondrial biogenesis: differences between in vitro and in vivo experiments?. <i>Journal of Applied Physiology</i> , 2012 , 112, 926-8	3.7	10
52	Minimal effect of walking before dinner on glycemic responses in type 2 diabetes: outcomes from the multi-site E-PARA DiGM study. <i>Acta Diabetologica</i> , 2019 , 56, 755-765	3.9	9
51	The Impact of Acute Ingestion of a Ketone Monoester Drink on LPS-Stimulated NLRP3 Activation in Humans with Obesity. <i>Nutrients</i> , 2020 , 12,	6.7	9
50	Exercise increases mitochondrial PGC-1 α content and promotes nuclear-mitochondrial cross-talk to coordinate mitochondrial biogenesis. <i>Journal of Biological Chemistry</i> , 2018 , 293, 4953	5.4	9
49	14-Day Ketone Supplementation Lowers Glucose and Improves Vascular Function in Obesity: A Randomized Crossover Trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e1738-e1754	5.6	9
48	Actions of the anti-angiogenic compound angiostatin in an animal model of Alzheimer's disease. <i>Current Alzheimer Research</i> , 2013 , 10, 252-60	3	9
47	Brief Exercise Counseling and High-Intensity Interval Training on Physical Activity Adherence and Cardiometabolic Health in Individuals at Risk of Type 2 Diabetes: Protocol for a Randomized Controlled Trial. <i>JMIR Research Protocols</i> , 2019 , 8, e11226	2	9

46	Circulating progenitor cells are positively associated with cognitive function among overweight/obese children. <i>Brain, Behavior, and Immunity</i> , 2016 , 57, 47-52	16.6	9
45	Short-term exercise training reduces glycaemic variability and lowers circulating endothelial microparticles in overweight and obese women at elevated risk of type 2 diabetes. <i>European Journal of Sport Science</i> , 2019 , 19, 1140-1149	3.9	9
44	Carbohydrate-Restriction with High-Intensity Interval Training: An Optimal Combination for Treating Metabolic Diseases?. <i>Frontiers in Nutrition</i> , 2017 , 4, 49	6.2	8
43	Walk, Talk and Listen: a pilot randomised controlled trial targeting functional fitness and loneliness in older adults with hearing loss. <i>BMJ Open</i> , 2019 , 9, e026169	3	7
42	The influence of high-intensity interval training and moderate-intensity continuous training on sedentary time in overweight and obese adults. <i>Applied Physiology, Nutrition and Metabolism</i> , 2018 , 43, 747-750	3	7
41	The impact of acute high-intensity interval exercise on biomarkers of cardiovascular health in type 2 diabetes. <i>European Journal of Applied Physiology</i> , 2017 , 117, 1607-1616	3.4	6
40	Short-Term Low-Carbohydrate High-Fat Diet in Healthy Young Males Renders the Endothelium Susceptible to Hyperglycemia-Induced Damage, An Exploratory Analysis. <i>Nutrients</i> , 2019 , 11,	6.7	6
39	Detection of Salivary Insulin Following Low versus High Carbohydrate Meals in Humans. <i>Nutrients</i> , 2017 , 9,	6.7	6
38	The Effect of Exogenous Ketone Monoester Ingestion on Plasma BDNF During an Oral Glucose Tolerance Test. <i>Frontiers in Physiology</i> , 2020 , 11, 1094	4.6	6
37	Effect of carbohydrate-restricted dietary interventions on LDL particle size and number in adults in the context of weight loss or weight maintenance: a systematic review and meta-analysis. <i>American Journal of Clinical Nutrition</i> , 2021 , 114, 1455-1466	7	6
36	Adipose-brain crosstalk: do adipokines have a role in neuroprotection?. <i>Neural Regeneration Research</i> , 2015 , 10, 1381-2	4.5	5
35	Keyto app and device versus WW app on weight loss and metabolic risk in adults with overweight or obesity: A randomized trial. <i>Obesity</i> , 2021 , 29, 1606-1614	8	5
34	Targeting functional fitness, hearing and health-related quality of life in older adults with hearing loss: Walk, Talk & Listen, study protocol for a pilot randomized controlled trial. <i>Trials</i> , 2017 , 18, 47	2.8	4
33	Role of amyloid β in the induction of lipolysis and secretion of adipokines from human adipose tissue. <i>Adipocyte</i> , 2015 , 4, 212-6	3.2	4
32	Glycemic and Metabolic Effects of Two Long Bouts of Moderate-Intensity Exercise in Men with Normal Glucose Tolerance or Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2017 , 8, 154	5.7	4
31	Regulating the regulators: the role of transcriptional regulatory proteins in the adaptive response to exercise in human skeletal muscle. <i>Journal of Physiology</i> , 2011 , 589, 1511-2	3.9	4
30	Potential Therapeutic Effects of Exogenous Ketone Supplementation for Type 2 Diabetes: A Review. <i>Current Pharmaceutical Design</i> , 2020 , 26, 958-969	3.3	4
29	Metabolic Effect of Breaking Up Prolonged Sitting with Stair Climbing Exercise Snacks. <i>Medicine and Science in Sports and Exercise</i> , 2021 , 53, 150-158	1.2	4

28	Prior ingestion of a ketone monoester supplement reduces postprandial glyceimic responses in young healthy-weight individuals. <i>Applied Physiology, Nutrition and Metabolism</i> , 2021 , 46, 309-317	3	4
27	A pragmatic randomized controlled trial testing the effects of the international scientific SCI exercise guidelines on SCI chronic pain: protocol for the EPIC-SCI trial. <i>Spinal Cord</i> , 2020 , 58, 746-754	2.7	3
26	A low-carbohydrate protein-rich bedtime snack to control fasting and nocturnal glucose in type 2 diabetes: A randomized trial. <i>Clinical Nutrition</i> , 2020 , 39, 3601-3606	5.9	3
25	Comparing the Keyto App and Device with Weight WatchersPWW App for Weight Loss: Protocol for a Randomized Trial. <i>JMIR Research Protocols</i> , 2020 , 9, e19053	2	3
24	Comment on Alarcon et al. Pancreatic β Cell Adaptive Plasticity in Obesity Increases Insulin Production but Adversely Affects Secretory Function. <i>Diabetes</i> 2016;65:438-450. <i>Diabetes</i> , 2016 , 65, e28	0.9	3
23	Last word on viewpoint: does SIRT1 determine exercise-induced skeletal muscle mitochondrial biogenesis: differences between in vitro and in vivo experiments?. <i>Journal of Applied Physiology</i> , 2012 , 112, 931	3.7	2
22	Short-term ketone monoester supplementation improves cerebral blood flow and cognition in obesity: A randomized cross-over trial. <i>Journal of Physiology</i> , 2021 , 599, 4763-4778	3.9	2
21	Exercise Snacks: A Novel Strategy to Improve Cardiometabolic Health. <i>Exercise and Sport Sciences Reviews</i> , 2021 , 50,	6.7	2
20	Interpreting Anti-inflammatoryCytokine responses to exercise: focus on interleukin-10. <i>Journal of Physiology</i> , 2021 , 599, 5163-5177	3.9	2
19	Genome wide association study of response to interval and continuous exercise training: the Predict-HIIT study. <i>Journal of Biomedical Science</i> , 2021 , 28, 37	13.3	2
18	Efficacy of Dietary and Supplementation Interventions for Individuals with Type 2 Diabetes. <i>Nutrients</i> , 2021 , 13,	6.7	2
17	Increased cardiorespiratory stress during submaximal cycling after ketone monoester ingestion in endurance-trained adults. <i>Applied Physiology, Nutrition and Metabolism</i> , 2021 , 46, 986-993	3	2
16	Exercise training remodels subcutaneous adipose tissue in adults with obesity even without weight loss.. <i>Journal of Physiology</i> , 2022 ,	3.9	2
15	Reply from M. J. Gibala, J. P. Little, M. J. MadDonald and J. A. Hawley. <i>Journal of Physiology</i> , 2012 , 590, 3391-3391	3.9	1
14	A Pilot Study on In-Task Affect Predicting Free-Living Adherence to HIIT and MICT. <i>Research Quarterly for Exercise and Sport</i> , 2020 , 1-10	1.9	1
13	Hourly staircase sprinting exercise "snacks" improve femoral artery shear patterns but not flow-mediated dilation or cerebrovascular regulation: a pilot study. <i>Applied Physiology, Nutrition and Metabolism</i> , 2021 , 46, 521-529	3	1
12	The effects of a high-fat/high-carbohydrate meal on leukocyte populations in adults with chronic spinal cord injury. <i>Spinal Cord Series and Cases</i> , 2021 , 7, 49	1.4	1
11	Pharmacist-led therapeutic carbohydrate restriction as a treatment strategy for type 2 diabetes: the Pharm-TCR randomized controlled trial protocol. <i>Trials</i> , 2019 , 20, 781	2.8	1

10	Use of an mHealth Ketogenic Diet App Intervention and User Behaviors Associated With Weight Loss in Adults With Overweight or Obesity: Secondary Analysis of a Randomized Clinical Trial.. <i>JMIR MHealth and UHealth</i> , 2022 , 10, e33940	5.5	1
9	Spinal cord injury impairs cardiac function due to impaired bulbospinal sympathetic control.. <i>Nature Communications</i> , 2022 , 13, 1382	17.4	1
8	Examining the Effect of Consuming C Medium-Chain Triglyceride Oil for 14 Days on Markers of NLRP3 Activation in Healthy Humans.. <i>Journal of Nutrition and Metabolism</i> , 2022 , 2022, 7672759	2.7	1
7	A randomized controlled trial of pharmacist-led therapeutic carbohydrate and energy restriction in type 2 diabetes. <i>Nature Communications</i> , 2021 , 12, 5367	17.4	0
6	Exercise Training Protocols to Improve Obesity, Glucose Homeostasis, and Subclinical Inflammation. <i>Methods in Molecular Biology</i> , 2022 , 2343, 119-145	1.4	0
5	The Effects of Creatine and Protein Supplementation on Bone Catabolism in Older Men. <i>Medicine and Science in Sports and Exercise</i> , 2008 , 40, S98	1.2	
4	Insulinotropic and Muscle Protein Synthetic Effects of Branched-Chain Amino Acids: Potential Therapy for Type Diabetes and Sarcopenia 2016 , 87-104		
3	Interval Walking Training Reduces T Cell CCR5 in Individuals with Type 2 Diabetes. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 197-198	1.2	
2	Reductions in RIP140 are not required for exercise and high fat diet mediated increases in mitochondrial enzymes. <i>FASEB Journal</i> , 2011 , 25, 1104.1	0.9	
1	Palmitate and ceramide induce human monocytic cell toxicity towards neuronal cells. <i>FASEB Journal</i> , 2012 , 26, 570.8	0.9	