

Victoria N Karetnikova

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7086076/publications.pdf>

Version: 2024-02-01

30
papers

166
citations

1162367

8
h-index

1125271

13
g-index

34
all docs

34
docs citations

34
times ranked

320
citing authors

#	ARTICLE	IF	CITATIONS
1	New coronavirus disease (COVID-19) and cardiovascular disease. <i>Complex Issues of Cardiovascular Diseases</i> , 2020, 9, 17-28.	0.3	24
2	Multivessel coronary artery disease, free fatty acids, oxidized LDL and its antibody in myocardial infarction. <i>Lipids in Health and Disease</i> , 2014, 13, 111.	1.2	23
3	Prognostic Value of Soluble ST2 During Hospitalization for ST-Segment Elevation Myocardial Infarction. <i>Annals of Laboratory Medicine</i> , 2016, 36, 313-319.	1.2	17
4	Glucose levels as a prognostic marker in patients with ST-segment elevation myocardial infarction: a caseâ€“control study. <i>BMC Endocrine Disorders</i> , 2016, 16, 31.	0.9	17
5	The role of adipose tissue and adipokines in the manifestation of type 2 diabetes in the long-term period following myocardial infarction. <i>Diabetology and Metabolic Syndrome</i> , 2016, 8, 24.	1.2	17
6	Early Effects of Treatment Low-Dose Atorvastatin on Markers of Insulin Resistance and Inflammation in Patients with Myocardial Infarction. <i>Frontiers in Pharmacology</i> , 2016, 7, 324.	1.6	16
7	Pre-diabetes as an interdisciplinary problem: definition, risks, approaches to the diagnostics and prevention of type 2 diabetes and cardiovascular complications. <i>Russian Journal of Cardiology</i> , 2019, , 83-91.	0.4	12
8	Dose-dependent effects of atorvastatin on myocardial infarction. <i>Drug Design, Development and Therapy</i> , 2015, 9, 3361.	2.0	8
9	Relationship key factor of inflammation and the development of complications in the late period of myocardial infarction in patients with visceral obesity. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 36.	0.7	8
10	Serum Galectin and Renal Dysfunction in ST-Segment Elevation Myocardial Infarction. <i>Disease Markers</i> , 2016, 2016, 1-6.	0.6	4
11	Predictors of myocardial fibrosis and loss of epicardial adipose tissue volume in the long-term period after myocardial infarction. <i>Russian Journal of Cardiology</i> , 2020, 25, 31-40.	0.4	4
12	Association of inflammatory markers and poor outcome in diabetic patients presenting with ST segment elevation myocardial infarction. <i>Journal of Inflammation Research</i> , 2015, 8, 107.	1.6	3
13	Fibrosis biomarkers and global myocardial strain in the diagnosis and prediction of diastolic dysfunction in patients with myocardial infarction and preserved ejection fraction. <i>Russian Journal of Cardiology</i> , 2021, 26, 4255.	0.4	2
14	Biological markers and cardiac remodelling following the myocardial infarction. <i>Aging</i> , 2019, 11, 3523-3535.	1.4	2
15	Visceral adiposity index in patients with coronary artery disease, obesity and type 2 diabetes. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2020, 19, 2311.	0.4	2
16	N-terminal propeptide of type III procollagen for predicting diastolic dysfunction in patients with myocardial infarction and preserved ejection fraction. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2021, 20, 2494.	0.4	1
17	The Congress of the American College of Cardiology in review. <i>Complex Issues of Cardiovascular Diseases</i> , 2020, 9, 29-37.	0.3	1
18	The importance of chronic kidney disease for the assessment of risk of adverse outcomes after myocardial infarction. <i>Klinicheskaia Meditsina</i> , 2017, 95, 563-570.	0.2	1

#	ARTICLE	IF	CITATIONS
19	The relationship of the epicardial fat and adipo-fibrokinases in myocardial infarction. <i>Klinicheskaya Laboratornaya Diagnostika</i> , 2020, 65, 533-540.	0.2	1
20	Participation of the C-terminal propeptide procollagen type I in the formation of cardiofibrosis in patients with myocardial infarction with preserved left ventricular ejection fraction. <i>Russian Journal of Cardiology</i> , 2021, 26, 4137.	0.4	0
21	Serum markers of cardiac fibrosis suffering from heart failure with preserved left ventricular ejection fraction upon ST-segment elevation myocardial infarction. <i>Fundamental and Clinical Medicine</i> , 2021, 6, 46-55.	0.1	0
22	CORRELATION OF INFLAMMATION MARKERS AND THE LEVEL OF DIFFUSIONAL CAPACITY OF LUNGS IN PATIENTS WITH MYOCARDIAL INFARCTION. <i>Siberian Medical Review</i> , 2018, , 11-16.	0.1	0
23	POST-INFARCTION REMODELLING AND DIFFUSION CAPACITY OF THE LUNGS: IS THERE ANY CONNECTION?. <i>Siberian Medical Review</i> , 2018, , 66-72.	0.1	0
24	RELATION OF THE MARKERS OF MYOCARDIAL DYSFUNCTION AND SYSTEMIC INFLAMMATION WITH STRUCTURAL AND FUNCTIONAL PARAMETERS OF THE LUNGS IN MYOCARDIAL INFARCTION PATIENTS. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2018, 17, 24-28.	0.4	0
25	The factors influencing the diffusion capacity of the lung in patients with myocardial infarction.. <i>Klinicheskaya Meditsina</i> , 2018, 96, 328-334.	0.2	0
26	Outcomes of long-term observation period in patients with myocardial infarction undergone radiopaque interventions in the acute period of the disease.. <i>Klinicheskaya Meditsina</i> , 2018, 96, 648-657.	0.2	0
27	Prevalence of electrocardiographic changes in Kemerovo region according to the data of the ESSE-RF study. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2019, 18, 120-126.	0.4	0
28	Serum procollagens, structural and functional characteristics of the heart in prediction of cardiac fibrosis after myocardial infarction. <i>Siberian Medical Review</i> , 2020, , 75-82.	0.1	0
29	In-hospital changes of echocardiographic parameters and their relationship with the procollagen I C-terminal propeptide in patients with myocardial infarction and preserved left ventricle systolic function. <i>Russian Journal of Cardiology</i> , 2020, 25, 45-51.	0.4	0
30	Educational approaches to teaching specialist program "Cardiology and cardiovascular surgery" during lockdown. <i>Complex Issues of Cardiovascular Diseases</i> , 2020, 9, 90-95.	0.3	0