

Ileana Ruxandra Botusan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7086583/publications.pdf>

Version: 2024-02-01

16
papers

1,039
citations

687363

13
h-index

996975

15
g-index

17
all docs

17
docs citations

17
times ranked

1915
citing authors

#	ARTICLE	IF	CITATIONS
1	Repression of hypoxia-inducible factor-1 contributes to increased mitochondrial reactive oxygen species production in diabetes. <i>ELife</i> , 2022, 11, .	6.0	31
2	GWAS for autoimmune Addison's disease identifies multiple risk loci and highlights AIRE in disease susceptibility. <i>Nature Communications</i> , 2021, 12, 959.	12.8	33
3	HypoxamiR-210 accelerates wound healing in diabetic mice by improving cellular metabolism. <i>Communications Biology</i> , 2020, 3, 768.	4.4	18
4	Triggering of a Dll4-Notch1 loop impairs wound healing in diabetes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 6985-6994.	7.1	58
5	Deficiency of liver-derived insulin-like growth factor-I (IGF-I) does not interfere with the skin wound healing rate. <i>PLoS ONE</i> , 2018, 13, e0193084.	2.5	15
6	Perlecan heparan sulfate deficiency impairs pulmonary vascular development and attenuates hypoxic pulmonary hypertension. <i>Cardiovascular Research</i> , 2015, 107, 20-31.	3.8	30
7	Hyperbaric oxygen therapy activates hypoxia-inducible factor 1 (HIF-1), which contributes to improved wound healing in diabetic mice. <i>Wound Repair and Regeneration</i> , 2015, 23, 98-103.	3.0	109
8	Impact of the Hypoxia-Inducible Factor-1 $\hat{\pm}$ (<i>HIF1A</i>) Pro582Ser Polymorphism on Diabetes Nephropathy. <i>Diabetes Care</i> , 2013, 36, 415-421.	8.6	56
9	Hypoxia and inflammation synergistically promote bone destruction. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, A61.1-A61.	0.9	0
10	Stability of mitochondrial DNA against reactive oxygen species (ROS) generated in diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2011, 27, 470-479.	4.0	21
11	Hyperglycemia and hypoxia inducible factor (HIF): A multifaceted story. <i>Cell Cycle</i> , 2010, 9, 1856-1856.	2.6	1
12	Stabilization of HIF-1 $\hat{\pm}$ is critical to improve wound healing in diabetic mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 19426-19431.	7.1	416
13	Anti-tumor necrosis factor therapy increases synovial osteoprotegerin expression in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2006, 54, 76-81.	6.7	93
14	Intraarticular corticosteroids decrease synovial RANKL expression in inflammatory arthritis. <i>Arthritis and Rheumatism</i> , 2006, 54, 1463-1472.	6.7	46
15	Hypoxia-Inducible Factor-1 $\hat{\pm}$ and Hypoxia-Inducible Factor-2 $\hat{\pm}$ Are Expressed in Kaposi Sarcoma and Modulated by Insulin-like Growth Factor-I. <i>Clinical Cancer Research</i> , 2006, 12, 4506-4514.	7.0	58
16	The value of an acute octreotide suppression test in predicting long-term responses to depot somatostatin analogues in patients with active acromegaly. <i>Clinical Endocrinology</i> , 2005, 62, 282-288.	2.4	54