

David A Mcdonald

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7087902/publications.pdf>

Version: 2024-02-01

13
papers

698
citations

1040056

9
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

713
citing authors

#	ARTICLE	IF	CITATIONS
1	Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS [®]) Society recommendations. Monthly Notices of the Royal Astronomical Society: Letters, 2020, 91, 3-19.	3.3	348
2	An enhanced recovery programme for primary total knee arthroplasty in the United Kingdom – follow up at one year. Knee, 2012, 19, 525-529.	1.6	118
3	The use of enhanced recovery after surgery (ERAS) principles in Scottish orthopaedic units – an implementation and follow-up at 1 year, 2010 – 2011: a report from the Musculoskeletal Audit, Scotland. Archives of Orthopaedic and Trauma Surgery, 2013, 133, 117-124.	2.4	70
4	Ropivacaine plasma levels following high-dose local infiltration analgesia for total knee arthroplasty. Anaesthesia, 2015, 70, 784-790.	3.8	36
5	Cost comparison of orthopaedic fracture pathways using discrete event simulation in a Glasgow hospital. BMJ Open, 2017, 7, e014509.	1.9	36
6	Level of adherence to prescribed exercise in spondyloarthritis and factors affecting this adherence: a systematic review. Rheumatology International, 2019, 39, 187-201.	3.0	22
7	An anatomic study of local infiltration analgesia in total knee arthroplasty. Knee, 2013, 20, 319-323.	1.6	19
8	The technique of delivery of peri-operative analgesia does not affect the rehabilitation or outcomes following total knee arthroplasty. Bone and Joint Journal, 2016, 98-B, 1189-1196.	4.4	14
9	Ropivacaine plasma levels following local infiltration analgesia for primary total hip arthroplasty. Anaesthesia, 2014, 69, 368-373.	3.8	13
10	Development of an occupational advice intervention for patients undergoing lower limb arthroplasty (the OPAL study). BMC Health Services Research, 2018, 18, 504.	2.2	8
11	Occupational advice to help people return to work following lower limb arthroplasty: the OPAL intervention mapping study. Health Technology Assessment, 2020, 24, 1-408.	2.8	8
12	Upright Time and Sit-To-Stand Transition Progression After Total Hip Arthroplasty: An Inhospital Longitudinal Study. Journal of Arthroplasty, 2016, 31, 735-739.	3.1	5
13	A hospital-based mixed-methods observational study to evaluate a hip and knee replacement quality improvement project. TQM Journal, 2021, 33, 272-294.	3.3	1