

Jason Matthews

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7092743/jason-matthews-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

79
papers

5,157
citations

30
h-index

71
g-index

86
ext. papers

5,795
ext. citations

6.2
avg, IF

5.46
L-index

| # | Paper | IF | Citations |
|----|---|------|-----------|
| 79 | LongITools: Dynamic longitudinal exposome trajectories in cardiovascular and metabolic noncommunicable diseases.. <i>Environmental Epidemiology</i> , 2022 , 6, e184 | 0.2 | 1 |
| 78 | PARP7 and Mono-ADP-Ribosylation Negatively Regulate Estrogen Receptor β Signaling in Human Breast Cancer Cells. <i>Cells</i> , 2021 , 10, | 7.9 | 7 |
| 77 | 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD)-Inducible Poly-ADP-Ribose Polymerase (TIPARP/PARP7) Catalytic Mutant Mice (TiparpH532A) Exhibit Increased Sensitivity to TCDD-Induced Hepatotoxicity and Lethality. <i>Toxicological Sciences</i> , 2021 , 183, 154-169 | 4.4 | 1 |
| 76 | The aryl hydrocarbon receptor reduces LC3II expression and controls endoplasmic reticulum stress. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2021 , 320, L339-L355 | 5.8 | 4 |
| 75 | Chemical genetics and proteome-wide site mapping reveal cysteine MARYlation by PARP-7 on immune-relevant protein targets. <i>ELife</i> , 2021 , 10, | 8.9 | 18 |
| 74 | ADP-ribosyltransferases, an update on function and nomenclature. <i>FEBS Journal</i> , 2021 , | 5.7 | 30 |
| 73 | LXR β Regulates ChREBP β Transactivity in a Target Gene-Specific Manner through an Agonist-Modulated LBD-LID Interaction. <i>Cells</i> , 2020 , 9, | 7.9 | 2 |
| 72 | 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) alters hepatic polyunsaturated fatty acid metabolism and eicosanoid biosynthesis in female Sprague-Dawley rats. <i>Toxicology and Applied Pharmacology</i> , 2020 , 398, 115034 | 4.6 | 2 |
| 71 | Environmental six-ring polycyclic aromatic hydrocarbons are potent inducers of the AhR-dependent signaling in human cells. <i>Environmental Pollution</i> , 2020 , 266, 115125 | 9.3 | 5 |
| 70 | DNA methylation repels binding of hypoxia-inducible transcription factors to maintain tumor immunotolerance. <i>Genome Biology</i> , 2020 , 21, 182 | 18.3 | 13 |
| 69 | Effects of antioxidant-rich foods on altitude-induced oxidative stress and inflammation in elite endurance athletes: A randomized controlled trial. <i>PLoS ONE</i> , 2019 , 14, e0217895 | 3.7 | 18 |
| 68 | 3-Methylcholanthrene Induces Chylous Ascites in TCDD-Inducible Poly-ADP-Ribose Polymerase () Knockout Mice. <i>International Journal of Molecular Sciences</i> , 2019 , 20, | 6.3 | 5 |
| 67 | Shared epitope is associated with the reactivity of Th17 cells to cigarette smoke extract regardless of smoking history. <i>Cellular and Molecular Immunology</i> , 2019 , 16, 674-675 | 15.4 | 1 |
| 66 | Characterization of Epigenetic Histone Activation/Repression Marks in Sequences of Genes by Chromatin Immunoprecipitation-Quantitative Polymerase Chain Reaction (ChIP-qPCR). <i>Methods in Molecular Biology</i> , 2019 , 1965, 389-403 | 1.4 | 6 |
| 65 | Molecular modelling, synthesis, and biological evaluations of a 3,5-disubstituted isoxazole fatty acid analogue as a PPAR β -selective agonist. <i>Bioorganic and Medicinal Chemistry</i> , 2019 , 27, 4059-4068 | 3.4 | 4 |
| 64 | Loss of Tiparp Results in Aberrant Layering of the Cerebral Cortex. <i>ENeuro</i> , 2019 , 6, | 3.9 | 4 |
| 63 | The aryl hydrocarbon receptor regulates the expression of TIPARP and its cis long non-coding RNA, TIPARP-AS1. <i>Biochemical and Biophysical Research Communications</i> , 2018 , 495, 2356-2362 | 3.4 | 11 |

| | | | |
|----|---|------|-----|
| 62 | Genome-wide mapping and analysis of aryl hydrocarbon receptor (AHR)- and aryl hydrocarbon receptor repressor (AHRR)-binding sites in human breast cancer cells. <i>Archives of Toxicology</i> , 2018 , 92, 225-240 | 5.8 | 20 |
| 61 | Methods to Study TCDD-Inducible Poly-ADP-Ribose Polymerase (TIPARP) Mono-ADP-Ribosyltransferase Activity. <i>Methods in Molecular Biology</i> , 2018 , 1813, 109-124 | 1.4 | 3 |
| 60 | Hepatocyte-Specific Deletion of TIPARP, a Negative Regulator of the Aryl Hydrocarbon Receptor, Is Sufficient to Increase Sensitivity to Dioxin-Induced Wasting Syndrome. <i>Toxicological Sciences</i> , 2018 , 165, 347-360 | 4.4 | 15 |
| 59 | Characterization of TCDD-inducible poly-ADP-ribose polymerase (TIPARP/ARTD14) catalytic activity. <i>Biochemical Journal</i> , 2018 , 475, 3827-3846 | 3.8 | 23 |
| 58 | Aryl Hydrocarbon Receptor-Dependent Metabolism Plays a Significant Role in Estrogen-Like Effects of Polycyclic Aromatic Hydrocarbons on Cell Proliferation. <i>Toxicological Sciences</i> , 2018 , 165, 447-461 | 4.4 | 21 |
| 57 | Aryl hydrocarbon receptor (AhR)-dependent regulation of pulmonary miRNA by chronic cigarette smoke exposure. <i>Scientific Reports</i> , 2017 , 7, 40539 | 4.9 | 29 |
| 56 | Convergence of hepcidin deficiency, systemic iron overloading, heme accumulation, and REV-ERB β activation in aryl hydrocarbon receptor-elicited hepatotoxicity. <i>Toxicology and Applied Pharmacology</i> , 2017 , 321, 1-17 | 4.6 | 19 |
| 55 | AHR toxicity and signaling: Role of TIPARP and ADP-ribosylation. <i>Current Opinion in Toxicology</i> , 2017 , 2, 50-57 | 4.4 | 9 |
| 54 | LXR β Regulates Hepatic ChREBP Activity and Lipogenesis upon Glucose, but Not Fructose Feeding in Mice. <i>Nutrients</i> , 2017 , 9, | 6.7 | 12 |
| 53 | Low levels of the AhR in chronic obstructive pulmonary disease (COPD)-derived lung cells increases COX-2 protein by altering mRNA stability. <i>PLoS ONE</i> , 2017 , 12, e0180881 | 3.7 | 7 |
| 52 | Pyruvate Kinase Isoform Switching and Hepatic Metabolic Reprogramming by the Environmental Contaminant 2,3,7,8-Tetrachlorodibenzo-p-Dioxin. <i>Toxicological Sciences</i> , 2016 , 149, 358-71 | 4.4 | 25 |
| 51 | TCDD-inducible poly-ADP-ribose polymerase (TIPARP/PARP7) mono-ADP-ribosylates and co-activates liver X receptors. <i>Biochemical Journal</i> , 2016 , 473, 899-910 | 3.8 | 30 |
| 50 | Constitutive aryl hydrocarbon receptor signaling constrains type I interferon-mediated antiviral innate defense. <i>Nature Immunology</i> , 2016 , 17, 687-94 | 19.1 | 113 |
| 49 | Dose-Dependent Metabolic Reprogramming and Differential Gene Expression in TCDD-Elicited Hepatic Fibrosis. <i>Toxicological Sciences</i> , 2016 , 154, 253-266 | 4.4 | 33 |
| 48 | Liver X receptor regulates hepatic nuclear O-GlcNAc signaling and carbohydrate responsive element-binding protein activity. <i>Journal of Lipid Research</i> , 2015 , 56, 771-85 | 6.3 | 32 |
| 47 | The aryl hydrocarbon receptor suppresses cigarette-smoke-induced oxidative stress in association with dioxin response element (DRE)-independent regulation of sulfiredoxin 1. <i>Free Radical Biology and Medicine</i> , 2015 , 89, 342-57 | 7.8 | 27 |
| 46 | Loss of the Mono-ADP-ribosyltransferase, Tiparp, Increases Sensitivity to Dioxin-induced Steatohepatitis and Lethality. <i>Journal of Biological Chemistry</i> , 2015 , 290, 16824-40 | 5.4 | 37 |
| 45 | Zinc finger nuclease-mediated knockout of AHR or ARNT in human breast cancer cells abolishes basal and ligand-dependent regulation of CYP1B1 and differentially affects estrogen receptor α transactivation. <i>Toxicological Sciences</i> , 2014 , 138, 89-103 | 4.4 | 19 |

| | | | |
|----|--|------|----|
| 44 | Aryl hydrocarbon receptor-dependent regulation of miR-196a expression controls lung fibroblast apoptosis but not proliferation. <i>Toxicology and Applied Pharmacology</i> , 2014 , 280, 511-25 | 4.6 | 32 |
| 43 | Aryl hydrocarbon receptor repressor and TiPARP (ARTD14) use similar, but also distinct mechanisms to repress aryl hydrocarbon receptor signaling. <i>International Journal of Molecular Sciences</i> , 2014 , 15, 7939-57 | 6.3 | 38 |
| 42 | Alternative Negative Feedback Control in the Aryl Hydrocarbon Receptor Signaling Pathway. <i>Journal of Drug Metabolism & Toxicology</i> , 2013 , 04, | | 2 |
| 41 | The aryl hydrocarbon receptor and estrogen receptor alpha differentially modulate nuclear factor erythroid-2-related factor 2 transactivation in MCF-7 breast cancer cells. <i>Toxicology and Applied Pharmacology</i> , 2013 , 270, 139-48 | 4.6 | 33 |
| 40 | Induction of multidrug resistance transporter ABCG2 by prolactin in human breast cancer cells. <i>Molecular Pharmacology</i> , 2013 , 83, 377-88 | 4.3 | 16 |
| 39 | AHR- and ER-Mediated Toxicology and Chemoprevention. <i>Advances in Molecular Toxicology</i> , 2013 , 1-38 | 0.4 | 2 |
| 38 | 2,3,7,8-Tetrachlorodibenzo-p-dioxin poly(ADP-ribose) polymerase (TiPARP, ARTD14) is a mono-ADP-ribosyltransferase and repressor of aryl hydrocarbon receptor transactivation. <i>Nucleic Acids Research</i> , 2013 , 41, 1604-21 | 20.1 | 85 |
| 37 | High-resolution genome-wide mapping of AHR and ARNT binding sites by ChIP-Seq. <i>Toxicological Sciences</i> , 2012 , 130, 349-61 | 4.4 | 84 |
| 36 | FOXA1 is essential for aryl hydrocarbon receptor-dependent regulation of cyclin G2. <i>Molecular Cancer Research</i> , 2012 , 10, 636-48 | 6.6 | 24 |
| 35 | Differential ligand-dependent activation and a role for Y322 in aryl hydrocarbon receptor-mediated regulation of gene expression. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 410, 859-65 | 3.4 | 10 |
| 34 | Identification of aryl hydrocarbon receptor binding targets in mouse hepatic tissue treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. <i>Toxicology and Applied Pharmacology</i> , 2011 , 257, 38-47 | 4.6 | 20 |
| 33 | Integration of genome-wide computation DRE search, AhR ChIP-chip and gene expression analyses of TCDD-elicited responses in the mouse liver. <i>BMC Genomics</i> , 2011 , 12, 365 | 4.5 | 83 |
| 32 | 3-methylcholanthrene induces differential recruitment of aryl hydrocarbon receptor to human promoters. <i>Toxicological Sciences</i> , 2010 , 117, 90-100 | 4.4 | 25 |
| 31 | Inhibition of aryl hydrocarbon receptor-dependent transcription by resveratrol or kaempferol is independent of estrogen receptor expression in human breast cancer cells. <i>Cancer Letters</i> , 2010 , 299, 119-29 | 9.9 | 46 |
| 30 | Estrogen receptor-dependent regulation of CYP2B6 in human breast cancer cells. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2010 , 1799, 469-79 | 6 | 30 |
| 29 | Flavin-containing monooxygenase-3: induction by 3-methylcholanthrene and complex regulation by xenobiotic chemicals in hepatoma cells and mouse liver. <i>Toxicology and Applied Pharmacology</i> , 2010 , 247, 60-9 | 4.6 | 30 |
| 28 | Functional analysis of six human aryl hydrocarbon receptor variants in human breast cancer and mouse hepatoma cell lines. <i>Toxicology</i> , 2010 , 277, 59-65 | 4.4 | 25 |
| 27 | A new class of estrogen receptor beta-selective activators. <i>Molecular Interventions: Pharmacological Perspectives From Biology, Chemistry and Genomics</i> , 2010 , 10, 133-6 | | 12 |

| | | | |
|----|---|------|------|
| 26 | Estrogen receptor subtype- and promoter-specific modulation of aryl hydrocarbon receptor-dependent transcription. <i>Molecular Cancer Research</i> , 2009 , 7, 977-86 | 6.6 | 40 |
| 25 | Activation function 2 mediates dioxin-induced recruitment of estrogen receptor alpha to CYP1A1 and CYP1B1. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 385, 263-8 | 3.4 | 11 |
| 24 | Dioxin increases the interaction between aryl hydrocarbon receptor and estrogen receptor alpha at human promoters. <i>Toxicological Sciences</i> , 2009 , 111, 254-66 | 4.4 | 61 |
| 23 | Aryl hydrocarbon receptor-dependent induction of flavin-containing monooxygenase mRNAs in mouse liver. <i>Drug Metabolism and Disposition</i> , 2008 , 36, 2499-505 | 4 | 38 |
| 22 | Estrogen receptors: how do they signal and what are their targets. <i>Physiological Reviews</i> , 2007 , 87, 905-317.9 | 17.9 | 1266 |
| 21 | Estrogen receptor beta2 negatively regulates the transactivation of estrogen receptor alpha in human breast cancer cells. <i>Cancer Research</i> , 2007 , 67, 3955-62 | 10.1 | 123 |
| 20 | Co-planar 3,3,4,4,5-pentachlorinated biphenyl and non-co-planar 2,2,4,6,6-pentachlorinated biphenyl differentially induce recruitment of oestrogen receptor alpha to aryl hydrocarbon receptor target genes. <i>Biochemical Journal</i> , 2007 , 406, 343-53 | 3.8 | 41 |
| 19 | Estrogen receptor (ER) beta modulates ERalpha-mediated transcriptional activation by altering the recruitment of c-Fos and c-Jun to estrogen-responsive promoters. <i>Molecular Endocrinology</i> , 2006 , 20, 534-43 | | 147 |
| 18 | Estrogen receptor and aryl hydrocarbon receptor signaling pathways. <i>Nuclear Receptor Signaling</i> , 2006 , 4, e016 | 1 | 178 |
| 17 | EID3 is a novel EID family member and an inhibitor of CBP-dependent co-activation. <i>Nucleic Acids Research</i> , 2005 , 33, 3561-9 | 20.1 | 40 |
| 16 | Estrogen receptor-alpha regulates SOCS-3 expression in human breast cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 335, 168-74 | 3.4 | 30 |
| 15 | Aryl hydrocarbon receptor-mediated transcription: ligand-dependent recruitment of estrogen receptor alpha to 2,3,7,8-tetrachlorodibenzo-p-dioxin-responsive promoters. <i>Molecular and Cellular Biology</i> , 2005 , 25, 5317-28 | 4.8 | 166 |
| 14 | Deoxyribonucleic acid response element-dependent regulation of transcription by orphan nuclear receptor estrogen receptor-related receptor gamma. <i>Molecular Endocrinology</i> , 2004 , 18, 312-25 | | 37 |
| 13 | The human RAP250 gene: genomic structure and promoter analysis. <i>Gene</i> , 2004 , 327, 233-8 | 3.8 | 4 |
| 12 | The Ah receptor inhibits estrogen-induced estrogen receptor beta in breast cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 320, 76-82 | 3.4 | 21 |
| 11 | The basic helix-loop-helix-PAS protein ARNT functions as a potent coactivator of estrogen receptor-dependent transcription. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 6517-22 | 11.5 | 119 |
| 10 | Estrogen signaling: a subtle balance between ER alpha and ER beta. <i>Molecular Interventions: Pharmacological Perspectives From Biology, Chemistry and Genomics</i> , 2003 , 3, 281-92 | | 625 |
| 9 | Distribution of androgen receptor mRNA expression and immunoreactivity in the brain of the green anole lizard. <i>Journal of Neuroendocrinology</i> , 2002 , 14, 19-28 | 3.8 | 56 |

| | | | |
|---|--|-----|-----|
| 8 | In silico approaches to mechanistic and predictive toxicology: an introduction to bioinformatics for toxicologists. <i>Critical Reviews in Toxicology</i> , 2002 , 32, 67-112 | 5.7 | 26 |
| 7 | Ability of structurally diverse natural products and synthetic chemicals to induce gene expression mediated by estrogen receptors from various species. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2002 , 82, 181-94 | 5.1 | 30 |
| 6 | Hydroxylated benzo[a]pyrene metabolites are responsible for in vitro estrogen receptor-mediated gene expression induced by benzo[a]pyrene, but do not elicit uterotrophic effects in vivo. <i>Toxicological Sciences</i> , 2001 , 59, 231-40 | 4.4 | 63 |
| 5 | Interaction of PAH-related compounds with the alpha and beta isoforms of the estrogen receptor. <i>Toxicology Letters</i> , 2001 , 121, 167-77 | 4.4 | 104 |
| 4 | Reciprocal mutagenesis between human alpha(L349, M528) and rainbow trout (M317, I496) estrogen receptor residues demonstrates their importance in ligand binding and gene expression at different temperatures. <i>Molecular and Cellular Endocrinology</i> , 2001 , 183, 127-39 | 4.4 | 17 |
| 3 | In vitro and in vivo interactions of bisphenol A and its metabolite, bisphenol A glucuronide, with estrogen receptors alpha and beta. <i>Chemical Research in Toxicology</i> , 2001 , 14, 149-57 | 4 | 369 |
| 2 | Differential estrogen receptor binding of estrogenic substances: a species comparison. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2000 , 74, 223-34 | 5.1 | 248 |
| 1 | Quantification of rainbow trout (<i>Oncorhynchus mykiss</i>) zona radiata and vitellogenin mRNA levels using real-time PCR after in vivo treatment with estradiol-17 beta or alpha-zearalenol. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2000 , 75, 109-19 | 5.1 | 94 |