

Carsten F Dormann

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7101978/carsten-f-dormann-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

146
papers

19,581
citations

52
h-index

139
g-index

164
ext. papers

24,086
ext. citations

6.1
avg, IF

6.76
L-index

#	Paper	IF	Citations
146	Quantitative Prediction of Interactions in Bipartite Networks Based on Traits, Abundances, and Phylogeny.. <i>American Naturalist</i> , 2022 , 199, 841-854	3.7	1
145	Population density estimates for terrestrial mammal species. <i>Global Ecology and Biogeography</i> , 2022 , 31, 978-994	6.1	0
144	Reply to: "Research on agroforestry systems and biodiversity conservation: what can we conclude so far and what should we improve?" by Boinot et al. 2022.. <i>Bmc Ecology and Evolution</i> , 2022 , 22, 65	2.1	
143	European agroforestry has no unequivocal effect on biodiversity: a time-cumulative meta-analysis. <i>Bmc Ecology and Evolution</i> , 2021 , 21, 193	2.1	3
142	Increasing connectivity enhances habitat specialists but simplifies plant-insect food webs. <i>Oecologia</i> , 2021 , 195, 539-546	2.9	5
141	Tree diversity reduces the risk of bark beetle infestation for preferred conifer species, but increases the risk for less preferred hosts. <i>Journal of Ecology</i> , 2021 , 109, 2649-2661	6	4
140	Curvature of LogsDevelopment of and Comparison between Different Calculation Approaches. <i>Forests</i> , 2021 , 12, 857	2.8	
139	Within-day dynamics of plant-pollinator networks are dominated by early flower closure: an experimental test of network plasticity. <i>Oecologia</i> , 2021 , 196, 781-794	2.9	2
138	Seeing through the static: the temporal dimension of plant-animal mutualistic interactions. <i>Ecology Letters</i> , 2021 , 24, 149-161	10	24
137	Temporal scale-dependence of plant-pollinator networks. <i>Oikos</i> , 2020 , 129, 1289-1302	4	25
136	A standard protocol for reporting species distribution models. <i>Ecography</i> , 2020 , 43, 1261-1277	6.5	141
135	Spatial conservation prioritisation in data-poor countries: a quantitative sensitivity analysis using multiple taxa. <i>BMC Ecology</i> , 2020 , 20, 35	2.7	3
134	Co-occurrence patterns and the large-scale spatial structure of benthic communities in seagrass meadows and bare sand. <i>BMC Ecology</i> , 2020 , 20, 37	2.7	2
133	The influence of camera trap flash type on the behavioural reactions and trapping rates of red deer and roe deer. <i>Remote Sensing in Ecology and Conservation</i> , 2020 , 6, 399-410	5.3	3
132	Evaluating the effectiveness of retention forestry to enhance biodiversity in production forests of Central Europe using an interdisciplinary, multi-scale approach. <i>Ecology and Evolution</i> , 2020 , 10, 1489-1509	2.8	27
131	Calibration of probability predictions from machine-learning and statistical models. <i>Global Ecology and Biogeography</i> , 2020 , 29, 760-765	6.1	9
130	Correlation and Association 2020 , 65-70		

129 Multiple Regression: Regression with Multiple Predictors **2020**, 207-226

128 Samples, Random Variables Histograms, Density Distribution **2020**, 1-12

127 Multiple Regression in R **2020**, 227-255

126 Evidence Ranking Needs to Reflect Causality. *Trends in Ecology and Evolution*, **2020**, 35, 94-95 10.9 1

125 Humpback whales extend their stay in a breeding ground in the Tropical Eastern Pacific. *ICES Journal of Marine Science*, **2020**, 77, 109-118 2.7 11

124 Insect abundance in managed forests benefits from multi-layered vegetation. *Basic and Applied Ecology*, **2020**, 48, 124-135 3.2 8

123 Plant species richness increases with light availability, but not variability, in temperate forests understorey. *BMC Ecology*, **2020**, 20, 43 2.7 17

122 Exploration of Concerns about the Evidence-Based Guideline Approach in Conservation Management: Hints from Medical Practice. *Environmental Management*, **2020**, 66, 435-449 3.1 4

121 Spatial validation reveals poor predictive performance of large-scale ecological mapping models. *Nature Communications*, **2020**, 11, 4540 17.4 92

120 Breaking the ecosystem services glass ceiling: realising impact. *Regional Environmental Change*, **2019**, 19, 2261-2274 4.3 4

119 Spatial behavior in rehabilitated orangutans in Sumatra: Where do they go?. *PLoS ONE*, **2019**, 14, e0215284 3.4 0

118 A new model explaining the origin of different topologies in interaction networks. *Ecology*, **2019**, 100, e02796 4.6 14

117 African elephant poaching rates correlate with local poverty, national corruption and global ivory price. *Nature Communications*, **2019**, 10, 2242 17.4 39

116 Refuges from fire maintain pollinator-plant interaction networks. *Ecology and Evolution*, **2019**, 9, 5777-5785 7.8 11

115 Better Model Transfers Require Knowledge of Mechanisms. *Trends in Ecology and Evolution*, **2019**, 34, 489-490 10.9 8

114 The interplay of landscape composition and configuration: new pathways to manage functional biodiversity and agroecosystem services across Europe. *Ecology Letters*, **2019**, 22, 1083-1094 10 171

113 Does Public Participation Shift German National Park Priorities Away from Nature Conservation?. *Environmental Conservation*, **2019**, 46, 84-91 3.3 5

112 Blind spots in ecosystem services research and challenges for implementation. *Regional Environmental Change*, **2019**, 19, 2151-2172 4.3 49

111	Standards for distribution models in biodiversity assessments. <i>Science Advances</i> , 2019 , 5, eaat4858	14.3	309
110	Forest-edge associated bees benefit from the proportion of tropical forest regardless of its edge length. <i>Biological Conservation</i> , 2018 , 220, 149-160	6.2	21
109	Current global risks to marine mammals: Taking stock of the threats. <i>Biological Conservation</i> , 2018 , 221, 44-58	6.2	93
108	Consistent set of additive biomass functions for eight tree species in Germany fit by nonlinear seemingly unrelated regression. <i>Annals of Forest Science</i> , 2018 , 75, 1	3.1	9
107	Disturbance intensity is a stronger driver of biomass recovery than remaining tree-community attributes in a managed Amazonian forest. <i>Journal of Applied Ecology</i> , 2018 , 55, 1647-1657	5.8	23
106	Wrong, but useful: regional species distribution models may not be improved by range-wide data under biased sampling. <i>Ecology and Evolution</i> , 2018 , 8, 2196-2206	2.8	37
105	Modelling the variation of bark thickness within and between European silver fir (<i>Abies alba</i> Mill.) trees in southwest Germany. <i>Forestry</i> , 2018 , 91, 283-294	2.2	7
104	Model averaging in ecology: a review of Bayesian, information-theoretic, and tactical approaches for predictive inference. <i>Ecological Monographs</i> , 2018 , 88, 485-504	9	105
103	Computing AIC for black-box models using generalized degrees of freedom: A comparison with cross-validation. <i>Communications in Statistics Part B: Simulation and Computation</i> , 2018 , 47, 1382-1396	0.6	10
102	Crop pests and predators exhibit inconsistent responses to surrounding landscape composition. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, E7863-E7870	11.5	265
101	Biotic interactions in species distribution modelling: 10 questions to guide interpretation and avoid false conclusions. <i>Global Ecology and Biogeography</i> , 2018 , 27, 1004-1016	6.1	111
100	Decaying trees improve nesting opportunities for cavity-nesting birds in temperate and boreal forests: A meta-analysis and implications for retention forestry. <i>Ecology and Evolution</i> , 2018 , 8, 8616-8626	2.8	24
99	Outstanding Challenges in the Transferability of Ecological Models. <i>Trends in Ecology and Evolution</i> , 2018 , 33, 790-802	10.9	213
98	Fragmentation of nest and foraging habitat affects time budgets of solitary bees, their fitness and pollination services, depending on traits: Results from an individual-based model. <i>PLoS ONE</i> , 2018 , 13, e0188269	3.7	32
97	An efficient method to exploit LiDAR data in animal ecology. <i>Methods in Ecology and Evolution</i> , 2018 , 9, 893-904	7.7	13
96	Quantifying forest structural diversity based on large-scale inventory data: a new approach to support biodiversity monitoring. <i>Forest Ecosystems</i> , 2018 , 5,	3.8	31
95	Improved species-occurrence predictions in data-poor regions: using large-scale data and bias correction with down-weighted Poisson regression and Maxent. <i>Ecography</i> , 2018 , 41, 1161-1172	6.5	36
94	Wrong, but useful: regional species distribution models may not be improved by range-wide data under biased sampling 2018 , 8, 2196		1

93	Comparison of models for estimating bark thickness of <i>Picea abies</i> in southwest Germany: the role of tree, stand, and environmental factors. <i>Annals of Forest Science</i> , 2017 , 74, 1	3.1	14
92	Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. <i>Ecography</i> , 2017 , 40, 913-929	6.5	566
91	Recruitment, growth and recovery of commercial tree species over 30 years following logging and thinning in a tropical rain forest. <i>Forest Ecology and Management</i> , 2017 , 385, 225-235	3.9	43
90	Parametrische Statistik 2017 ,		2
89	Identifying Causes of Patterns in Ecological Networks: Opportunities and Limitations. <i>Annual Review of Ecology, Evolution, and Systematics</i> , 2017 , 48, 559-584	13.5	96
88	No consistent effect of plant species richness on resistance to simulated climate change for above- or below-ground processes in managed grasslands. <i>BMC Ecology</i> , 2017 , 17, 23	2.7	5
87	Influence of Forest Harvest on Nitrate Concentration in Temperate Streams A Meta-Analysis. <i>Forests</i> , 2017 , 8, 5	2.8	13
86	The former Iron Curtain still drives biodiversity-profit trade-offs in German agriculture. <i>Nature Ecology and Evolution</i> , 2017 , 1, 1279-1284	12.3	76
85	The influence of floral traits on specialization and modularity of plant-pollinator networks in a biodiversity hotspot in the Peruvian Andes. <i>Annals of Botany</i> , 2016 , 118, 415-29	4.1	44
84	Effectiveness of light-reflecting devices: A systematic reanalysis of animal-vehicle collision data. <i>Accident Analysis and Prevention</i> , 2016 , 97, 242-260	6.1	16
83	Environmental management: Synthesize evidence to steer decisions. <i>Nature</i> , 2016 , 529, 466	50.4	4
82	Measurement and prediction of bark thickness in <i>Picea abies</i> : assessment of accuracy, precision, and sample size requirements. <i>Canadian Journal of Forest Research</i> , 2016 , 46, 39-47	1.9	16
81	Dispersal Ecology Informs Design of Large-Scale Wildlife Corridors. <i>PLoS ONE</i> , 2016 , 11, e0162989	3.7	15
80	An evidence assessment tool for ecosystem services and conservation studies 2016 , 26, 1295-1301		36
79	Ecological networks are more sensitive to plant than to animal extinction under climate change. <i>Nature Communications</i> , 2016 , 7, 13965	17.4	118
78	Detection probabilities for sessile organisms. <i>Ecosphere</i> , 2016 , 7, e01546	3.1	9
77	Squares of different sizes: effect of geographical projection on model parameter estimates in species distribution modeling. <i>Ecology and Evolution</i> , 2016 , 6, 202-11	2.8	9
76	Effects of warming and drought on potential N ₂ O emissions and denitrifying bacteria abundance in grasslands with different land-use. <i>FEMS Microbiology Ecology</i> , 2015 , 91,	4.3	27

75	Community structure and ecological specialization in plant–ant interactions. <i>Journal of Tropical Ecology</i> , 2015 , 31, 325-334	1.3	4
74	Medium-term dynamics of tree species composition in response to silvicultural intervention intensities in a tropical rain forest. <i>Biological Conservation</i> , 2015 , 191, 577-586	6.2	39
73	Cross-Scale Variation in Biodiversity-Environment Links Illustrated by Coastal Sandflat Communities. <i>PLoS ONE</i> , 2015 , 10, e0142411	3.7	12
72	Ecological, historical and evolutionary determinants of modularity in weighted seed-dispersal networks. <i>Ecology Letters</i> , 2014 , 17, 454-63	10	125
71	Choices of abundance currency, community definition and diversity metric control the predictive power of macroecological models of biodiversity. <i>Global Ecology and Biogeography</i> , 2014 , 23, 468-478	6.1	8
70	Accounting for geographical variation in species–area relationships improves the prediction of plant species richness at the global scale. <i>Journal of Biogeography</i> , 2014 , 41, 261-273	4.1	35
69	EDITOR'S CHOICE: REVIEW: Effects of land use on plant diversity – A global meta-analysis. <i>Journal of Applied Ecology</i> , 2014 , 51, 1690-1700	5.8	72
68	A method for detecting modules in quantitative bipartite networks. <i>Methods in Ecology and Evolution</i> , 2014 , 5, 90-98	7.7	257
67	Interannual variation in land-use intensity enhances grassland multidiversity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 308-13	11.5	166
66	The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility and relatedness. <i>Ecology Letters</i> , 2014 , 17, 1389-99	10	112
65	The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. <i>Ecology and Evolution</i> , 2014 , 4, 4701-35	2.8	132
64	Stacking species distribution models and adjusting bias by linking them to macroecological models. <i>Global Ecology and Biogeography</i> , 2014 , 23, 99-112	6.1	196
63	Mass-flowering crops enhance wild bee abundance. <i>Oecologia</i> , 2013 , 172, 477-84	2.9	138
62	Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. <i>Ecography</i> , 2013 , 36, 27-46	6.5	4125
61	Temporal variability of ecological niches: a study on intertidal macrobenthic fauna. <i>Oikos</i> , 2013 , 122, 754-760	4	10
60	Bee diversity effects on pollination depend on functional complementarity and niche shifts. <i>Ecology</i> , 2013 , 94, 2042-54	4.6	182
59	The role of biotic interactions in shaping distributions and realised assemblages of species: implications for species distribution modelling. <i>Biological Reviews</i> , 2013 , 88, 15-30	13.5	931
58	Process, correlation and parameter fitting in species distribution models: a response to Kriticos et al. <i>Journal of Biogeography</i> , 2013 , 40, 612-613	4.1	6

57	Does model-free forecasting really outperform the true model?. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, E3975	11.5	13
56	Organic farming favours insect-pollinated over non-insect pollinated forbs in meadows and wheat fields. <i>PLoS ONE</i> , 2013 , 8, e54818	3.7	21
55	Parametrische Statistik 2013 ,		7
54	A quantitative index of land-use intensity in grasslands: Integrating mowing, grazing and fertilization. <i>Basic and Applied Ecology</i> , 2012 , 13, 207-220	3.2	240
53	Correlation and process in species distribution models: bridging a dichotomy. <i>Journal of Biogeography</i> , 2012 , 39, 2119-2131	4.1	414
52	Towards novel approaches to modelling biotic interactions in multispecies assemblages at large spatial extents. <i>Journal of Biogeography</i> , 2012 , 39, 2163-2178	4.1	282
51	Less than eight (and a half) misconceptions of spatial analysis. <i>Journal of Biogeography</i> , 2012 , 39, 995-998.	4.1	57
50	What's on the horizon for macroecology?. <i>Ecography</i> , 2012 , 35, 673-683	6.5	129
49	Landscape moderation of biodiversity patterns and processes - eight hypotheses. <i>Biological Reviews</i> , 2012 , 87, 661-85	13.5	1121
48	Mapping water quality-related ecosystem services: concepts and applications for nitrogen retention and pesticide risk reduction. <i>International Journal of Biodiversity Science, Ecosystem Services & Management</i> , 2012 , 8, 35-49		17
47	Climatic niche evolution and species diversification in the Cape flora, South Africa. <i>Journal of Biogeography</i> , 2012 , 39, 2201-2211	4.1	52
46	Specialization of mutualistic interaction networks decreases toward tropical latitudes. <i>Current Biology</i> , 2012 , 22, 1925-31	6.3	223
45	Spatial and temporal trends of global pollination benefit. <i>PLoS ONE</i> , 2012 , 7, e35954	3.7	208
44	Spillover of functionally important organisms between managed and natural habitats. <i>Agriculture, Ecosystems and Environment</i> , 2012 , 146, 34-43	5.7	298
43	The responses of grassland plants to experimentally simulated climate change depend on land use and region. <i>Global Change Biology</i> , 2012 , 18, 127-137	11.4	35
42	Mind the gap! How well does Natura 2000 cover species of European interest?. <i>Nature Conservation</i> , 2012 , 3, 45-62		52
41	Predator richness increases the effect of prey diversity on prey yield. <i>Nature Communications</i> , 2012 , 3, 1305	17.4	81
40	Modelling Species Distributions 2011 , 179-196		9

39	Landscape elements as potential barriers and corridors for bees, wasps and parasitoids. <i>Biological Conservation</i> , 2011 , 144, 1816-1825	6.2	82
38	Microsite conditions dominate habitat selection of the red mason bee (<i>Osmia bicornis</i> , Hymenoptera: Megachilidae) in an urban environment: A case study from Leipzig, Germany. <i>Landscape and Urban Planning</i> , 2011 , 103, 15-23	7.7	37
37	Linné's floral clock is slow without pollinators--flower closure and plant-pollinator interaction webs. <i>Ecology Letters</i> , 2011 , 14, 896-904	10	38
36	A quantitative review of ecosystem service studies: approaches, shortcomings and the road ahead. <i>Journal of Applied Ecology</i> , 2011 , 48, 630-636	5.8	637
35	Species abundance distributions and richness estimations in fungal metagenomics--lessons learned from community ecology. <i>Molecular Ecology</i> , 2011 , 20, 275-85	5.7	134
34	Set-aside management: How do succession, sowing patterns and landscape context affect biodiversity?. <i>Agriculture, Ecosystems and Environment</i> , 2011 , 143, 37-44	5.7	80
33	Crop-noncrop spillover: arable fields affect trophic interactions on wild plants in surrounding habitats. <i>Oecologia</i> , 2011 , 166, 433-41	2.9	29
32	On managing the red mason bee (<i>Osmia bicornis</i>) in apple orchards. <i>Apidologie</i> , 2011 , 42, 564-576	2.3	49
31	Food web structure and biocontrol in a four-trophic level system across a landscape complexity gradient. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2011 , 278, 2946-53	4.4	104
30	Expansion of mass-flowering crops leads to transient pollinator dilution and reduced wild plant pollination. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2011 , 278, 3444-51	4.4	154
29	TaqMan real-time PCR assays to assess arbuscular mycorrhizal responses to field manipulation of grassland biodiversity: effects of soil characteristics, plant species richness, and functional traits. <i>Applied and Environmental Microbiology</i> , 2010 , 76, 3765-75	4.8	62
28	Evolution of climate niches in European mammals?. <i>Biology Letters</i> , 2010 , 6, 229-32	3.6	50
27	Reassessing Neotropical angiosperm distribution patterns based on monographic data: a geometric interpolation approach. <i>Biodiversity and Conservation</i> , 2010 , 19, 1523-1546	3.4	12
26	Ecological networks--beyond food webs. <i>Journal of Animal Ecology</i> , 2009 , 78, 253-69	4.7	615
25	Static species distribution models in dynamically changing systems: how good can predictions really be?. <i>Ecography</i> , 2009 , 32, 733-744	6.5	100
24	Response to Comment on "Methods to account for spatial autocorrelation in the analysis of species distributional data: a review" <i>Ecography</i> , 2009 , 32, 379-381	6.5	26
23	Indices, Graphs and Null Models: Analyzing Bipartite Ecological Networks. <i>Open Ecology Journal</i> , 2009 , 2, 7-24	2	830
22	Prediction uncertainty of environmental change effects on temperate European biodiversity. <i>Ecology Letters</i> , 2008 , 11, 235-44	10	70

21	Components of uncertainty in species distribution analysis: a case study of the Great Grey Shrike. <i>Ecology</i> , 2008 , 89, 3371-86	4.6	153
20	Application of species richness estimators for the assessment of fungal diversity. <i>FEMS Microbiology Letters</i> , 2008 , 282, 205-13	2.9	56
19	Effects of incorporating spatial autocorrelation into the analysis of species distribution data. <i>Global Ecology and Biogeography</i> , 2007 , 16, 129-138	6.1	425
18	Effects of landscape structure and land-use intensity on similarity of plant and animal communities. <i>Global Ecology and Biogeography</i> , 2007 , 16, 774-787	6.1	133
17	Promising the future? Global change projections of species distributions. <i>Basic and Applied Ecology</i> , 2007 , 8, 387-397	3.2	332
16	Assessing the validity of autologistic regression. <i>Ecological Modelling</i> , 2007 , 207, 234-242	3	85
15	Competition hierarchy, transitivity and additivity: investigating the effect of fertilisation on plant-plant interactions using three common bryophytes. <i>Plant Ecology</i> , 2007 , 191, 171-184	1.7	10
14	Methods to account for spatial autocorrelation in the analysis of species distributional data: a review. <i>Ecography</i> , 2007 , 30, 609-628	6.5	2078
13	Occurrence pattern of <i>Pararge aegeria</i> (Lepidoptera: Nymphalidae) with respect to local habitat suitability, climate and landscape structure. <i>Landscape Ecology</i> , 2006 , 21, 989-1001	4.3	17
12	Experimental evidence rejects pairwise modelling approach to coexistence in plant communities. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2005 , 272, 1279-85	4.4	45
11	Neighbour identity modifies effects of elevated temperature on plant performance in the High Arctic. <i>Global Change Biology</i> , 2004 , 10, 1587-1598	11.4	32
10	Consequences of manipulations in carbon and nitrogen supply for concentration of anti-herbivore defence compounds in <i>Salix polaris</i> . <i>Ecoscience</i> , 2003 , 10, 312-318	1.1	25
9	No evidence for adaptation of two <i>Polygonum viviparum</i> morphotypes of different bulbil characteristics to length of growing season: abundance, biomass and germination. <i>Polar Biology</i> , 2002 , 25, 884-890	2	12
8	Flowering, growth and defence in the two sexes: consequences of herbivore exclusion for <i>Salix polaris</i> . <i>Functional Ecology</i> , 2002 , 16, 649-656	5.6	21
7	Facilitation and competition in the high Arctic: the importance of the experimental approach. <i>Acta Oecologica</i> , 2002 , 23, 297-301	1.7	46
6	Competition and herbivory during salt marsh succession: the importance of forb growth strategy. <i>Journal of Ecology</i> , 2000 , 88, 571-583	6	50
5	Food Webs versus Interaction Networks: Principles, Pitfalls, and Perspectives		9-18
4	A new model explaining the origin of different topologies in interaction networks		1

3	European agroforestry is no universal remedy for biodiversity: a time-cumulative meta-analysis	2
2	Neither global nor consistent: a technical comment on the tree diversity-productivity analysis of Liang et al. (2016)	4
1	A systematic map of demographic data from elephant populations throughout Africa: implications for poaching and population analyses. <i>Mammal Review</i> ,	5